



Bio Ecology of Aquatic and Semi-Aquatic Insects Of Order Coleoptera In The World

Khaghani Ramin^{1,2}, Vatandoost Hassan^{1,3*}, Tashakori Ghazal¹, Jalilian Ali¹

¹Department of Medical Entomology & Vector Control, School of Public Health, Tehran University of Medical Sciences, Iran

²Department of Medical Parasitology, Faculty of Medicine, AJA University of Medical Sciences, Tehran, Iran

³Department of Chemical Pollutants and Pesticides, Institute for Environmental Research, Tehran University of Medical Sciences, Iran

*Corresponding author

Hassan Vatandoost, Department of Medical Entomology & Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Submitted: 26 Jul 2022; Accepted: 12 Sep 2022 Published: 21 Sep 2022

Citation: Khaghani, Ramin., Vatandoost, Hassan., Tashakori, Ghazal., Jalilian, Ali. (2022). Bio Ecology of Aquatic and Semi-Aquatic Insects Of Order Coleoptera In The World. *J Mari Scie Res Ocean*, 5(3),157-178.

Abstract

Background: Aquatic Coleoptera plays a major role in freshwater ecosystems and is regarded as an effective bioindicator. At least 23 beetle families, from three of the four extant suborders, are predominantly aquatic as adults, larvae, or both. They play important role in ecosystem functioning, nutrient cycling primary production, decomposition and materials translocation.

Methods: In this study, various databases including PubMed, Web of Knowledge, Scopus, Google Scholar and sciencedirect were used.

Results: Aquatic Coleoptera known as water beetles, with more than 13,000 described species, is one of the most abundant aquatic insects. Out of 4 suborders of Coleoptera, the suborder Myxophaga is truly aquatic whereas 8 of the 11 extant families of Adephaga are regarded as truly. As far as Polyphaga is concerned, the largest suborder of Coleoptera, only 13 of the 150 families are regarded as truly aquatic.

Discussion and Conclusion: Water beetles, although defined by their affinity for aquatic ways of life, occupy a broad array of habitats, and have shifted secondarily back to their terrestrial roots (either as adults, larvae or both) on multiple occasions. This ecological variability coupled with repeated, parallel transitions has positioned water beetles as a premier study group for questions related to dispersal, ecological speciation, and diversification rates. The characteristics of a bioindicator are: richness and diversity species, easy handling, ecological faithfulness, fragility to small environmental changes and good organism responses, and water beetles are very integral parts of any biotic component of any water bodies or wetlands. They are indicators of ecological diversity and habitat characteristics

Key words: Aquatic Coleoptera, Bioecology, Biodiversity, Habitat

Introduction

More species of insect have been described than of any other life form, and beetles are the greatest proportion of these. Beetles are an evolutionary success story par excellence, being by far the most species order of animals on earth. The Coleoptera are an old radiation, whose evolutionary origins may date back to the Permian or even Carboniferous [1, 2]. Water Beetles common name for any of numerous aquatic beetles including the true or diving water beetles, the whirligig beetles, and the water scavengers beetles. There are approximately 2000 species of true water beetles native to land

areas throughout the world [3]. Water beetles are very integral parts of the biotic component of any water bodies or wetlands. They are indicator of ecological diversity and habitat characteristics [4]. It has been recognized as a vital component for an integrated assessment of water quality. Use of aquatic insects as bio monitor provides important insights into changes in water and habitat quality [5]. These changes are valuable in demonstrating the effects of anthropogenic disturbances in the aquatic ecosystems. Numerous taxa of aquatic insects groups have been used as biological parameter to know the ecological status of aquatic systems. Of these

aquatic beetles provides all the requirements needed for the assessment that is, large diversity, various adaptations (tolerance and sensitivity), long life, and use of different types of habitats. All these qualities suit them perfectly to use as biological indicators [3]. Slipinski et al. estimated that there are about 386500 species belonging to 29500 genera and 176 families under 4 suborders (Archostemata, Myxophaga, Adephaga, and Polyphaga), representing 40% of the total animal diversity of the world [6]. Aquatic Coleoptera known as water beetles, with more than 13000 described species, is one of the most abundant aquatic insects [7]. They play an important role in freshwater ecosystems and are considered as a suitable bioindicator [8]. Grove & Stork (2000) hypothesized that about 70–95% of all the beetle species are yet to be discovered and described and also emphasized that it would take 200 years to explore the entire beetle fauna of the world [9]. The beetles are found everywhere and in almost all ecosystems where animals can thrive with the exceptions of arctic snow and the sea water. Aquatic beetles or water beetles live in almost all kinds of aquatic habitats, such as rivers, springs, lakes, ditches, puddles, seepages, and ground water [10]. However, beetles do not inhabit the oceans, although numerous species live at their shores, where they can be found in hypersaline rock pools of the supralittoral, i.e., the spray (or splash) zone slightly above the intertidal zone. In contrast to other insects, water beetles prefer small, richly vegetated ditches. In larger lakes, they prefer the swampy margins, as for instance the reed belt of the Central Europe, where water beetle biomass is probably higher than anywhere on earth [11]. Water beetle can live in the water as larvae and as adults, but the pupae are generally on land (Some, such as *Noterus* of the *Noteridae* family, pupate underwater in air-filled cocoons). Nearly all aquatic Coleoptera go to land to pupate, then return to the water as adults. Water beetles, together with some Diptera and Heteroptera, are among the few insects able to tolerate hypersaline waters, tolerating concentrations of more than 200 g/L of total solutes. This tolerance has developed independently and recurrently in several lineages, mostly Hydrophilidae (e.g., *Berosus*, *Enochrus*, *Paracymus*), *Hydraenidae* (*Ochthebius*), and *Dytiscidae* (some species of *Nebrioporus*, *Hygrotus*, and in the *Boreonectes* group of genera). Coastal, most commonly saltmarshes or rock pools, but they may also be found inland, with saline streams forming one of the most unusual aquatic environments [12]. Although comprehensive water beetle surveys are still lacking for large parts of the Neotropical and Afrotropical Realms, it is estimated that the Palearctic region contains 3350 described species out 3900 total estimated, the Neotropical region 2510 out of 3900, and the Afrotropical Region contains 2700 out of 3750 total estimation, followed by the Oriental 2200 out of 3580 and the Australian/Pacific Realm 1340 out of 2100 estimated species. Undoubtedly, the Nearctic 1420 out of 1550 species is by far the poorest region in terms of water beetle diversity. Within the Palearctic Region, the Mediterranean countries and Anatolia are to be regarded as biodiversity hot spots, at least for certain families. In the comparatively well-explored Oriental Region, Borneo was found to be a hot spot of paramount significance for many water beetle families [13]. Based on their association with aquatic life as of

adults and of immature stages, the aquatic beetles can be classified into following six groups: 1. True Water Beetles, 2. False Water Beetles, 3. Phytophilous Water Beetles, 4. Parasitic Water Beetles, 5. Facultative Water Beetles, and 6. Shore Beetles [14]. The adults of the true water beetles remain submerged for most of the time in water whereas their larvae and pupae may be aquatic or terrestrial. The species in this group are highly adapted to aquatic life showing important morphological adaptations such as swimming hairs on legs, divided eyes, plastron, large claws, and streamlined body form. In false water beetles, only the juvenile stages are aquatic, the adult stage is always predominantly terrestrial. Parasitic water beetles come in association with water only when their hosts are submerged whereas facultative water beetles and shore beetles are predominantly terrestrial beetle families with their juveniles live in very wet habitats [14].

Materials and Methods

In this study, various databases including PubMed, Web of Knowledge, Scopus, Google Scholar and science direct were used and information was extracted based on keywords: Aquatic Coleoptera, bioecology, biodiversity, Habitat and categorized and analyzed based on content.

Results

Out of 4 suborders of Coleoptera, the suborder Myxophaga is truly aquatic whereas 8 of the 11 extant families of Adephaga are regarded as truly aquatic (*Gyrinidae*, *Haliplidae*, *Meruidae*, *Noteridae*, *Amphizoidae*, *Aspidytidae*, *Hygrobiidae*, *Dytiscidae*). As far as Polyphaga is concerned, the largest suborder of Coleoptera, only 13 of the 150 families are regarded as truly aquatic (*Helophoridae*, *Epimetopidae*, *Hydrochidae*, *Spercheidae*, *Hydrophilidae*, *Hydraenidae*, *Scirtidae*, *Elmidae*, *Dryopidae*, *Lutrochidae*, *Psephenidae*, *Cneoglossidae*, and *Eulichadidae*) [15].

Suborder Adephaga

Adephaga includes 44990 species (18% aquatic) under 11 families globally [6, 15]. The families, *Gyrinidae*, *Haliplidae*, *Meruidae*, *Noteridae*, *Amphizoidae*, *Hygrobiidae*, and *Dytiscidae* are truly aquatic in nature

1. Family Gyrinidae (Whirligig Beetles)

Gyrinidae with its worldwide distribution consists of about 1000 species under 25 genera globally [6]. The beetles in this family show peculiar swimming behavior where adults rapidly revolve around a fixed point on the surface of the water. In static or moderately running water and preferably live in the habitats with rich oxygen contents. The beetles are commonly known as whirligig beetles and can be distinguished from the other Adephagan families by following characters: compound eyes divided completely, so placed with upper pair on the dorsal surface of the head, remains above the water line and the lower pair on the ventral surface of the head, remains below the water line when the beetle swims; antennae short with a broad, cup-shaped scape, sub triangular pedicel and elongate but compact flagellum, and meso and metathoracic legs broadly expanded and fringed with setae for swimming [16].

All stages except for the pupa are aquatic, with the adults spending the major part of their life on the water surface, being half submerged; the larvae, in contrast, are always fully submerged. Larvae creep about on the vegetation or bottom substrate. They can also swim by up- and downward undulation of the body. They often form huge aggregations of individuals. Interspecific swarms of up to eight species may occur [17]. In ponds and lakes, gyri-

nids are often found between stands of emergent plants such as reed, and also between Pandanus stems. Many species, especially in the tropics, are, however, rheobiotic and will be found in different kinds of streams, often hiding in undercut river banks or in crevices of submerged wood during daytime. Several species are attracted to light traps [18] (Figure.1).



Figure 1: Family of Gyrinidae beetle

2. Family Haliplidae (Crawling Water Beetles)

Haliplidae consists of 220 species under 5 genera: Algophilus, Apteraliplus, Brychius, Peltodytes, and Haliplus [19]. The genus Haliplus is widely distributed in Australian, Afrotropical, Nearctic, Neotropical, Oriental and Palaearctic regions. The family Haliplidae is a comparatively small group of inconspicuous, small water-dwelling beetles, which are commonly termed as crawling water beetles. The features that distinguish these beetles from other families are: extremely large meta-coxal plates which cover most of the abdominal ventrites, tarsal formula 5-5-5 and body form

adapted for aquatic life [20]. All stages except for the pupa are aquatic. Larvae cannot swim, rather they can be observed creeping over the substrate or on water plants. Adults move rather quickly in the water by alternately moving their hind legs. This produces the impression of water walking. Larvae do not need to access atmospheric air, they breathe via the integument and their gills [21]. Adults carry an air supply beneath the elytra and under the coxal plates; this supply has to be renewed at the surface from time to time (Figure.2).



Figure 2: Family of Haliplidae beetle

3. Family Meruidae

Meruidae is a family of aquatic beetles in the suborder Adephaga, with only one genus and species, *Meru phyllisae*. The term *meru* comes from the local indigenous population's (the Pemon people) word for "waterfall." Appropriately so, this beetle is only found in the natural rock water slide in El Tobogán, Las Amazonas, Vene-

zuela. These beetles are morphologically highly distinct and at a body length of 0.8 mm represent the perhaps smallest individuals of Adephaga. A survey of aquatic beetles of Venezuela indicated that *Meru* is most common during the wet season, when larger areas of granitic rock surface are covered with water film, which the adult beetles as well as the larvae inhabit [22] (Figure .3).



Figure 3: Family of Meruidae beetle

4. Family Noteridae (Burrowing Water Beetles)

Globally the family includes 258 species belonging to 16 genera under 3 subfamilies; Noterinae (240 species), Notomicrinae (11 species), and Phreatodytinae (7 species) [23]. The members of this family are called as burrowing water beetles. They are usually found in large numbers in sunny, and shallow lentic habitats, and generally avoid the situations where they need to burrow [24]. The larvae cannot swim, but crawl rapidly on the substrate or among

water plants, and larvae of some species (*Noterus*) are active burrowers [25]. Adults and larvae are aquatic; *Noterus* pupates under water in air-filled cocoons. Noteridae are commonly found in stagnant water between roots of water plants. Most species inhabit stagnant or slowly flowing water rich in emergent and floating vegetation. The dense network of roots latter are often inhabited by numerous noterids [26] (Figure.4) .



Figure 4: Family of Noteridae beetle

5. Family Amphizoidae (Trout Stream Beetle)

Monogeneric family with five described species, known only from North America and China. Larvae and adults of all species are aquatic living in rather fast flowing rivers. The members of this family lives in North America and China. Larvae and adults of all species are aquatic. Both larvae and adults live in fast-flowing mountain and foothill streams where they feed primarily on stonefly (Plecoptera) naiads. They may be abundant on driftwood

and floating trash or along undercut banks among roots and accumulations of Pine needles. Both larvae and adults live in fast flowing mountain and foothill streams where they feed primarily on stonefly (Plecoptera) naiads. Amphizoids live at the interface between aquatic and terrestrial environments and seem at home in both. Larvae remain just out of the water on twigs or other wood. They enter the water to seize prey, but quickly return to the twig to eat the victim [27] (Figure.5).



Figure 5: Family of Amphizoidae beetle

6. Family Hygrobiidae

Monogeneric family with six described species, occurring in Europe, China and Australia. Larvae and adults of all species are aquatic lives in stagnant water [28]. Their size is moderate (8.5-10

mm). They are most common in stagnant water, where they walk or swim; they do not dive. They obtain their oxygen from the air collected and stored under the elytra [29] (Figure.6).



Figure 6: Family Hygrobiidae beetle

7. Family Dytiscidae (Diving Beetles)

This family is worldwide in distribution with about 4,303 species belonging to 190 genera so far described globally [30]. The beetles in this family are adapted well to aquatic life. The adults and larvae live in water and can be found in any aquatic ecosystems. They mostly feed on aquatic invertebrates and fish eggs. This is why they are commonly called as 'predacious diving beetles'. Eggs,

larvae and adults are aquatic. However, adults of a few species from India, Australia and New Caledonia were described as terrestrial [18]. Adult Dytiscidae produce antimicrobial secretions with their pygidial glands. These secretions are applied to the body on land, where the beetle cleans and brushes his body with the legs [31] (Figure.7).



Figure 7: Family of Dytiscidae beetle

8. Family Aspidytidae (cliff water beetles)

Aspidytidae is a family of aquatic beetles of the suborder Adephaga, described in 2002 from specimens in South Africa and China [32]. There are only two known species in the family and these were originally described in the genus *Aspidytes* [32, 33]. But later the new genus *Sinaspidytes* was erected for the species found in

China [34]. Larvae and adults of both *A. niobe* and *S. wrasei* are found in or in close proximity of hygropetric habitats. Meaning they require a rock surface covered by a thin layer of water. Both adults and larvae are likely predacious. Eggs and pupa are yet unknown [33] (Figure.8).



Figure 8: Family Aspidytidae beetle

Suborder Myxophaga

Myxophaga includes 84 species under 4 families from the world: Lepiceridae (2 species), Torridincolidae (60 species), Hydroscaaphidae (22 species), and Sphaeriusidae (23 species) [6, 14, 15]. The two known species of Lepiceridae are reported from Mexico and Venezuela, whereas Torridincolidae occurs in South America, Africa and with few species from Asia [14].

1. Family Lepiceridae

Monogeneric family with two New World species. *Lepicerus inaequalis* and *L. bufo* both usually found on sand banks very close to streams but assumed they might get submerged under water [35]. In 2010 *Lepicerus pichilingue* new species (type locality: Quevedo, Los Ríos, Ecuador) is described from leaf litter in mixed plantings of plantain and cacao in western Ecuador. *L. pichilingue* is very similar to *L. inaequalis*, but differs distinctively in the structure of the aedeagus (Figure.9) [36].



Figure 9: Family Lepiceridae beetle

2. Family Torridincolidae

Torridincolidae has about 40 species distributed in seven genera, but only three are found on Neotropics. Torridincolidae are usually dark colored, some of which bearing iridescent scales, with smooth surface or finely sculptured with punctuation and carinae [37]. The species are found in mountain streams (Figure.10) [38].




Figure 10: Family of Torridincolidae beetle

3. Family Hydroscaphidae (Skiff Beetles)

The world fauna of this family includes about 22 species under 3 genera: Scaphydra (3 species), Yara (2 species), and Hydroscapha (17 species) globally [39]. The beetles in this family are small sized and commonly called as skiff beetles. They can be distinguished by the presence of a distinct notopleural suture and of their aquatic life [40]. Larvae, pupa and adults are aquatic, which is probably also true for the egg. They are found on algal mats, over

which a thin water film flows [41]. Adults are Hydroscapha able to move very quickly on water films; it seems, however, that they cannot swim freely. Because of its relatively large size, only one egg matures at a time. Larvae and adults feed on algae. Members of this family are found on hygropetric rocks and in the interstitial of gravelly river margins and residual pools. They are also reported from hot springs up to a temperature of 46°C [42] (Figure.11).



Figure 11: Family of Hydroscaphidae beetle

4. Family Sphaeriusidae

The beetles in this family are called as minute bog beetles which can be distinguished from other families of the suborder by their comparatively large and prominent head, capitate antennae, design of meso and metasternum, large posterior coxal plates and unequal length of visible abdominal sternites. They are found in mud, under

stones, or on algae along the edge of streams and rivers, among the roots of plants, in mosses associated bogs or inhabiting moist leaf litter further away from the bodies of water [40]. This monogeneric family is widely distributed, and includes 23 species belonging to the genus, *Sphaerius* from the world [15] (Figure.12).



Figure 12: Family Sphaeriusidae of beetle

Suborder Polyphaga

Only 13 of the 150 recognized families of the large suborder Polyphaga are regarded as predominantly aquatic.

1. Family Helophoridae

The Helophoridae is a relatively small hydrophiloid family with ca. 190 known species classified into a single genus, *Helophorus* Fabricius, 1775. Adults of the genus are easily recognizable by the elongate body and five longitudinal furrows on the pronotum.

The majority of species occur mostly in Palearctic and Nearctic regions and only a few species are known from the Afrotropical and Oriental regions [43, 44]. Adult beetles are saprophagous and usually inhabit shallow standing water bodies, only a few species are terrestrial. In contrast, larvae are terrestrial, living in moist habitats near water, preying on various small invertebrates. A few species have herbivorous larvae feeding on various Brassicaceae or Poaceae [45] (Figure.13).



Figure 13: Family of Helophoridae beetle

2. Family Epimetopidae

Epimetopidae are a small beetle family of the superfamily Hydrophiloidea, comprising 72 described species in three genera: the American *Epimetopus* Lacordaire, 1854 (56 species), Asian *Eumetopus* Balfour-Browne, 1949 (eight species) and African *Eupotemus* Ji & Jäch, 1998 (eight species, of which six are described as new here) [46]. Eggs, larvae and adults are aquatic. Eggs are carried by the female in an egg case underneath the abdomen (brood care); these cases contain up to 17 eggs as far as is known [47]. Besides that, hardly anything appears to be known about the biology

of Epimetopidae. Most species of Epimetopidae live in similar habitats: wet sand or gravel at sides of various types of streams and rivers, occasionally also with algal mats or accumulations of plant debris. The observation of living specimens of *Eumetopus acutumontis* from Vietnam indicate that the specimens were digging and hiding in the wet sandy or gravelly substrate at the margin of a small river. In contrast, some *Epimetopus* (e.g., *E. venezuelensis* Perkins, 2012) were collected in standing water, typically well vegetated shallow marshes [46] (Figure.14).





Figure 14: breeding places of Family of Epimetopidae beetle

3. Family Hydrochidae (water scavenger beetle)

Monogeneric family with about 180 species; hydrochids occur on all continents. All species are truly aquatic, living in well-vegetated stagnant water and/or at the edges of very slowly flowing water [48]. All inhabit well vegetated marginal lentic habitats where both

the larvae and adults are aquatic but do not swim; they crawl on vegetation where the adults are herbivores and detritivores. The larval feeding habits are not known. Adults are generally slow moving and cryptic and will often remain motionless for some time in the water net and so may be overlooked (Figure.15).



Figure 15: Family of Hydrochidae beetle

4. Family Spercheidae

Monogeneric family with 18 species. The genus occurs on all continents. Larvae and adults generally live in stagnant water [48]. Most of them from the Ethiopian and Oriental zoogeographic re-

gions [49]. They occur in stagnant water with rich vegetation, often in the mud on the edge of small ponds. Both adults and larvae are among the few coleopteran filter-feeders, feeding on detritus or possibly small crustaceans and insect larvae (Figure.16).



Figure 16: Family of Spercheidae beetle

5. Family Hydrophilidae (water scavenger beetle)

There are a total of 1740 species in 66 genera, on all continents, adults and larvae of most species are living in stagnant water, run-

ning water, in phytotelmata or seepages, numerous species are reportedly riparian or terrestrial (humicolous) [15] (Figure.17).




Figure 17: Family of Hydrophilidae beetle

6. Family Hydraenidae (Minute Moss Beetles)

Hydraenidae is a cosmopolitan family of tiny beetles, ranging in size from 0.5 to less than 4.0 mm in length. The majority of species are adapted for marginal life, mainly in the freshwater aquatic world [50, 51]. The family includes more than 1500 species, but several hundreds await formal description [52]. The family is currently divided into 4 subfamilies: Orchymontiinae Perkins, Prosthetopinae Perkins, Ochthebiinae Thomson, and Hydraeninae Mulsant [50-54]. Globally, Hydraenidae includes 1600 species under 42 genera, occurring on all continents. Eggs and adults of most species are aquatic; there is however a number of strictly terrestrial and riparian genera. Larvae appear to be terrestrial or riparian throughout, few of them (e.g. *Enicocerus*) seem to be amphibious

(facultatively aquatic); they usually inhabit damp sites close to the water (14, 55). Adult hydraenids cannot swim well. They crawl on the substrate and if one washes them loose, they float to the surface, where they usually cling upside down to the underside of the water surface trying to reach the next stone, leaf and the like to search for foothold and shelter. Certain adults, especially those living in well oxygenated streams use an incompressible gas gill (or microplastron), which enables them to stay permanently submerged. The aquatic species may be encountered in a variety of habitats: e.g. saline coastal rock pools, swamps, streams, seepage water. Often, numerous beetles can be encountered in leaf packs, between roots or any kind of wood in lentic and lotic situations [50, 56] (Figure.18).




Figure 18: Family of Hydraenidae beetle

7. Family Scirtidae (marsh beetles)

These beetles are commonly referred to as marsh beetles, as the larvae are typically associated with stagnant water, but can be found in flowing water. Adults prefer decomposing plant material near the water's edge [57]. About 900 species in 30 genera. Scirtids occur on all continents. Larvae are usually aquatic, although there

are reports about scirtid larvae found in wet soil and on rotten logs. Imagos are generally terrestrial, but adults of *Hydrocyphon* are occasionally collected under water; pupae of *Hydrocyphon* are also reported to be aquatic. Scirtid larvae are found in running water (about 20%), in stagnant water, phytotelmata, and in groundwater [57, 58] (Figure.19).



Figure 19: Family of Scirtidae beetle

8. Family Elmidae

The beetles in this family are commonly called as riffle beetles and they can be both aquatic and semi-aquatic in nature. Till date, 1498 species belonging to 147 genera under 2 subfamilies Elminae and Larinae have been described from the world [59]. The adults

and juvenile stages of the subfamily Elminae live totally aquatic habitats and on occasions leave water. The subfamily Larinae is centered in Afrotropical and Neotropical regions with lower diversity in the Oriental region [60] (Figure.20).






Figure 20: Family Elmidae beetle adults, habitus in dorsal view. (A) *Neocylloepus boeseli* Brown; *Neoelmis* sp.; (C) *Neolimnius palpalis* Hinton; (D) *Notelmis* sp.; (E) *Onychelmis* sp.; (F) *Pagelmis amazônica* Spangler; (G) *Pilielmis sara* Hinton; (H) *Portelmis nevermanni* Sanderson; (I) *Stethelmis kaszabi* Hinton; (J) *Stegoelmis* sp.; (K) *Stenhelmoides rufulus*-group; (L) *Tolmerelmis pubipes* (Hinton); (M) *Tyletelmis* sp.; (N) *Xenelmis* sp.

9. Family Dryopidae (Long-toed water beetle)

Dryopidae is an ecologically diverse family, with some members being aquatic, while others are completely terrestrial or semi-aquatic. Only five aquatic genera are recognized in the Neotropical Region. They are found on the sandy banks of lotic habitats in soft stream substrate, while others occur on emergent vegetation and sticks protruding from streams. They are often collected near lights at night; for these the habitat is unknown. In the terrestri-

al dryopids, adults and larvae can be found in forest leaf litter. This family contains about 300 species in 33 genera worldwide. Dryopids occur in all biogeographic regions, but they are absent from the Australian continent.. Larvae are generally riparian or terrestrial; adults of about 75% of the species are regarded as aquatic (lotic and lentic habitats), the remaining ones are riparian or terrestrial [61] (Figure.21).



Figure 21: Family of Dryopidae beetle

10. Family Lutrochidae (Travertine Beetles)

About 15 species, all confined to the New World. Larvae and adults are reported to be aquatic in lotic habitats (riparian gravel, emergent rocks or submerged wood) [62] (Figure.22).



Figure 22: Family of Lutrochidae beetle

11. Family Psephenidae (Water-penny beetle)

About 272 species in 35 genera. Psephenids occur on all continents. They are comprised of four subfamilies: Eubrianacinae, Eu-

briinae, Psepheninae, Psephenoidinae. Larvae are always aquatic, almost exclusively occurring in running water, with few exceptions, adults and pupae are strictly terrestrial [63] (Figure.23).





Figure 23: Family of Psephenidae beetle

12. Family Cneoglossidae

This family contains ten species in a single genus. Small beetles 3-6 mm long. Larvae of *Cneoglossa edsoni* were collected in submerged rotting brushwood in shallow flowing streams (Figure.24) [64]. The main species are : *Cneoglossa brevis* Champion, 1897,

Cneoglossa collaris Guérin-Méneville, 1849, *Cneoglossa elongata* Pic, 1916, *Cneoglossa gournellei* Pic, 1916, *Cneoglossa lampyroides* Champion, 1897, *Cneoglossa longipennis* (Pic, 1915), *Cneoglossa peruviana* Pic, 1916, *Cneoglossa rufifrons* Pic, 1916, *Cneoglossa testacericollis* Pic, 1916




Figure 24: Family of Cneoglossidae beetle

13. Family Eulichadidae (Forest Stream Beetles)

The small elateriform family Eulichadidae comprises of two extant genera: the monotypic Californian genus *Stenocolus* LeConte, 1853, and the predominately Oriental genus *Eulichas* Jacobson, 1913, with 30 described and more than 10 undescribed species [65, 66]. The larvae of both genera are aquatic, while the adults live on

vegetation near water, and especially the genus *Eulichas* is often attracted at light [66]. these five genera belong to the family Eulichadidae: *Cretasyne* Yan, Wang & Zhang, 2013, *Eulichas* Jacobson, 1913, *Mesaplus* Hong, 1983, *Mesodascilla* Martynov, 1926, *Stenocolus* LeConte, 1853. (Figure. 25)






Figure 25: Family of Eulichadidae beetle

Eulichas genus of family Eulichadidae . 1—*E. alesbezdeki* s, 2—*E. bertiae*, 3—*E. dudgeon*, 4—*E. jendeki* , 5—*E. siamensis* sp. , 6—*E. similis* sp., 7—*E. uniformis*, 8—*E. baeri*, 9—*E. baeri* , 10—*E. gigantea* , 11—*E. incisicollis* , 12—*E. incisicollis* , 13—*E. oborili* sp, 14—*E. robusta* , 15—*E. sausai* , 16—*E. serricornis* , 17—*E. sikkimensis* (, 18—*E. sundaensis* , 19—*E. sundaensis* , 20—*E. wewalkai* , 21—*E. fasciolata* , 22—*E. jakli* , 23—*E. subocellata* , 24—*E. villosa* , Gunung Emas).

Discussion

Water beetles, although defined by their affinity for aquatic ways of life, occupy a broad array of habitats, and have shifted secondarily back to their terrestrial roots (either as adults, larvae or both) on multiple occasions. This ecological variability coupled

with repeated, parallel transitions has positioned water beetles as a premier study group for questions related to dispersal, ecological speciation, and diversification rates [67, 68]. The development of comprehensive online specimen and fieldwork databases also has helped anchor our knowledge of water beetle. Many important surveys that focus in total or in part on water beetles have been carried out in the last quarter of a century. Ecologically, focusing on under sampled habitats such as seepages, underground waters and, to a lesser degree, the margins of river and streams, will likely yield new discoveries regardless of their geographical location [69]. Although regional research and generic revisions exist to a certain extent for some groups, the recent publication of comprehensive resources on aquatic coleoptera biology and identification put into sharp relief the varying degrees to which water

beetle identification is feasible. Of the 10 water beetle families with more than three described genera, modern global genus level keys now exist for just half: Dytiscidae, Noterida, Gyrinidae, Hydros-caphidae and Psephenidae [24, 70-71]. The diversity assessment and preparation of the water beetle inventories are considered an essential tasks now a day, due to the importance of wetlands in the of conservation planning and endeavours. Some species of water beetles have aquatic larvae and terrestrial adults. Water beetles from family Gyrinidae, Haliplidae, Noteridae, Amphizoidae, Dytiscidae and Hydros-caphidae are aquatic in all life stages. The

adult water beetles from family Hydros-caphidae, Hydrophilidae, Lutrochidae, Dyropidae, Elmidae, Eulichadidae, Heteroceridae, Limmichidae, Psephenidae, Ptilodactylidae and Sphaeriusidae are not aquatic. The characteristics of a bioindicator are: richness and diversity species, easy handling, ecological faithfulness, fragility to small environmental changes and good organism responses (Figure.26) and water beetles are very integral parts of any biotic component of any water bodies or wetlands. They are indicators of ecological diversity and habitat characteristics [72].




Figure 26: Important were beetles in the world

Conflict of Interest: The authors declare that there is no conflict of interest

Author Contributions: All the authors were involved

References

1. Toussaint, E. F., Seidel, M., Arriaga-Varela, E., Hájek, J. I. Ř. Í., Kral, D., Sekerka, L. U. K. A. Š., ... & Fikáček, M. A. R. T. I. N. (2017). The peril of dating beetles. *Systematic Entomology*, 42(1), 1-10.
2. Zhang, S. Q., Che, L. H., Li, Y., Liang, D., Pang, H., Ślipiński, A., & Zhang, P. (2018). Evolutionary history of Coleoptera revealed by extensive sampling of genes and species. *Nature communications*, 9(1), 1-11.
3. Sharma, S., Pandey, P., & Dave, V. (2013). Research article role of aquatic beetles for water quality assessment. *International Journal of Recent Scientific Research*, 4(11), 1673-1676.
4. Sánchez-Fernández, D., Abellán, P., Velasco, J., & Millán, A. (2004). Selecting areas to protect the biodiversity of aquatic ecosystems in a semiarid Mediterranean region using water beetles. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 14(5), 465-479.
5. Rosenberg, D. M. (1993). Introduction to freshwater biomonitoring and benthic macroinvertebrates. *Freshwater biomonitoring and benthic macroinvertebrates*, 1-9.
6. Zhang, Z. Q. (2011). Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Magnolia press*.
7. Short, A. E. Z. (2018). Systematics of aquatic beetles (Coleoptera): current state and future directions. *Systematic Entomology*, 43(1), 1-18.
8. Dong, B., Geng, C., Cai, Y., & Ji, L. (2014). Aquatic Coleoptera response to environmental factors of freshwater ecosystems in Changbai Mountain, northeast China. *Aquatic ecosystem health & management*, 17(2), 171-178.
9. Grove, S. J., & Stork, N. E. (2000). An inordinate fondness for beetles. *Invertebrate Systematics*, 14(6), 733-739.
10. Yule, C. M., & Sen, Y. H. (2004). Freshwater invertebrates of the Malaysian region. *Academy of Sciences Malaysia*.
11. Biswas, S., Mukhopadhyay, P., & Saha, S. K. (1995). Insecta: Coleoptera: Adephaga, Family Dytiscidae. *State fauna Series*, 5, 77-120.
12. Millán, A., Velasco, J., Gutiérrez-Cánovas, C., Arribas, P., Picazo, F., et al. (2011). Mediterranean saline streams in south-east Spain: What do we know? *J. Arid Environ*, 75, 1352-59.
13. Balian, E. V., Lévêque, C., Segers, H., Martens, K. (2008). Fresh water animal diversity assessment, *Hydrobiologia*, 595, 419-442.
14. Jäch, M. A. (1998). Annotated check list of aquatic and riparian/littoral beetle families of the world (Coleoptera). *Water beetles of China. Volume II.*, 25-42.
15. Jäch, M. A., & Balke, M. (2008). Global diversity of water beetles (Coleoptera) in freshwater. *Hydrobiologia*, 595(1), 419-442.
16. Thomas, M. C. (2000). *American Beetles, Volume I: Archostemata, Myxophaga, Adephaga, Polyphaga: Staphyliniformia*. CRC Press.
17. Stehr, F. W. (1991). *Immature insects volume 2*. Iowa, USA, Kendall/Hunt Publishing Company.
18. Yule, C. M., & Sen, Y. H. (2004). *Freshwater invertebrates of the Malaysian region*. Academy of Sciences Malaysia.
19. Nilsson, A. N., and Vondel, B. J. van. (2005a). *World Catalogue of Insects volume7, Amphizoidae, Aspidytidae, Haliplidae, Noteridae and Paelobiidae (Coleoptera, Adephaga)*, 171.
20. Thomas, M. C. (2000). *American Beetles, Volume I: Archostemata, Myxophaga, Adephaga, Polyphaga: Staphyliniformia*. CRC Press.
21. Spangler, P. J., & Steiner JR, W. E. (2005). A new aquatic beetle family, Meruidae, from Venezuela (Coleoptera: Adephaga). *Systematic Entomology*, 30(3), 339-357.
22. Nilsson, A. N. (2011). *A World Catalogue of the Family Noteridae*. Version 16, 54.
23. Miller, K. B. (2009). On the systematics of Noteridae (Coleoptera: Adephaga: Hydradephaga): Phylogeny, description of a new tribe, genus and species, and survey of female genital morphology. *Systematics and Biodiversity*, 7(2), 191-214.
24. Dettner K. (1997). Insecta: Coleoptera: Noteridae. In: Schwoerbel, J. & Zwick, P. (eds), *Süßwasserfauna von Mitteleuropa*. 20(2, 3 and 4), 98-126.
25. Sharma, S., Sharma, G., & Pir, F. A. (2019). Diversity and habitat selection of aquatic beetles (Coleoptera). *IOSR J Pharm Biol Sci*, 14, 31-37.
26. Nilsson, A. N. (2005). Family Amphizoidae (Coleoptera, Adephaga). In *Amphizoidae, Aspidytidae, Haliplidae, Noteridae and Paelobiidae (Coleoptera, Adephaga)* (pp. 15-18). Brill.
27. Nilsson, A. N. (2005). Family Paelobiidae (Coleoptera, Adephaga). In *Amphizoidae, Aspidytidae, Haliplidae, Noteridae and Paelobiidae (Coleoptera, Adephaga)* (pp. 154-163). Brill.
28. Glime, J. M. (2017). *Aquatic Insects: Holometabola–Diptera, Suborder Nematocera*. *Bryophyte Ecology. Volume 2. Bryological Interaction*, 1-22.
29. Nilsson, A. N., & Hájek, J. (2015). A world catalogue of the family Dytiscidae, or the diving beetles (Coleoptera, Adephaga). Version, 1, 2015.
30. Kovac, D., & Maschwitz, U. (1990). Secretion-grooming in aquatic beetles (Hydradephaga): a chemical protection against contamination of the hydrofuge respiratory region. *Chemoecology*, 1(3), 131-138.
31. Ribera, I., Beutel, R. G., Balke, M., & Vogler, A. P. (2002). Discovery of Aspidytidae, a new family of aquatic Coleoptera. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 269(1507), 2351-2356.
32. Beutel, R. G., Balke, M., & Ribera, I. (2011). 3.1. Aspidytidae Ribera, Beutel, Balke and Vogler, 2002. In *Volume 2 Morphology and Systematics (Elateroidea, Bostrichiformia, Cucujiformia partim)* (pp. 21-28). De Gruyter.
33. Toussaint, E. F., Beutel, R. G., Morinière, J., Jia, F., Xu, S., Michat, M. C., ... & Balke, M. (2016). Molecular phylogeny

- of the highly disjunct cliff water beetles from South Africa and China (Coleoptera: Aspidytidae). *Zoological Journal of the Linnean Society*, 176(3), 537-546.
34. Reichardt, H. (1976). Revision of the Lepiceridae (Coleoptera, Myxophaga).
35. Flowers, R. W., Shepard, W. D., & MERA, R. T. (2010). A new species of *Lepicerus* (Coleoptera: Lepiceridae) from Ecuador. *Zootaxa*, 2639(1), 35-39.
36. Sampaio, B. H., & Short, A. E. (2018). Families Hydroscaphidae and Torridincolidae. In Thorp and Covich's *Freshwater Invertebrates* (pp. 519-525). Academic Press.
37. Jäch, M. A. (1998). Torridincolidae: I. First record of Torridincolidae from China (Coleoptera). *Water beetles of China*, 2, 51-52.
38. Fikáček, M. A. R. T. I. N., & Šípková, H. A. N. A. (2009). New Asian Hydroscapha, with comments on male-female association of co-occurring species (Coleoptera, Myxophaga, Hydroscaphidae). *Zootaxa*, 2286(1), 31-48.
39. Thomas, M. C. (2000). *American Beetles, Volume I: Archostemata, Myxophaga, Adephaga, Polyphaga: Staphyliniformia*. CRC Press.
40. Vanin, S. A., Beutel, R. G., & Arce-Pérez, R. (2011). 6.4. Hydroscaphidae LeConte, 1874. Volume 1: Morphology and Systematics (Archostemata, Adephaga, Myxophaga, Polyphaga partim), 4, 49.
41. Merritt, R. W., & Cummins, K. W. (Eds.). (1996). *An introduction to the aquatic insects of North America*. Kendall Hunt.
42. Hansen, M. (1999). World catalogue of insects. Volume 2: Hydrophiloidea (s. str.)(Coleoptera). World catalogue of insects. Volume 2: Hydrophiloidea (s. str.)(Coleoptera).
43. Short, A. E. Z., & Hebauer, F. (2006). World catalogue of the Hydrophiloidea—additions and corrections, 1 (1999–2005). *Koleopterologische Rundschau*, 76, 315-359.
44. Angus, R. (1992). *Insecta: Coleoptera: Hydrophilidae: Helophorinae*. G. Fischer.
45. Fikáček, M., Matsumoto, K., Perkins, P., Prokin, A., Sazhnev, A., Litovkin, S., & Jaech, M. A. (2021). The family Epimetopidae (Coleoptera: Hydrophiloidea): review of current knowledge, genus-level phylogeny, and taxonomic revision of *Eupotemus*. *Acta Entomologica Musei Nationalis Pragae*, 61(1), 1-34.
46. Archangelsky, M. (1997). Studies on the biology, ecology, and systematics of the immature stages of New World Hydronhiloidea (Coleoptera: Staphyliniformia). *Bull. Ohio Biol. Surv.*, (n. ser.), 12, 1-207.
47. Darilmaz, M. C., & Kiyak, S. (2011). A study of the family Spercheidae (Coleoptera) from Turkey. *Turkish Journal of Zoology*, 35(3), 441-444.
48. Hansen, M. (1998). *Hydraenidae* (Coleoptera). In *Hydraenidae* (Coleoptera). Brill.
49. Perkins, P. D. (1997). Life on the effective bubble: exocrine secretion delivery systems (ESDS) and the evolution and classification of beetles in the family Hydraenidae (Insecta: Coleoptera). *Annals of Carnegie Museum*, 66(2), 89-207.
50. Jäch, M. A., & Díaz, J. A. (2001). New and little known Palearctic species of the genus *Hydraena* Kugelann VII (Insecta: Coleoptera: Hydraenidae). *Annalen des Naturhistorischen Museums in Wien. Serie B für Botanik und Zoologie*, 363-373.
51. Löbl, I., & Smetana, A. (2004). Catalogue of Palearctic Coleoptera. Volume 2: Hydrophiloidea-Histeroidea-Staphylininoidea. Catalogue of Palearctic Coleoptera. Volume 2: Hydrophiloidea-Histeroidea-Staphylininoidea.
52. Beutel, R. G., Anton, E., & Jäch, M. A. (2003). On the evolution of adult head structures and the phylogeny of Hydraenidae (Coleoptera, Staphyliniformia). *Journal of Zoological Systematics and Evolutionary Research*, 41(4), 256-275.
53. Hansen, M. (1997). Evolutionary trends in "staphyliniform" beetles (Coleoptera). *Steenstrupia*, 23, 43-86.
54. Jäch, M. A., Beutel, R. G., Díaz, J. A., & Kodada, J. (2000). Subgeneric classification, description of head structures, and world check list of *Hydraena* Kugelann (Insecta: Coleoptera: Hydraenidae). *Annalen des Naturhistorischen Museums in Wien. Serie B für Botanik und Zoologie*, 177-258.
55. Epler, J. H. (2010). *The Water Beetles of Florida—an identification manual for the families Chrysomelidae, Curculionidae, Dryopidae, Elmidae, Gyrinidae, Haliplidae, Helophoridae, Hydraenidae, Hydrochidae, Hydrophilidae, Noteridae, Psephenidae, Ptilodactylidae and Scirtidae*. Florida Department of Environmental Protection, Tallahassee, FL, 399.
56. Lawrence, J. F. (2005). Scirtidae Fleming, 1821. In Beutel, R. G. & R. A. B. Leschen (eds), *Handbook of Zoology, Vol. IV (Part 38), Coleoptera, Beetles, Vol. 1: Morphology and Systematics (Archostemata, Adephaga, Myxophaga, Polyphagapartim)*. Walter de Gruyter, Berlin, 443-450.
57. Jäch, M., Kodada, J., Brojer, M., Shepard, W. D., & Čiampor Jr, F. (2016). *Coleoptera: Elmidae and Protelmidae*. Brill.
58. Arnett Jr, R. H., Thomas, M. C., Skelley, P. E., & Frank, J. H. (2002). *American beetles: Vol 2. Polyphaga: Scarabaeoidea through Curculionoidea*. CRC Press LLC.
59. Kodada, J., & Jäch, M. A. (2005). Elmidae Curtis, 1830. In Beutel, R. G., & R. A. B. Leschen (eds), *Handbook of Zoology, Vol. IV (Part 38), Coleoptera, Beetles, Vol. 1 Morphology and Systematics (Archostemata, Adephaga, Myxophaga, Polyphagapartim)*. Walter de Gruyter, Berlin: 471-496.
60. Ide, S., Costa, C., & Vanin, S. A. (2005). Lutrochidae Kasap & Crowson, 1975. *Coleoptera, beetles. Vol. 1, Morphology and systematics (Archostemata, Adephaga, Myxophaga, Polyphaga partim)*.
61. Lee, C. F. (2005). Psephenidae Lacordaire, 1854. *Morphology and Systematics*, 521-533.
62. Costa, C., Vanin, S. A., & Ide, S. (1999). Systematics and biometrics of Cneoglossidae with a cladistic analysis of Byrrhoidea sensu Lawrence & Newton (1995)(Coleoptera, Elateriformia). *Arquivos de Zoologia*, 35(3), 231-300.
63. Jäch, M. A. (1995). Eulichadidae: Synopsis of the species of the genus *Eulichas* Jacobson from China, Laos and Vietnam (Coleoptera). *Water Beetles of China*, 1, 359-388.

-
64. HÁJEK, J. (2007). Revision of the genus *Eulichas* Jacobson, 1913 (Coleoptera: Eulichadidae) I. Introduction, morphology of adults, key to subgenera and species groups, and taxonomy of *E. funebris* species group. *Zootaxa*, 1620(1), 1-35.
 65. Ribera, I., Vogler, A. P., & Balke, M. (2008). Phylogeny and diversification of diving beetles (Coleoptera: Dytiscidae). *Cladistics*, 24(4), 563-590.
 66. Bloom, D. D., Fikáček, M., & Short, A. E. (2014). Clade age and diversification rate variation explain disparity in species richness among water scavenger beetle (Hydrophilidae) lineages. *Plos one*, 9(6), e98430.
 67. Short, A. E. Z. (2018). Systematics of aquatic beetles (Coleoptera): current state and future directions. *Systematic Entomology*, 43(1), 1-18.
 68. Miller K B & Bergsten J (2016). *Diving Beetles of the World*. Johns Hopkins University Press, Baltimore, Maryland, 336pp.
 69. Miller, K. B., & Bergsten, J. (2012). Phylogeny and classification of whirligig beetles (Coleoptera: Gyrinidae): relaxed-clock model outperforms parsimony and time-free Bayesian analyses. *Systematic Entomology*, 37(4), 706-746.
 70. Short, A. E. Z., Joly, L. J., Garcia, M., Wild, A. L., Bloom, D. D., & Maddison, D. R. (2015). Molecular phylogeny of the Hydroscaphidae (Coleoptera: M yxophaga) with description of a remarkable new lineage from the Guiana Shield. *Systematic Entomology*, 40(1), 214-229.
 71. LEE, C. F., Sato, M., Shepard, W. D., & Jaech, M. A. (2007). Phylogeny of Psephenidae (Coleoptera: Byrrhoidea) based on larval, pupal and adult characters. *Systematic Entomology*, 32(3), 502-538.
 72. Sharma, S., Pandey, P., Dave, V. (2013). Role of aquatic beetles for water quality assessment. *International Journal of Recent Scientific Research*. 4(11),1673-1676.

Copyright: ©2022 Hassan Vatandoost. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.