

Enriched Agriculture Horticulture Science Technology Socio-Economy-Communication-Issue by Biomedicines Suppressing Tomato-Disease and Coronavirus 2-Like-Pandemic-Diseases!

Subhas Chandra Datta^{1*}

¹Headmaster, Secretary and Researcher, Kanchannagar D.N.Das High School (HS), Kanchannagar, Burdwan Municipality, Purba Bardhaman, Bardhaman-713102, West Bengal, India

*Corresponding author

Dr. Subhas Chandra Datta, PhD in VisvaBharati, C/O- Rajendra Nath Nag, House No.-430A, Bajeprotappur (Katwa Road), Burdwan Municipality, Purba Bardhaman, Bardhaman-713101, West Bengal, India

Submitted: 30 Apr 2021; **Accepted:** 17 May 2021; **Published:** 20 May 2021

Citation: Dr. Subhas Chandra Datta (2021). Enriched Agriculture Horticulture Science Technology Socio-Economy-Communication-Issue by Biomedicines Suppressing Tomato-Disease and Coronavirus 2-Like-Pandemic-Diseases!. *J Agri Horti Res*, 4(2): 74-77.

Abstract

To avoid traditional control of root-knot disease, there remains a need for developing effective biomedicines of animal origins. The animal biomedicines; Nematode Extract (NE) and Gall Protein (GP) when applied by foliar spray, are highly effective at 1.3mg/plant and 2.01mg/plant, respectively in ameliorating root-knot disease of tomato (*Lycopersicon lycopersicum*, Cv. Pusa Ruby) caused by *Meloidogyne incognita* (Kofoid & White) Chitwood, promoting plant growth and fruit production. The nematode extract (NE) is more effective than the gall protein (GP) in this respect. Both the extract is thought to induce systemic acquired resistance response of the treated plants through the expression of pathogenesis-related proteins, and may prevent 21st-century pandemics COVID-19 like virus diseases by boosting immunity, resisting toxic effects on the environment, and improved agriculture green-socio-economy aquatic-science-technology-communication application issues, and it may also give a good scope for new advanced-development and research in the field "Agricultural, Horticulture, Earth-Environmental-Clinical-Toxicology-Health-Research, Science-Technology-Communication-Socio-Economy and May be Controlled 21st-Century Pandemic Diseases in Future New Normal Situation Also".

Keywords: Enriched; Agriculture-Horticulture; Biomedicines; Tomato-and-Coronavirus 2-Diseases'

Introduction

Root-knot nematodes infest almost all kinds of crops, mostly vegetables, and in India the amount of damage, they inflict on plants, accounts for 10-40% of the total crop losses annually [1]. Thus the disease caused by the plant-parasitic nematodes is a serious problem in our country and this indirectly affects our agricultural economy. A number of chemical pesticides have been extensively used by the farmers; early 80,000 tons of chemical pesticides are applied on the Indian soil annually [2]. The safe alternative method to the conventional method of nematode control is to induce the natural defense response of the plants against nematode attack [3-6]. In the present study, it is exploited systematic signaling and induced natural defense in the host plants by applying animal-biomedicines; nematode extract (NE), and gall protein (GP).

MATERIALS AND METHODS

Preparation of Biomedicines:

The animal biomedicines; nematode extract (NE) and total gall

protein (GP), *Meloidogyne incognita* females, and gall roots were collected from roots of the tomato plant grown in the experimental garden of the Department of Zoology, VisvaBharati University, Santiniketan – 731235. They were washed with sterile water, homogenizer and extracted with 90% ethanol at room temperature (25±2°C) for five days and centrifuged at @3500 rpm for 5 minutes. Both the extract supernatant was collected and allowed to evaporate at room temperature (25±2°C) and both the biomedicines residues were kept over anhydrous CaCl₂ for dehydration and stored at 4°C. The biomedicines residue was mixed with sterile distilled water just before application on the test plants [3-7].

Pot Test and Inoculation:

The aseptically germinated seeds of tomato, *Lycopersicon lycopersicum*, Cv. Pusa Ruby was shown one seed/pot (32 cm diameter) containing a mixture of clay soil and composted manure (2:1v/v and 4.5kg), and the soil-filled were earlier treated with boiling water five times. The pots were divided into four groups of 10

pots each. The groups were: uninoculated untreated, inoculated untreated, GP treated inoculated, and NE treated inoculated. When the plants were at the 8-leaf stage they were inoculated with *M. incognita* juveniles (J2) @ 2580±55 J2/plant [3-6].

Treatments:

All the treatments were applied by foliar spray @ 10ml / plant in five steps spanned with an interval of 1 hr., and the treatments were given three days before inoculation. In the NE treated group was received 1.3 mg/plant nematode extract and in the gall protein treated group were received 2.01 mg/plant respectively. Control plants were sprayed with an equal amount of sterile distilled water. The plants were irrigated regularly and the experiment was conducted outdoors in the experimental garden of Department of Zoology, Visva-Bharati University, Santiniketan – 731235, West Bengal, India, at an ambient atmospheric temperature (28±5°C) and humidity (80±5%).

Result Harvesting:

All the plants were uprooted 53 days after the sowing of seeds. The following measure measurement was taken: biomass of shoot,

root, and fruits, root gall number, nematode population in roots (2g) and soil (200g), protein content of root and fruits. Proteins were estimated by the Folin-phenol method [8]. All the data were analyzed by analysis of variance (ANOVA). The experiment was repeated five times with similar results and the data from the third experiment were represented in Table 1.

Pretreatment with NE and GP:

The pretreated nematode extract (NE) and gall protein (GP) extract significantly ($P \leq 0.05$ by ANOVA) increased plant growth in terms of shoot weight, fruit weight as compared to the inoculated and untreated plants [Table 1]. Root galls, nematode population in the root and soil, and root protein content were significantly ($P \leq 0.05$ by ANOVA) reduced in pretreated plants as compared to the untreated ones [Table 1]. Protein content in green fruit was significantly reduced in inoculated plants as compared to the uninoculated ones, and both the pretreatments with biomedicines (NE and GP) restored the condition to some extent [Table 1]. NE-bio-medicine treatment showed better plant growth and lesser intensity of root-knot disease as compared to the GP-bio-medicine pretreatment [Table 1].

Table 1: Effect of the GP and NE on tomato plantsX growth, development and protein

±	Fresh Biomass (g)X			Number of FruitsX	Number of Root-GallsX	Nematode PopulationX		ProteinX Content (%)	
	Shoot	Root	Fruits			Root (2g)	Soil (200g)	Root	Fruits
Uninoculated Untreated	206.48bZ±12.02	30.02d ±1.92	132.16a±10.02	12.32a ±4.04	00			3.94b ±0.22	0.25a ±0.05
Inoculated Untreated	98.03d ±12.01	76.86a ±6.42	43.97c ±4.63	05.97c ±4.03	3500.65a ±196.15	4389a ±265	769a ±49	6.03a ±0.01	0.14c ±0.02
GP-Treated Inoculated	193.31c±6.29	43.28b ±8.02	81.45b ±6.25	08.86b ±4.02	875.11b ±83.71	329b ±32	187b ±29	3.42b ±0.24	0.19b ±0.03
NE-Treated Inoculated	208.02a±4.68	36.02c ±2.04	133.56a ±9.28	13.98a ±3.04	58.65c ±67.35	178c±22	119c ±29	2.97c ±0.19	0.21b ±0.03

‘X’-Mean of 10 replicates with S.E.

‘Y’-Tomato plants inoculated at 8-leaf stage with *M. incognita* juveniles (2580±55 J2/pot) and harvested 53 days after sowing of germinated seeds.

‘Z’-Means carrying same letters in a column are not significantly different ($P \leq 0.05$) by analysis of variance.

DISCUSSION

The present study very clearly shows that the NE and GP could induce some resistance in tomato to *M. incognita* infection, and NE treatment was more effective than GP treatment. Results of the pot test further indicate that both the treatments (NE and GP) could promote the growth of the plants and the resistance induced by NE and GP were systematic. It is reported that nematode extract had no toxic direct effects on nematodes [3-6]. It can be assumed that NE and GP could induce the synthesis of some antagonistic substances. Lectin accumulated in galled regions of root of *Hibiscus esculentus* infected with *M. incognita* [9]. Systemic acquired resistance can be induced in different crop plants by localized virus infection, non-pathogenic and pathogenic microorganisms or their

culture filtrates, or by salicylic acid and this form of resistance protects plants from a broad spectrum of pathogens and work systematically in many cases [3-6,10-20]. *M. incognita* is known to share common antigens with its host plants [21]. It appears that during natural infection with the nematode, host plants show minimal defense responses to the nematode because of this antigenic similarity. NE and GP containing various antigens may induce defense responses involving a number of pathogenesis-related proteins (PR-proteins) which the nematode fails to tolerate [22-23]. It is primarily observed that in lady's finger disease treated with NE, showed the highest number of root protein (13 and 24) but in inoculated untreated root was 18 numbers and in uninoculated root was 11 numbers. Those showed that NE served as the stimulus for

the expression of many proteins particularly the defense-related proteins which later provide resistance to nematode infection [5, 22-23].

Thus, nematode extract and gall protein will serve as very effective biomedicines which would be cheap, non-phytotoxic, non-pollutant, and conserve our biodiversity, and scholars which give a good scope for new development and future research in the field 'Agricultural and Horticulture Sciences' also[24-29].

Future Research:

The nematode extract and gall protein could induce the production of new defense-related genes in the test plant may be confirmed by further biochemical analysis [3-6], and in near future, synthetic production of both the biomedicines; NE and GP will be the potential cost-effective personalized-biomedicine OR social vaccine OR vaccine against coronavirus-2 like pandemic diseases by boosting immunity, and helping policy initiative clinical research in all areas in the field of agricultural and horticulture Sciences also [30-31].

Conclusion

The animal biomedicines; nematode extract (NE) and gall protein (GP), boosted the growth of tomato plants by inducing defense-resistance PR-proteins and may prevent 21st-century pandemics like virus diseases by increasing immunity, resisting toxic effects, and improved potential animal-biomedicines, advanced in agriculture, aquaculture, clinical toxicology, green-socio-economy, aquatic-science technology, and communication application issues, and it may also give a good scope for new development and research in the field "Agricultural, Horticulture, Earth-Environmental-Clinical-Toxicology-Health-Research, Science-Technology-Communication-Socio-Economy and May be Controlled 21st-Century Pandemic Diseases in Future New Normal Situation Also".

Acknowledgements

The work described here is the part of the UGC and CSB project, Government of India to NCS. I thank Ex- Prof. Dr. N.C. Sukul, Parasitology Laboratory, Department of Zoology, Visva-Bharati, Santiniketan-731235, and West Bengal, India for all kinds of support. I also thankful to Prof. Dr. N. Banerjee and Dr. K. Ghosh, Plant Biotechnology Laboratory, Department of Botany, Visva-Bharati University, Santiniketan for needful helps.

References

1. Sen J (1983) Survey of common plant-parasitic nematodes in Birbhum. Ph.D. thesis, Dept of Zool, Visva-Bharati University, Santiniketan – 731235, West Bengal, India.
2. Paul A, Sinhababu SP and Sukul NC (1995). Effect of Meloidogyne incognita infected mulberry plants and their treatment with nematicides on feeding silkworms, Bombyx mori L. Sericologia 35: 577-581.
3. Datta SC, Sinhababu SP, Banerjee N, Ghosh K and Sukul NC (1998 a). Meloidogyne incognita extract reduces M. incognita infestation of tomato. Indian Journal of Nematology 28:1-5.
4. Datta SC, Datta R, Sinhababu SP and Sukul NC (1998 b). Acaciasides and root-knot nematode extract suppress Meloidogyne incognita infection in lady's finger plants. Proceeding

- of National Seminar on Environmental Biology, Visva-Bharati, Santiniketan, West Bengal, India, April 03-05, 1998, (Ed.) by Aditya, A.K. & Haidar, P. 205-209.
5. Datta SC (1999). Bionematicides in the control of root-knot nematode. Ph.D. thesis, Deptt. of Zoology, Visva-Bharati, Santiniketan, West Bengal, India.
6. Mukherjee A, Mondal P and Sinhababu SP (2020). Nematode extract-induced resistance in tomato against Meloidogyne incognita. Indian Journal of Science and Technology 13: 1476-1479.
7. Datta SC (2020a). Enriched Science and Technology Communication Economy in Agriculture by Use of Acacia sides as Potential Bio-Agents against Various Pathogens. Advances in Agriculture, Horticulture and Entomology 2: 1-13.
8. Lowry OH, Rosebrough NJ, Farr AL and Randall RJ (1951) Protein measurement with Folin-phenol reagent. Journal of Biological Chemistry 193: 265-275.
9. Das S, Sukul NC, Mitra D and Sarkar H (1989) Distribution of lectin in nematode infested and uninfested roots of Hibiscus esculentus. Nematologica Mediterranea 17: 123-125.
10. Kiessig DF and Malamy J (1994). The salicylic acid signal in plants. Plant Molecular Biology 26: 1439-1458.
11. Kiessig DF, Durner J, Noad R, Navarre DA and Wendehenne D, et al. (2000). Nitric oxide and salicylic acid signaling in plant defense. Proceedings of the National Academy of Science, USA 97: 8849-8855.
12. Descalzo RC, Rahe JE and Mauza B (1990). Comparative efficacy of induced resistance for selected diseases of greenhouse cucumber. Canadian Journal of Plant Pathology 12: 16-24.
13. Kuc J and Strobel NE (1992). Induced resistance using pathogens and nonpathogens. In: Biological Control of plant Diseases, (Eds. E.C. Tjamos) Plenum Press, New York: 295-300.
14. Mauch-Mani B and Metraux JP (1998). Salicylic acid and systemic acquired resistance to pathogen attack. Annals of Botany 82: 535-540.
15. Merra MS, Shivanna MB, Kageyama K and Hyakumachi M (1994). Plant growth promoting fungi from Zoysia grass rhizosphere as potential inducers of systemic resistance in cucumbers. Phytopathology 84: 1399-1406.
16. Nandi B, Sukul NC, Banerjee N, Sengupta S and Das P, et al. (2002). Salicylic acid enhances resistance in cowpea against Meloidogyne incognita. Phytopathologia Mediterranea 41, 39-44.
17. Nandi B, Kundu K, Banerjee N and Sinhababu SP (2003). Salicylic acid-induced suppression of Meloidogyne incognita infestation of okra and cowpea. Nematology 5: 747-752.
18. Ross AF (1961). Systemic acquired resistance induced by localized virus infection in plants. Virology 14: 340-358.
19. Schneider M, Schweizer P, Meuwly P and Metraux JP (1996). Systemic acquired resistance in plants. International Journal of Cytology 168: 303-340.
20. Sukul NC (1970). Inhibition of nematode infestation of wheat seedlings by Polygonum hydropiper. Nature 226: TIX-TTL.
21. McClure, Michael A, Misaghi I and Night EJ (1973). Shared antigens of parasitic nematodes and host plant. Nature 244: 306.
22. Datta SC (2020b). Acacia auriculiformis-Extract Synthesis PR-Proteins Developed Potential Biomedicines-Vaccine against Okra-Diseases and COVID-19: Improved Science

- Technology Communications Bio-Economy Applications. 8(10): 249-270.
23. Datta SC (2020c). Cina-Pretreatments Act as Potential-Bio-medicine-Vaccine against COVID-19 and Okra-Plant-Diseases: Synthesis PR-Proteins Increased-Immunity Improved Biomedicines-Economy Applications Science-Technology-Communications. International Journal of Ayurveda. 5: 05-26.
 24. Datta SC (2020d). Okra Maybe Potential Cost-Effective Personalized-Biomedicines Social-Vaccine against COVID-19: Improved Immunity Food-Security Green-Economy Science-and-Technology-Communication Applications. Innovative Journal of Medical Sciences 4: 5-20.
 25. Datta SC (2020e). Intercropped Cowpea Maybe Use as Bio-medicine Improved Immunity against COVID-19: Enriching Science and Technology Communication Applications Food Security Economy. Diagnosis and Therapies Complementary and Traditional Medicine. 2020: 35-48.
 26. Datta SC (2020f). Weeds-Vegetables and Fruits Act as Potential Biomedicines against COVID-19: Enriched Agriculture Biodiversity Socio-Economy Science Technology Communications by Controlling Plants Diseases. Journal of Experimental Biology and Agricultural Sciences, October - 2020; 8(Spl-1-SARS-CoV-2): S139-S157.
 27. Datta SC (2020g). Biomedicines-Aakashmini Cost-Effective COVID-19 vaccine: Reduced Plant-Diseases Enriched Science Technology Communications Socio-Economy Bio-Applications. Global Journal of Bioscience and biotechnology. 9: 127-144.
 28. Datta SC (2020b). Discovery of COVID-19 Vaccine by Using Acaciades as a Phytomedicine Improving Science and Technology Communication Applications- An Ideas. Open Access Journal of Biogeneric Science and Research 2: 1-30.
 29. Datta SC (2021a). Weed-Plant Act as Vaccine against Plant-and-COVID-19 Diseases: Enriched-Agriculture-Health-Development Socio-Economy Sciences-Technology-Communication-Application. International Journal of Pharmaceutical Sciences and Clinical Research.1:1-17.
 30. Datta SC (2021b). Nematode Extract and Acaciasides Use as Preventive Biomedicines Against Plant Diseases: Improved Earth-Environmental-Health-Research Science-Technology-Communication and May be Controlled 21st-Century Pandemic Diseases! Eart & Envi Scie Res & Rev. 4: 55-60.
 31. Datta SC (2021c). Animal-Biomedicine Controls Root-Knot-Disease in Lentil-Callus-Culture: Enriched Advanced-Clinical-Toxicology Socio-Economy Science-Technology-Communication by Preventing 21st-Century-COVID-19-Like-Pandemic-Diseases. Advances in Clinical Toxicology (Accepted).

Copyright: ©2021 Dr. Subhas Chandra Datta. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.