

## Another Look at the Water Phases That Exist Under Room Conditions

T. A. Yakhno\* and V. G. Yakhno

Federal Research Center Institute of Applied Physics of the Russian Academy of Sciences (IAP RAS), Russia

### \*Corresponding Author

T. A. Yakhno, Federal Research Center Institute of Applied Physics of the Russian Academy of Sciences (IAP RAS), Russia.

Submitted: 2025, Oct 20; Accepted: 2025, Nov 21; Published: 2025, Dec 04

**Citation:** Yakhno, T. A., Yakhno, V. G. (2025). Another Look at the Water Phases That Exist Under Room Conditions. *J Applied Surf Sci*, 3(2), 01-12.

### Abstract

A brief review of the history of the problem is intended to trace the progress in the development of scientific views on the structure of water. The discoveries in this area made in the first half of the last century are considered in comparison with modern data. It is shown that due to the de facto ban on the study of the structure of water in 1971, work in this direction was frozen for a long time. The paradox is that many forgotten truths have been independently "rediscovered" in our time, which speaks of their authenticity. On the basis of scientific literature and their own research, the authors come to the conclusion that water under room conditions is a micro dispersed system, one phase of which is represented by continuous water, and the other - by polywater, discovered in the 70s of the last century.

**Keywords:** Phases of Water; Microstructure, Dispersion, Physical and Chemical Properties, Polywater

### 1. Introduction

From the standpoint of world academic science, water at room temperature and humidity is still an association of H<sub>2</sub>O molecules, molecular chains and nanometer-sized clusters dynamically changing their structure in femtosecond time [1,2]. By introducing certain assumptions, the authors of such views were able to explain certain anomalous properties of water and ice with the help of molecular modeling. At the same time, information about the presence of other phases of water coexisting with the continuum one was not taken into account.


Meanwhile, more than 70 years ago (see review), convincing experimental facts were obtained that indicate that water near hydrophilic surfaces forms a new phase - a nearsurface layer, which is very different in terms of the complex of physical properties from bulk water [3]. The thickness of this layer is up to hundreds of microns. Direct evidence of the existence of a new water phase was presented, including in terms of refractive index, diffraction of X-ray and neutron radiation, viscosity, adhesion, and the presence of long-range order in structural ordering. This information has been repeatedly confirmed by many researchers

using different techniques of physical experiments. Experimental evidence was obtained for the fact that the structure of water formed at hydrophilic surfaces corresponds to the structure of ice and represents layers of hexagonal structures superimposed on each other in one, two, three or four layers [4-7]. It has been shown that thin ( $h < 150 \mu\text{m}$ ) layers of water located between glass surfaces have shape elasticity [8]. Films with a thickness of  $0.35 \mu\text{m}$  had the rigidity of metallic lead [9]. The viscosity of water in the gap between glass plates separated by  $1 \mu\text{m}$  was many times higher than the viscosity of water in the volume, and at a distance between the plates of  $2.5 \mu\text{m}$  it was times higher [10,11].

The data on the existence of a new phase of water under room conditions were not accepted by representatives of "classical science". Suffice it to recall the opening and closing of 1 "polivoda" (water-II) [12-14]. The water obtained by Soviet scientists during the condensation of water vapor in ultra-thin quartz capillaries had a significantly higher viscosity, density, refractive index, and surface tension than ordinary water. Its dissolving power was less. Experiments on obtaining polywater have been successfully reproduced in many research laboratories around the world,

including Europe, the USA and Australia, and were widely discussed in the press and at international forums [15,16]. The molecular weight of water-II determined from the vapor density exceeded this indicator for ordinary water by 8-10 times, and the absorption in the IR region was characterized by two peaks that are not characteristic of any other known substance [17]. Based on the data obtained, it was assumed that the new substance is a polymer consisting of water molecules combined into hexagonal cells (Figure 1) [17]. Upon the formation of water-II from ordinary water vapor on the surface of fused quartz or glass B.V. Deryagin

assumed the existence of a special kind of condensation catalysis [16]. However, in water samples in a number of laboratories, inorganic and organic impurities were found [16,18-20]. In addition, unique IR absorption bands characteristic of polywater ( $1595\text{ cm}^{-1}$  and  $1400\text{ cm}^{-1}$ ) were also found in a sample of human sweat [21]. All this, ultimately, served as a reason for accusing the authors of the discovery of insufficient purity of the experiment and called into question the very existence of the phenomenon [22]. At the same time, for example, the hypothesis about the possible existence of polywater in biological fluids was ignored.



**Figure 1:** Representation of the authors about the molecular structure of polywater: a - structural diagram of polywater, consisting of a network of hexagonal blocks; b - monolayer [17]

Under the onslaught of skeptical colleagues, the authors were forced to publicly “repent” [23,24]. However, it turned out that the “impurities”, interacting with the surrounding water, significantly change its properties at the interface. In the work [25, p. 68] we read: “*A new stage in the study of surface phenomena was the discovery of a special structure of layers of polar liquids, bordering with the surface of an adjacent phase. Under the influence of this surface, polar molecules are oriented relative to it in a certain way, and this orientation is transferred from layer to layer over a certain distance. A similar phenomenon was observed for the first time in water layers adjacent to the surface of a hydrophilic body. In this case, the degree of orientation decreases with distance from the substrate, becoming insignificant at a distance of the order of the length of hundreds of molecules. As a result of a change in the structure, the properties of the boundary layers of water (refractive index, density, viscosity, dissolving ability) differ from the normal one’s characteristic of the volume of water. The freezing point also changes. It is for this reason, as we have shown, that in permafrost regions the layers of water adjacent to the particles of hydrophilic clays do not freeze, at least at temperatures not lower than  $-10\text{ }^{\circ}\text{C}$ .*” And further (p. 70): “*According to its optical properties, the boundary layer is homogeneous and anisotropic, like liquid crystals, and is separated from the bulk phase by a sharp interface. Thus, in addition to the previously known solid, liquid and gaseous phases, there are “boundary phases” of a strictly defined thickness, which naturally decreases with increasing temperature.*”

## 2. Current Data

After the “exposure” of the researchers of the new phase of water, the work in this direction was frozen and partially forgotten. However, after half a century, the opinion that liquid water is two-phase resurfaced. According to modern data, the simplest model of the structure of liquid water assumes the existence of two separate microscopic phases [26–31]. Many physical quantities exhibit behavior that suggests the presence of two states in liquid water. For example, infrared and Raman spectra, as well as the presence of an isobestic point in the Raman spectra, are considered as a clear sign confirming the model of a mixture of two phases [31]. According to Preparata et al, liquid water is a two-component liquid system consisting of a coherent phase (about 40% of the total volume at room temperature) and an incoherent phase [32]. Thin films of water of ice-like structure with a thickness of several molecular layers on hydrophilic surfaces have been rediscovered [33–37]. Obviously, these water films are formed from water vapor present in the ambient air, since their thickness depends on humidity.

At this nanoscale level, interfacial water near hydrophilic surfaces exhibits a viscosity of about 2 to 106 times that of bulk water, while no significant changes in the viscosity of water near hydrophobic surfaces are observed [38]. When a hydrophilic surface is immersed in water, a layer of structured liquid crystalline water up to  $500\text{ }\mu\text{m}$  thick forms near it in a few minutes, displacing ions and microparticles from it, which are usually present in an aqueous


medium [39,40]. An exclusion zone (EZ) is formed, due to which the concentration of ions and particles in free water increases, increasing also its osmotic pressure [41]. It is believed that the thin ice-like layer that covered the previously dry surface is used as a matrix for EZ growth [42–44]. Molecular dynamics has shown that water / solid and water / air interfaces strongly affect the structure and intermolecular dynamics of sessile water droplets, and suggests that intermolecular dynamics in other systems with interfaces differs from that in liquid water [45]. Interfacial water EZ is much more viscous than bulk water [46].

Despite the abundance of experimental evidence for the existence of the near-wall phase of water, a number of researchers stubbornly continue to deny this fact [47-51]. Several alternative explanations for EZ have been presented, in particular, Schurr's theory based on chemotaxis and diffusionophoresis [47]. This theory was strongly supported by Elton et al and Esplandiu et al [48,49]. Another view of the formation of EZ is associated with swelling in water and partial destruction of Nafion, a solid polymer electrolyte, a popular owner of hydrophilic surfaces [50,51]. When this polymer swells, its fibers partially untwist and go out into bulk water. Such elongated strands can also form a rigid, parallel palisade that can repel colloidal microparticles; this, according to the authors of, creates the effect of an exclusion zone near Nafion [51]. Analysis of alternative opinions is beyond the scope of this work. Pollack clearly commented on skeptical statements about the formation of a special near wall layer of water: “In sum, the four issues of

challenge fail to derail the original line of interpretation, that the EZ is a distinct, ordered structure that excludes solutes. Ample evidence for this interpretation (which Schurr seems to have missed) lends strength to this view” [52]. Further, we present works that, in our opinion, clearly indicate the existence of an additional phase in liquid water.

### 3. Additional Research Methods and Materials Used


Two decades ago, sensational information appeared that liquid water is a heterogeneous polydisperse system [53-58]. To detect this fact, the following methods were used: visualization using laser radiation, which makes it possible to capture an insignificant difference in the refractive indices of two phases of water; laser modulation interference phase microscopy and scattering of laser radiation in aqueous media purified from solid impurities; static and dynamic small-angle light scattering; method of small scattering angles in bidistilled water [53-56,58]. Measurements showed the presence in water of a spectrum of clusters in the range of 1.5 - 6.0 microns, as well as giant clusters tens and hundreds of microns in size. In Figure 2 shows a microphoto of ultrapure water preparations (OST 34-70- 953.2-88, specific conductivity 0.04-0.05  $\mu\text{S} / \text{cm}$ , pH 5.4-7.0), distilled water (TU 2384-009-48326337-2015, specific 84 electrical conductivity 4.5  $\mu\text{S} / \text{cm}$ , pH 7.0) and mineral water "Seraphim Dar" (mineralization 0.05-0.12 g / l). The thickness of the liquid layer in the preparations is 8  $\mu\text{m}$ , the width of each frame is 3 mm.



**Figure 2:** The structure of ultrapure water (a, b) immediately after depressurization of the container is also a microdispersion and fundamentally corresponds to the structure of distilled (c) and mineral (d) water. The width of each frame is 3 mm [59]

The fact that water is a polydisperse system can be verified by observing under a microscope a drop suspended in a micro-hole


0.5 mm in diameter made in a plastic substrate (Figure 3).




**Figure 3:** Microstructure of liquids in the "hanging drop" preparation. Liquid placed in a hole in a plastic plate with a diameter of 0.5 mm. Microscopy "through liquid": a - distilled water; b - tap water; c - water of the Black Sea [59]

After evaporation of equal volumes of water samples presented in Figure 3, it turned out that the mass of the solid sediment corresponds to the proportion a: b: c = 0.2: 0.45: 2.5. In other words, the degree of dispersion of water in the volume turned out to be inversely proportional to the mass of substances dissolved

in it. The fact that the dispersed phase of water is not an optical illusion can also be seen by examining the layer of water between the slide and the cover glass during the evaporation of the liquid phase (Figures 4, 5).




**Figure 4:** The "crushed drop" preparation is a thin layer of liquid between the slide and the cover glass during the evaporation of the liquid phase of water through the slits between the glasses. Evaporating through the edges of the preparation, the liquid recedes to the center, leaving a solid sediment on its way (Figure 5): L - Liquid phase, S - Solid sediment




**Figure 5:** Microphoto of a part of the "crushed drop" preparation (Figure 4) in the process of evaporation of the liquid phase. Fragment a: on the right (light part) - an evaporating liquid; on the left (dark part) the dispersed phase of water settling on a substrate; b - fragment a, limited by a light square, at higher magnification. Frame width "a" is 3 mm [60]

In smears prepared from water and aqueous solutions, the dispersed phase is visible in the form of aggregates (Figure 6) [61].




**Figure 6:** Aggregates of the dispersed phase in dried smears: a- distilled water; b - dry white wine; c, d - water solution of “Nescafe Gold” freeze-dried coffee. Microphoto. The width of each frame is 3 mm [61]

In Figure 7 clearly visible aggregates of the dispersed phase of water in a thin layer of an aqueous solution of freeze-dried coffee “Nescafe Gold” (250 mg / 100 ml).



**Figure 7:** A layer of coffee solution between the slide and the cover slip is 8 μm thick. Aggregates of the dispersed phase are visible


Below is a diagram of the proposed organization of the structural unit of the micro-aggregate of the dispersed phase (Figure 8).



**Figure 8:** Scheme of the proposed organization of the structural unit of the microaggregate of the dispersed phase

It is obvious that individual microaggregates can be combined into larger formations. Earlier, we put forward a hypothesis, which received experimental confirmation that the basis of the dispersed phase of water are microcrystals of NaCl salt, surrounded by a thick liquid crystal shell of hydrated water [60,62]. The evolution


of water sediments drying in the open air has been traced [62,63]. It was shown that the aggregates do not evaporate at room temperature, exhibit the properties of a viscous liquid, do not dissolve in organic solvents (ether, hexane), but break down in strong salt solutions (Figure 9).



**Figure 9:** Dried sediment of coffee solution: a - after two hours of extraction with hexane, the dispersed phase aggregates retain their integrity; b - after adding NaCl to the coffee solution at a dose of 4.0 g / 100 ml, the dispersed phase aggregates are destroyed

The fact that an increase in NaCl concentration destroys the EZ structure was also reported by the authors of publications [64–66]. This fact confirms the common nature of the dispersed phase of water and EZ.

According to the results of gas chromatography-mass spectrometric analysis of a sample of dried coffee for water content, in the temperature range of 200 - 300 ° C, the amount of water released from the analyzed sample increases abruptly, which may be associated with the destruction of its polymer structure (Figure 10, Table 1.) [67].




**Table 1.**  
Dependence of the mass fraction of released water on the temperature of thermal desorption  $\eta$

№	Thermal desorption temperature	Sample weight (mg)	S (H <sub>2</sub> O)	w(H <sub>2</sub> O), %
1	100	7.2	4921843	1.6
2	150	11.7	33178809	6.5
3	200	10.6	33876750	7.4
4	300	11.2	97991324	20.1

**Figure 10:** Results of gas chromatography-mass spectrometric analysis of a sample of dried coffee for water content. A sharp increase in water evaporation occurs in the range of 200 - 300 ° C, which may be associated with the destruction of the polymer structure of hydrated water

To determine the relative density of dispersed phase aggregates, we centrifuged a Nescafe Gold coffee solution (250 mg / 100 ml of water) for 20 min at 1.5 thousand rpm and prepared smears from

the upper, middle, and lower parts of the centrifugate. The results are shown in Figure 11.



**Figure 11:** Microphoto of smears of Nescafe Gold coffee (250 mg / 100 ml of water), before and after centrifugation for 20 min at 1.5 thousand rpm. Samples were taken before centrifugation (a) and after, from different heights of the centrifuge tube: b - from the upper part, c - from the middle part, d - from the lower part. Frame width - 3 mm


Most of the complete aggregates were contained in the upper and middle parts of the tube. From this fact, it can be concluded that the dispersed phase is close in density to continuous water. More accurate measurements are needed.

#### 4. Discussion

The authors reported the discovery of isolated stable water clusters ranging in size from tens of nanometers to microns as a result of the evaporation of a highly dilute sodium chloride solution ( $10^{-7}$  M) at room temperature and normal atmospheric pressure [68]. Stable water clusters are electrically charged (from 1.257 V to  $-0.246$  V), as shown by studies using an electric force microscope. Raman scattering and infrared spectra of sediments show similar but not identical characteristics to liquid water. We will return to this information later. A review of modern works on the existence of isolated stable water clusters was made by Ho [69]. In studies, water structures are described, some of which, according to the authors' assumption, are accumulations of gas microbubbles [70,71]. The above facts leave no doubt that water is a polydisperse system. According to Borilo [72, p. 18] “*Dispersion (or specific surface area) is an independent thermodynamic parameter of the state of the system, a change in which causes corresponding changes in*

*other equilibrium properties of the system. Dispersion acts as an intense feature of the system. This value can be compared with concentration, that is, with the amount of surface per unit volume. Thus, the surface acts as a separate component. In this case, you can draw an analogy with temperature or pressure. Preliminary consideration shows that dispersion is an independent and full-fledged thermodynamic parameter of the system. ... Thus, the surface tension is a partial derivative of any thermodynamic potential with respect to the interfacial area at constant corresponding parameters.*” This pattern was also demonstrated with specific examples in [73].

The destruction of water units by mechanical action does not raise questions. It turned out that low-intensity light also leads to the dissociation of the microdispersed phase of water. In Figure 7 shows the result of irradiating water with a 20 mW green LED laser for 10 minutes. Natural mineral drinking water "Serafimov Dar" was used in the experiment. Water was poured into a 100 ml glass beaker; the light source was placed vertically directly above the water surface in the central zone equidistant from the glass walls. The microdispersed phase of water was examined under a microscope before and after irradiation (Figure 12).



**Figure 12:** Microstructure of natural mineral drinking water "Serafimov Dar". Microscopy of a liquid layer ~ 16  $\mu\text{m}$  thick before (a) and after (b) irradiation of a glass with water with a 20 mW green laser for 10 minutes

It is obvious that irradiation led to the dissociation of large aggregates of the microdispersed phase. Simultaneously, there was also a decrease in the electrical capacity of the liquid by ~ 1 pF [74]. The changes were reversible and returned to baseline within an hour. A review of the available literature on the results of the impact on water of various disturbing factors of a physical nature made it possible to conclude that they all lead to the same consequences - an increase in pH, a decrease in ORP, an increase in electrical conductivity, and a decrease in viscosity [73]. We associate these changes with the destruction of aggregates of the dispersed phase of water and an increase in the specific surface area of the system.

The development of an original technique for obtaining samples of "solid-state water" by the method of repeated rinsing of hydrophilic surfaces in high-purity water and subsequent lyophilization of the used liquid allowed the authors to obtain solid-state microstructured sediment [75-82]. The structure of the sediment depended on the nature of the hydrophilic substrate used (nafion, cellophane, cotton wool, filter paper). The physicochemical properties of water after repeated contact with different hydrophilic surfaces had a number of similar features: all samples contained associates  $10^{-4}$  m in size, but in different concentrations, had a higher density and electrical conductivity than the initial water. According to the authors, the associates were a detached from the surface and crumpled EZ, as a result of mixing and deformation of the surfaces [81]. When water was disturbed by cellulose preparations, in particular, pH, electrical conductivity, heat of mixing with acids or bases, and density changed. It is natural to assume that the nature of these changes is also associated with an increase in the specific surface area of the system caused by multiple mechanical disturbances.

In particular, this fact is confirmed by the linear correlation between electrical conductivity and other physicochemical variables [75]. In this case, the use of quantum electrodynamic field theory to


explain the observed changes seems to us excessive, since the determining events, as we see, develop not at the atomic, but at the micrometer level [83,84]. Thus, the classical concepts of water as a liquid homogeneous at the micrometer level must be revised, since they do not correspond to reality.

Based on the features of the IR spectrum, the deconvolution of the OH stretching peak between  $2880$  and  $3800\text{ cm}^{-1}$  shows the complete disappearance of the higher energy component by about  $3600\text{ cm}^{-1}$  relative to bulk water [75]. This feature is characteristic of OH stretching in ice. It is also noted that "*Other features appearing at  $1418\text{ cm}^{-1}$  and  $1105\text{ cm}^{-1}$ , that are not present in liquid water, are difficult to attribute univocally at the present stage of knowledge.*" [74, p. 22]. Let us recall that the authors of the discovery of polywater faced a similar problem more than 50 years ago: "*the complete absence of absorption from  $2500$  to  $4000\text{ cm}^{-1}$ , the presence of a strong band at about  $1595\text{ cm}^{-1}$  and an intense doublet in the region of  $1400\text{ cm}^{-1}$* " [17]. ... And further: "*... we believe that water is restructured on the fused quartz or Pyrex surface into a polymeric form with bonds and bond energies completely different from that of ordinary water. Its properties are not those of water, and it should not be considered to be or even called water; any more than the properties of the polymer polyethylene can be directly correlated to the properties of the gas ethylene. In the case of the formation of polyethylene, a bond change occurs. In the case of polywater, a bond change also occurs, although it has previously been unknown. ... Vibrational spectra indicate unique stable polymeric structure*" [17, p. 1486]. As you can see, the authors of the discovery of "polywater" also obtained it upon contact of ordinary water with hydrophilic surfaces - fused quartz and glass. Let us return to the publication by Lo et al [68]. The structures discovered by the authors - stable water clusters - at room temperature also had characteristic differences in the IR spectrum from ordinary water: the absorption peaks for pure water are  $3283.5\text{ cm}^{-1}$  and  $1634.5\text{ cm}^{-1}$ . The absorption peaks for stable-

water-clusters are  $3371.4\text{ cm}^{-1}$ ,  $1639.5\text{ cm}^{-1}$ ,  $1342\text{ cm}^{-1}$ ,  $822.5\text{ cm}^{-1}$ . We see the same "red shift" and characteristic absorption for "water structures" obtained by other authors. The IR spectrometric analysis of the Nescafe Gold coffee solution<sup>1</sup>, in which we clearly observed aggregates of the dispersed phase of liquid crystalline water, revealed similar shifts [67]. To obtain the IR spectrum, a part of the sample dried to constant weight was ground in an agate mortar with a certain amount of KBr powder and pressed into a tablet. The IR spectrum was recorded on an IRRAFINITY<sup>1</sup> instrument (Shimadzu) in the range of wave numbers 400 - 4000  $\text{cm}^{-1}$ . The spectrum of a pure KBr tablet was used as a background. In Figure 13 shows the IR spectrum of the sample at room

temperature. It can be seen that, along with residues of free water ( $3381.2\text{ cm}^{-1}$ ), there are bands characteristic of structured water ( $\sim 1600\text{ cm}^{-1}$ ,  $\sim 1400\text{ cm}^{-1}$ ,  $\sim 1100\text{ cm}^{-1}$ ).

Thus, all the detected stable aqueous structures have similarities in the features of the IR spectrum characteristic of polywater. Does this mean that there are only two phases of water - continuous and near-wall (EZ), and the latter can also exist in the form of "water structures": EZ near hydrophilic impurities or EZ, exfoliated from the substrate [60,64]. This second phase of water has a unified structure and corresponds to the previously described polywater [17].



**Figure 13:** IR spectrum of dried powder of aqueous coffee solution at room temperature

By the way, certain differences in the appearance of "dry water" obtained by iteration and lyophilization from different substrates (Nafion, cellophane, cotton wool, paper filter) can be explained by the presence of microparticles of these substrates in the sediments (see, for example, Figure. 2b; Figure. 1-3; Figure. 6.7; Figure. 6.7 [74,78,80,81]. All used substrates have a fibrous structure, the micro-fragments of which can be torn off during repeated friction against water [82,83]. Since all the substrates used are hydrophilic, EZ is formed on their surface, which is revealed through the characteristic absorption in the IR spectrum.

This brief review is devoted to the discovery and development of only one idea about the interaction of water with hydrophilic surfaces and the existence of two phases of water at room temperature. These are continual water and structured water EZ, which is similar to polywater. This, in our opinion, is sufficient for a consistent explanation of all the phenomena described to date.

We have not touched upon the alternative opinion of a number of physicists - theorists on the structuring of water by the action

of a "vacuum electromagnetic field" on it. According to Giuliano Preparata, Emilio Del Giudice and colleagues, liquid water is a two-fluid system, consisting of a coherent and incoherent phases [83]. In the coherent phase, water molecules vibrate between two electronic configurations in phase with a resonating electromagnetic field (EMF) and, as a result, large stable coherent domains (CD) with a diameter of about 100 nm are formed in it at ordinary temperature and pressure. It is argued that these CDs can be responsible for all the special properties of water etc. Not all authors agreed with the adequacy of the presented model [69,83-86]. An attempt to combine two different views on the structure and dynamics of water was made in [86]. Undoubtedly, the involvement of quantum electromagnetic dynamics in the study of Nature is a promising theoretical tool, however, in the application to the problem considered here, we do not find a place for it. Let us summarize what has been said.

## 5. Conclusion

Under room conditions, water is a microdispersed system with all its inherent properties. There are two phases of water: this is

the usual continuous water and water, which is a cyclic polymer H<sub>2</sub>O, consisting of flat layers of hexagonal cells superimposed on each other (polywater). Polywater can exist as a near wall layer at hydrophilic surfaces (EZ), and as "micron-sized structures" floating in continual water. The structures are prone to the formation of aggregates (coagulates) tens and hundreds of microns in size. The degree of water dispersion determines its physicochemical properties. The downsizing of aggregates by any disturbance leads to a decrease in viscosity, an increase in electrical conductivity, an increase in pH, and a decrease in ORP. That is, the reason for the change in properties lies at the micrometer level, accessible for observation under an ordinary light microscope, and this reason is a change in the "surface concentration". We consider the continuation of research in this area to be promising.

In conclusion, let us remember the statement of A. Schweitzer: "Any truth is first ridiculed and then recognized."

**Funding:** The study has been supported in the frames of the Governmental Project of the Institute of Applied Physics RAS (Project #FFUF-2024-0037)

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Eisenberg, D., & Kauzmann, W. (2005). *The structure and properties of water*. Oxford University Press.
2. Pang, X. F. (2014). *Water: molecular structure and properties*. World Scientific.
3. Henniker, J. C. (1949). The depth of the surface zone of a liquid. *Reviews of modern physics*, 21(2), 322.
4. Macey, H. H. (1942). Clay-water relationships and the internal mechanism of drying. *Trans. Br. Ceram. Soc*, 41, 73-121.
5. Kolkmeijer, N. H., & Favejee, J. C. L. (1933). Structure of emulsoid sol particles and their hydration film. *Nature*, 132(3337), 602-603.
6. Mering, J. (1946). On the hydration of montmorillonite. *Transactions of the Faraday Society*, 42, B205-B219.
7. Williamson, W. O. (1948). *The clay/water relation*. ReSearch, London, 1, 363-371.
8. Derjaguin, B. (1933). Die Formelastizität der dünnen Wasserschichten. *Zeitschrift für Physik*, 84(9), 657-670.
9. Derjaguin, B. (1936). Range of action of surface forces. *Nature*, 138(3486), 330-331.
10. Terzaghi, C., *Angew. Z.* (1924). *Math. Mech.* 4, 10<sup>7</sup>. (As quoted in Henniker's review [3]).
11. Norton, F. H., & Johnson, A. L. (1944). Fundamental study of clay: v, nature of water film in plastic clay. *Journal of the American Ceramic Society*, 27(3), 77-80.
12. Derjaguin, B. V., Churaev, N. V., Fedyakin, N. N., Talaev, M. V., & Ershova, I. G. (1967). The modified state of water and other liquids. *Bulletin of the Academy of Sciences of the USSR. Division of Chemical Sciences*, 16(10), 2095-2102.
13. Ice-nine, Russian style, in the book: S. J. Gould. (1996). *The Mismeasure of Man*. Norton, 1996, 447 pp.
14. Franks, F. (1982). *Polywater*. Cambridge, MA: The MIT Press, pp 208.
15. Deryagin, V. B. (1970). New data on superdense water. *Soviet Physics Uspekhi*, 13(2), 305.
16. Derjaguin, B. Abnormal water - hypotheses and facts. 11.03.2007. (In Russian). Available online: [www.o8ode.ru/article/water/udivit/anomalwater.htm](http://www.o8ode.ru/article/water/udivit/anomalwater.htm) (accessed on 7 May 2008).
17. Lippincott, E. R., Stromberg, R. R., Grant, W. H., & Cessac, G. L. (1969). Polywater: Vibrational spectra indicate unique stable polymeric structure. *Science*, 164(3887), 1482-1487.
18. Rousseau, D. L., & Porto, S. P. S. (1970). Polywater: polymer or artifact? *Science*, 167(3926), 1715-1719.
19. Davis, R. E., Rousseau, D. L., & Board, R. D. (1971). "Polywater": Evidence from electron spectroscopy for chemical analysis (ESCA) of a complex salt mixture. *Science*, 171(3967), 167-170.
20. Kurtin, S. L., Mead, C. A., Mueller, W. A., Kurtin, B. C., & Wolf, E. D. (1970). "Polywater": a hydrosol? *Science*, 167(3926), 1720-1722.
21. Rousseau, D. L. (1971). "Polywater" and sweat: similarities between the infrared spectra. *Science*, 171(3967), 170-172.
22. Ginzburg, V.L. (1971). What problems of physics and astrophysics seem to be especially important and interesting? *Uspehi fizicheskikh nauk*, 103, 87, 93-94. (In Russian).
23. Derjaguin, B. V., & Churaev, N. V. (1973). Nature of "anomalous water". *Nature*, 244(5416), 430-431.
24. Derjaguin, B. V., Zorin, Z. M., Rabinovich, Y. I., & Churaev, N. V. (1974). Results of analytical investigation of the composition of "anomalous" water. *Journal of colloid and Interface science*, 46(3), 437-441.
25. Derjaguin, B.V. (1990). The world of colloid-superficial phenomena. *Vestnik Akademii Nauk SSSR*. The heading "From the notebook of the researcher" 9, 68-75. (In Russian).
26. Pershin, S. M. (2005). Two-liquid water. *Physics of Wave Phenomena*, 13(4), 192.
27. Huang, C., Wikfeldt, K. T., Tokushima, T., Nordlund, D., Harada, Y., Bergmann, U., ... & Nilsson, A. (2009). The inhomogeneous structure of water at ambient conditions. *Proceedings of the National Academy of Sciences*, 106(36), 15214-15218.
28. Maestro, L. M., Marqués, M. I., Camarillo, E., Jaque, D., Solé, J. G., Gonzalo, J. A., ... & Stanley, H. E. (2016). On the existence of two states in liquid water: impact on biological and nanoscopic systems. *International Journal of Nanotechnology*, 13(8-9), 667-677.
29. Gallo, P., Amann-Winkel, K., Angell, C. A., Anisimov, M. A., Caupin, F., Chakravarty, C., ... & Pettersson, L. G. M. (2016). Water: A tale of two liquids. *Chemical reviews*, 116(13), 7463-7500.
30. Russo, J., & Tanaka, H. (2014). Understanding water's anomalies with locally favoured structures. *Nature communications*, 5(1), 3556.
31. Renati, P., Kovacs, Z., De Ninno, A., & Tsenkova, R. (2019). Temperature dependence analysis of the NIR spectra of liquid water confirms the existence of two phases, one of which is in a coherent state. *Journal of Molecular Liquids*, 292, 111449.

32. Roseghini, M. L. The role of QED (Quantum Electro Dynamics) in medicine.
33. Hu, J., Ogletree, D. F., & Salmeron, M. (1995). The structure of molecularly thin films of water on mica in humid environments. *Surface science*, 344(3), 221-236.
34. Aarts, I. M. P., Pipino, A. C. R., Hoefnagels, J. P. M., Kessels, W. M. M., & Van de Sanden, M. C. M. (2005). Quasi-ice monolayer on atomically smooth amorphous SiO<sub>2</sub> at room temperature observed with a high-finesse optical resonator. *Physical review letters*, 95(16), 166104.
35. Jinesh, K. B., & Frenken, J. W. M. (2006). Capillary condensation in atomic scale friction: how water acts like a glue. *Physical review letters*, 96(16), 166103.
36. Jinesh, K. B., & Frenken, J. W. M. (2008). Experimental evidence for ice formation at room temperature. *Physical review letters*, 101(3), 036101.
37. Striolo, A. (2011). From interfacial water to macroscopic observables: A review. *Adsorption Science & Technology*, 29(3), 211-258.
38. Sendner, C., Horinek, D., Bocquet, L., & Netz, R. R. (2009). Interfacial water at hydrophobic and hydrophilic surfaces: Slip, viscosity, and diffusion. *Langmuir*, 25(18), 10768-10781.
39. Pollack, G. (2013). The fourth phase of water: beyond solid, liquid, and vapor. Ebner and Sons Publishers, 320 p.
40. Zheng, J. M., Chin, W. C., Khijniak, E., Khijniak Jr, E., & Pollack, G. H. (2006). Surfaces and interfacial water: evidence that hydrophilic surfaces have long-range impact. *Advances in colloid and interface science*, 127(1), 19-27.
41. Yakhno, T.A. Yakhno, V.G. (2017). Water-induced self-oscillatory processes in colloidal systems by the example of instant coffee. *Journal of Basic and Applied Research International*, 20(2), 70-83.
42. Fenter, P., & Sturchio, N. C. (2004). Mineral–water interfacial structures revealed by synchrotron X-ray scattering. *Progress in Surface Science*, 77(5-8), 171-258.
43. Verdaguier, A., Sacha, G. M., Bluhm, H., & Salmeron, M. (2006). Molecular structure of water at interfaces: Wetting at the nanometer scale. *Chemical reviews*, 106(4), 1478-1510.
44. Maccarini, M. (2007). Water at solid surfaces: A review of selected theoretical aspects and experiments on the subject. *Biointerphases*, 2(3), MR1-MR15.
45. Maurya, M., Metya, A. K., Singh, J. K., & Saito, S. (2021). Effects of interfaces on structure and dynamics of water droplets on a graphene surface: A molecular dynamics study. *The Journal of Chemical Physics*, 154(16).
46. Del Giudice, E., Tedeschi, A., Vitiello, G., & Voeikov, V. (2013, June). Coherent structures in liquid water close to hydrophilic surfaces. In *Journal of Physics: Conference Series* (Vol. 442, No. 1, p. 012028). IOP Publishing.
47. Schurr, J. M. (2013). Phenomena associated with gel–water interfaces. Analyses and alternatives to the long-range ordered water hypothesis. *The Journal of Physical Chemistry B*, 117(25), 7653-7674.
48. Elton, D. C., Spencer, P. D., Riches, J. D., & Williams, E. D. (2020). Exclusion zone phenomena in water—A critical review of experimental findings and theories. *International Journal of Molecular Sciences*, 21(14), 5041.
49. Esplandiu, M.J.; Reguera, D. and Fraxedas, J. Electrophoretic origin of long-range repulsion of colloids near water/Nafion interfaces. *Soft Matter*, 2020,16, 3717-3726.
50. Huszár, I. N., Mártonfalvi, Z., Laki, A. J., Iván, K., & Kellermayer, M. (2014). Exclusion-zone dynamics explored with microfluidics and optical tweezers. *Entropy*, 16(8), 4322-4337.
51. Ninham, B. W., Bolotskova, P. N., Gudkov, S. V., Juraev, Y., Kiryanova, M. S., Kozlov, V. A., ... & Bunkin, N. F. (2020). Formation of water-free cavity in the process of Nafion swelling in a cell of limited volume; effect of polymer fibers unwinding. *Polymers*, 12(12), 2888.
52. Pollack, G. H. (2013). Comment on “A theory of macromolecular chemotaxis” and “Phenomena associated with gel–water interfaces. analyses and alternatives to the long-range ordered water hypothesis”. *The Journal of Physical Chemistry B*, 117(25), 7843-7846.
53. Smirnov, A.N., Syroeshkin, A.V. (2004). Supratmolecular water complexes. *Ros. Chem. J.*, XLVIII(2), 125-135 (In Russ.).
54. Smirnov, A.N. Lapshin, V.B. Balshev, A.V. Lebedev, I.M. Goncharuk, V.V. Syroeshkin, A.V. (2005). Water Structure: Giant Heterophase Water Clusters. *Chemistry and Water Technology*, 27(2), 111136 (In Russian).
55. Bunkin, N. F., Suyazov, N. V., & Tsipenyuk, D. Y. (2005). Small-angle scattering of laser radiation by stable micron particles in twice-distilled water. *Quantum electronics*, 35(2), 180.
56. Sedlák, M. (2006). Large-scale supramolecular structure in solutions of low molar mass compounds and mixtures of liquids: I. Light scattering characterization. *The Journal of Physical Chemistry B*, 110(9), 4329-4338.
57. Bukatyi, V.I. Nesteryuk, P.I. (2011). The study of optical inhomogeneities (clusters) in bidistilled water with an optical method of small angles. *Polzunovsky Vestnik*, 3(1), 106-108. (In Russian).
58. Yakhno, T.A.; Yakhno, V.G. (2019). Microstructure and phase transitions of water according to optical microscopy. *Proceedings of the second all-Russian conference "Physics of aqueous solutions"*, October 17-18, 2019, Moscow, 44-45. (in Russian).
59. Yakhno, T., & Yakhno, V. (2019). A study of the structural organization of water and aqueous solutions by means of optical microscopy. *Crystals*, 9(1), 52.
60. Yakhno, T. A., & Yakhno, V. G. (2020). Study of the role of microdispersed phase of water during its transition to activated state. *Russian journal of biological physics and chemistry*.
61. Yakhno, T., Drozdov, M., & Yakhno, V. (2019). Giant water clusters: where are they from?. *International Journal of Molecular Sciences*, 20(7), 1582.
62. Yakhno, T. A., & Yakhno, V. G. (2019). Physical-chemical evolution of the dispersed phase of water when it dries. *Russian journal of biological physics and chemistry*, 4(1), 9-16.
63. Zhang, Y., Takizawa, S., & Lohwacharin, J. (2015). Spontaneous Particle Separation and Salt Rejection by

Hydrophilic Membranes. *Water Journal*.

64. Zheng, J. M., Chin, W. C., Khijniak, E., Khijniak Jr, E., & Pollack, G. H. (2006). Surfaces and interfacial water: evidence that hydrophilic surfaces have long-range impact. *Advances in colloid and interface science*, 127(1), 19-27.
65. Zheng, J. M., & Pollack, G. H. (2003). Long-range forces extending from polymer-gel surfaces. *Physical Review E*, 68(3), 031408.
66. Yakhno, T. A., Yakhno, V. G., & Zanozina, V. F. (2017, October). Phase transitions of water as a source of slow oscillatory processes in liquid media. In *Proceedings of the XII International Science-Technical Conference "Modern Trends in Biological Physics and Chemistry"*, Sevastopol, Russia (pp. 2-6).
67. Lo, S. Y., Geng, X., & Gann, D. (2009). Evidence for the existence of stable-water-clusters at room temperature and normal pressure. *Physics Letters A*, 373(42), 3872-3876.
68. Ho, M. W. (2014). Large supramolecular water clusters caught on camera-a review. *Water*, 6, 1-12
69. Seddon, J. R., & Lohse, D. (2011). Nanobubbles and micropancakes: gaseous domains on immersed substrates. *Journal of physics: Condensed matter*, 23(13), 133001.
70. Zhang, L., Zhang, X., Fan, C., Zhang, Y., & Hu, J. (2009). Nanoscale multiple gaseous layers on a hydrophobic surface. *Langmuir*, 25(16), 8860-8864.
71. Borilo, L. P. (2012). *Thin-film inorganic nanosystems* (ed. prof. V.V.Kozik). - Tomsk: Tomsk State University, 134 p. ISBN978-5-94621-362-2 (In Russian.).
72. Yakhno, T. A., & Yakhno, V. G. (2020). Study of the role of microdispersed phase of water during its transition to activated state. *Russian journal of biological physics and chemistry*.
73. Elia, V., Ausanio, G., De Ninno, A., Gentile, F., Germano, R., Napoli, E., & Niccoli, M. (2013). Experimental evidence of stable aggregates of water at room temperature and normal pressure after iterative contact with a Nafion® polymer membrane. *Water*, 5, 16-26.
74. Elia, V., Napoli, E., & Niccoli, M. (2013). Physical-chemical study of water in contact with a hydrophilic polymer: Nafion. *Journal of thermal analysis and calorimetry*, 112(2), 937-944.
75. Elia, V., Lista, L., Napoli, E., & Niccoli, M. (2014). A thermodynamic characterization of aqueous nanostructures of water molecules formed by prolonged contact with the hydrophilic polymer Nafion. *Journal of Thermal Analysis and Calorimetry*, 115(2), 1841-1849.
76. Elia, V., Germano, R., & Napoli, E. (2015). Permanent dissipative structures in water: the matrix of life? Experimental evidences and their quantum origin. *Current topics in medicinal chemistry*, 15(6), 559-571.
77. Elia, V., Yinnon, T. A., Oliva, R., Napoli, E., Germano, R., Bobba, F., & Amoresano, A. (2017). Chiral micron-sized H<sub>2</sub>O aggregates in water: circular dichroism of supramolecular H<sub>2</sub>O architectures created by perturbing pure water. *Water*, 8, 1-29.
78. Yinnon, T. A., Elia, V., Napoli, E., Germano, R., & Liu, Z. Q. (2016). Water ordering induced by interfaces: an experimental and theoretical study. *Water*, 7, 96-128..
79. Elia, V.; Napoli, E.; Germano, R.; Roviello, V.; Oliva, R.; Niccoli, V.; Amoresano, A.; Toscanesi, M.; Trifuoggi, M.; Fabozzi, A.; Yinnon, T.A. Water perturbed by cellophane: comparison of its physicochemical properties with those of water perturbed with cotton wool or Nafion. *Journal of Thermal Analysis and Calorimetry*, 2020. <https://doi.org/10.1007/s10973-020-10185-0>
80. Elia, V., Napoli, E., Germano, R., Roviello, V., Oliva, R., Niccoli, M., ... & Yinnon, T. A. (2021). Water perturbed by cellophane: comparison of its physicochemical properties with those of water perturbed with cotton wool or Nafion. *Journal of Thermal Analysis and Calorimetry*, 146(5), 2073-2088.
81. Loelovich, M. Ya. (2016). Models of supermolecular structure and properties of cellulose. *High-molecular compounds, Ser. A*, 58(6), 604-624. (In Russian).
82. Gallyamov, M. O., Khokhlov, A., R. (2014). *Polymer Membrane Fuel Cells: Materials for the Fuel Cell Fundamentals course*. Moscow: Faculty of Physics, Moscow State University, 72 p. ISBN 9785-600-00346-0 (In Russian).
83. Del Giudice, E., Preparata, G., & Vitiello, G. (1988). Water as a free electric dipole laser. *Physical review letters*, 61(9), 1085.
84. Yinnon, T. A., & Liu, Z. Q. (2015). Domains Formation Mediated by Electromagnetic Fields in Very Dilute Aqueous Solutions: 3. Quantum Electrodynamical Analyses of Experimental Data on Solutions of Weak Electrolytes and Non-electrolytes. *Water Journal*.
85. Bier, M., & Pravica, D. (2018). LIMITS ON QUANTUM COHERENT DOMAINS IN LIQUID WATER. *Acta Physica Polonica B*, 49(9).
86. Messori, C., Prinzer, S. V., & di Bardone, F. B. (2019). Deep into the water: exploring the hydro-electromagnetic and quantum-electrodynamical properties of interfacial water in living systems. *Open Access Library Journal*, 6(05), 1.

**Copyright:** ©2025 T. A. Yakhno, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.