

## Imaging of Mitral Valve Area by Cardiac Magnetic Resonance in Patients with Rheumatic Mitral Stenosis

Ahmed Ibrahim Bedier<sup>1\*</sup> and Amal Sakrana<sup>2</sup>.

<sup>1</sup>Cardiovascular medicine department, Mansoura University

<sup>2</sup>Diagnostic and interventional radiology department, Mansoura University

### \*Corresponding author

Ahmed Ibrahim Bedier, Cardiovascular medicine department, Mansoura University

Submitted: 15 Jul 2022; Accepted: 22 Jul 2022; Published: 30 Jul 2022

**Citation:** Ahmed Ibrahim Bedier, (2022). Imaging of Mitral Valve Area by Cardiac Magnetic Resonance in Patients with Rheumatic Mitral Stenosis, *World J Clin Med Img*, 1(1), 14-18.

### Abstract

**Background:** Rheumatic heart disease which is a result of rheumatic fever is still a major health problem in developing countries. Rheumatic mitral stenosis (MS) is the commonest delayed valvular affection as a consequence of rheumatic fever. The assessment of MS severity by measuring mitral valve area (MVA) is very essential for patient management. Different imaging modalities are available for MVA assessment including echocardiography and cardiovascular magnetic resonance (CMR).

**Objectives:** The aim of this study is to compare the planimetric MVA between CMR and two dimensional echocardiography in MS patients.

**Patients and methods:** A forty adult patients with symptomatic mitral stenosis were included in the study. Other significant valvular lesions, atrial fibrillation, poor echocardiographic window, contraindications to CMR, and NYHA IV were excluded from the study. All patients were assessed by 2D echocardiography and CMR for MVA measurement.

**Results:** The mean 2D TTE MVA was  $1.2 \pm 0.26 \text{ cm}^2$ , while the mean CMR MVA  $1.2 \pm 0.28 \text{ cm}^2$ . No significant statistical difference was found between both methods ( $P$  value 0.842) with a very strong correlation between both methods ( $r = 0.93$  and  $p$ -value  $< 0.0001$ ). The mean difference of MVA between the two methods was  $0.012 \text{ cm}^2$ .

**Conclusion:** CMR is a non-invasive imaging modality that provides MVA measurement and is a reliable method in the diagnosis of MS patients.

### Introduction

Rheumatic heart disease which is a result of rheumatic fever that is caused by group A beta hemolytic streptococci, is still a major health problem in developing countries. Although the health related burden of rheumatic heart disease has been declined worldwide, it is still considered a cause of young population morbidity and mortality [1-3].

Rheumatic mitral stenosis (MS) which is defined as diastolic narrowing of the mitral valve (MV) orifice is the commonest delayed valvular affection as a consequence of rheumatic fever. Characteristic changes that occur in mitral valve as a result of rheumatic fever are leaflet edges thickening, commissural fusion, and chordal thickening, shortening and fusion [4].

Mitral stenosis diagnosis and assessment of its severity are very essential for timing and selection of the method of treatment. Echocardiography represents the corner stone for diagnostic

assessment in patients with MS by assessment of mitral valve area (MVA) and is also helpful in selection of patients for valvuloplasty [5].

Although echocardiography is the main imaging modality used to assess MS severity, cardiovascular magnetic resonance (CMR) also may be performed as a complementary non-invasive technique especially, when TTE and TEE evaluations are of suboptimal quality and in whom Doppler studies are inconsistent with the clinical data. CMR has the advantage of being non-invasive, reproducible, not limited by air and bone conduction, more sensitive in detecting thrombus, and has a high 3D spatial resolution and freedom of access to any location in any position. [6].

To date, no studies have been done on the role of CMR in evaluating rheumatic mitral stenosis (MS) in Egypt and comparing the results with echocardiographic results.

## Objective of the study

The aim of this study is to compare the planimetric MVA between CMR and two dimensional echocardiography in patients with MS.

## Patients and methods

This prospective study was carried out between January 2019 and January 2020 on forty adult patients with severe symptomatic MS admitted to Specialized Medical Hospital, Cardiovascular Medicine Department, Mansoura University and referred to Magnetic Resonance Unit, Radiology Department, Mansoura University Hospital for CMR. Patients with other significant valvular lesions, atrial fibrillation, poor echocardiographic window, contraindications to CMR, and NYHA IV were excluded from the study.

Written consent was obtained from all patients and the study was accepted by IRB committee. A detailed history including age, sex, residence, NYHA class, history of previous commissurotomy or PBMV, antibiotic prophylaxis, anticoagulants, and previous history of TIA, stroke or embolization. A detailed physical examination was done also.

The study included 40 patients, 5 males (12.5%) and 35 females (87.5%). The age ranged from 24 to 60 years with a mean of  $37.3 \pm 8.4$  years. As regard presence of other medical disease; 38 patients (95%) had no medical diseases while one patient (2.5%) had diabetes mellitus and another one had chronic kidney disease (2.5%). All patients were from rural areas in Dakahlia and neighboring governorates.

**Table 1: Demographic data of studied cases**

Age/y	Mean $\pm$ SD (min-max)	
	37.3 $\pm$ 8.4 (24-60)	
Sex	(N%)	
Male	5	12.5
Female	35	87.5
Medical disease		
No	38	95
Yes DM	1	2.5
Yes CKD	1	2.5
Residence		
Rural	40	100

## Echocardiography

was done by (GE vivid E9 XDClear, GE Medical Systems, General Electric Company, Manufacturer GE Vingmed Ultrasound AS, Horten, Norway), with the probe (4V Hz). Assessment of mitral valve morphology was done in parasternal long axis, short axis and Apical views, with evaluation of subvalvular apparatus, MV leaflets mobility, thickness and calcification. Assessment of mitral valve area (MVA) by MV planimetry was performed in short-axis view, in the diastolic frame with maximum diastolic opening of the MV, with identifying the smallest orifice at the leaflet tips. Estimation of MVA less than 1.5 cm<sup>2</sup> is considered an indication for intervention according to the latest ESC guidelines (7).

## CMR assessment

CMR was done by a clinical MR scanner 1.5-T scanner, Philips, Ingenia. During CMR study the procedure was explained to the patient with training for breath hold technique with continuous monitoring of the patient heart rate. A 16-channel torso phased-array receiver coil was used for signal reception. All acquisition of data was retrospective ECG gated and with respiratory gating. The scan protocol was carried out in the following order:

- A Scout images (axial, coronal, sagittal) using Real-time interactive planning (FOV 450 X 450, mm<sup>2</sup>, slice thickness 10 mm acquisition matrix 220 x 176, voxel size 1.6 mm x 1.9 mm x 10 mm, echo/repetition time (TE/TR) shortest, and flip angle 50°).
- Cine steady state free precession (SSFP) sequences were acquired on the long axis cardiac planes: 4 chambers, 2 chambers, and 3 chambers (FOV 350 x 350 mm<sup>2</sup>, slice thickness 6-8 mm, acquisition matrix 220 x 176, voxel size 1.7 mm x 1.7 mm x 10 mm, and flip angle 60°), followed by a "stack" of contiguous SSFP cine images, with the same technical parameters, acquired along cardiac short axis, to cover the whole LV from base to apex.
- The MV was visualized with 4 chamber, 2 chamber and 3 chamber views. Short axis LV images were performed parallel to the mitral valve plane; 4-6 cross sections were obtained (with the same previous parameters but with slice thickness of 5 mm, with 5-6 slices, with slice gap -1). Then the minimal diastolic area was chosen as the planimetric MVA.
- Depending on the heart rate, and patient ability to hold breath the average scanning time was 10 - 20 minutes.

## CMR imaging analysis

- Cardiac morphology and function were quantitatively evaluated on the cine images with the workstation (Circle CMR cvi42 cardiovascular imaging Inc 2016, Calgary, Canada). Two CMR radiologists assessed the CMR findings independently and parameters were recorded. The CMR images were assessed for mitral valve morphology, motion, and thickness. Calculation of planimetric MVA was done by measuring the smallest orifice at mid diastole in different slices.
- All CMR data were blinded to echocardiographic data.

## Statistical analysis

The history, 2D data, and CMR data were recorded on an investigation report form, and tabulated, coded then analyzed. All statistical analyses were performed using SPSS (statistical package of social sciences) version 22 for windows (SPSS Inc., Chicago, IL, USA). Normality of data was first tested by Shapiro-Wilk test. Parametric data were presented as mean  $\pm$  standard deviation (SD), while non parametric data were expressed in median, minimum and maximum. Categorical data were presented as absolute numbers and percentages (%). Correlation analysis between echocardiography and CMR was done by Pearson coefficient of correlation test. Scatter plot graphs were used to represent the significant correlation. A 2 tailed P-values  $\leq$  0.05 were considered to be statistically significant, and  $\leq$  0.001 were considered to be statistically highly significant. The smaller the p-value obtained, the more significant are the results.

## Results

The study included forty MS patients; thirty three of them were NYHA II 33 (82.5%), while 6 patients (15%) were NYHA III, and only one patient was NYHA I (2.5%). Only one patient had history of closed commissurotomy (2.5%), and 7 cases had previous PBMV (17.5%). Only one patient had history of TIA (2.5%). 5 patients were on anticoagulation therapy (12.5%) while 35 patients were not on anticoagulation therapy (87.5%). 36 patients were on penicillin antibiotic prophylaxis (90%), while 4 patients did not receive penicillin antibiotic prophylaxis (10%).


**Table 2: patient clinical characteristics**

Variable	Mean ±SD	
BSA	1.94±0.14	
NYHA	(N%)	
Class I	1	2.5
Class II	33	82.5
Class III	6	15
<b>Previous commissurotomy</b>		
No	39	97.5
Yes,24y	1	2.5
<b>Previous PBMV</b>		
No	33	82.5
Yes	7	17.5
<b>Antibiotic prophylaxis</b>		
Yes	36	90
No	4	10
<b>Oral anticoagulants</b>		
Yes	5	12.5
No	35	87.5
<b>Previous stroke or TIA</b>		
No	39	97.5
Yes	1	2.5

**Table 3: Mitral valve area by direct planimetry in all studied cases**

		2D TTE	CMR
N	Valid	40	40
	Missing	0	0
Mean		1.20	1.20
Median		1.20	1.20
Std. Deviation		0.26	0.28
Minimum		0.70	0.62
Maximum		1.80	1.80

There was a significant positive correlation between both modalities in direct MVA planimetry measurement (P value < 0.0001). The correlation had a high linearity between both modalities (r > 0.9).



**Figure 1: The scatter plot graph of the planimetric MVA using CMR and 2D TTE.**

## Discussion

Direct planimetry of mitral valve area is the corner stone for diagnosis of mitral stenosis and for decision making as presence of a symptomatic patient with MVA less than 1.5 cm<sup>2</sup> is an indication for intervention either with MV replacement or PBMV. Planimetric MVA can be assessed by different modalities. In our study we assessed MV by 2D TTE, and CMR. In our study; the mean 2D TTE MVA was 1.2±0.26 cm<sup>2</sup> (range 0.7-1.8 cm<sup>2</sup>), while the mean CMR MVA 1.2±0.28 cm<sup>2</sup> (range 0.62-1.8 cm<sup>2</sup>). No statistically significant difference in MVA was found between the 2D TTE and CMR (P value 0.842) with a very strong correlation between both methods (r = 0.93 and p-value < 0.0001). The mean difference of MVA between both methods was 0.012 cm<sup>2</sup>, with slight overestimation of CMR MVA by 0.12%. In our study we observed the advantage of CMR in patients with heavily calcific valves; as MVA can be measured easily without limitation as in echocardiography (Figure 2).

This significant correlation between 2D TTE, and CMR in this study is concordant with a study done by Mutnuru, et al; who performed 2DTTE and CMR on 50 Indian patients with different rheumatic valvular affection with MS was the predominant valve affection in the study. They found that the mean MVA by 2D TTE was 1.79±0.43 cm<sup>2</sup> and by CMR 1.82±0.47 cm<sup>2</sup> (r= 0.98, p-value <0.00001) and they described a highly significant strong positive association between the results by 2D TTE and CMR [8].

Our results are also concordant with another study done by Kim et al on 102 MS patients with AF; where they compared MVA by TTE and PHT with MVA by CMR and MSCT. The mean MVA by 2D TTE was 1.16 ± 0.28 cm<sup>2</sup> and CMR 1.15 ± 0.28 cm<sup>2</sup>; the correlation between CMR and TTE planimetry (r = 0.67, P <0.05). The mean difference between the 2D TTE and CMR MVAs was 0.01 (P = 0.61) with an overestimation of 0.9 % by CMR [9].


Another CMR study done by Helvacioğlu, et al; who performed 2DTTE and CMR on 30 patients with rheumatic mitral stenosis

in Turkey; the mean 2D TTE MVA in their study was  $1.50 \pm 0.53 \text{ cm}^2$  and the mean CMR MVA was  $1.50 \pm 0.26 \text{ cm}^2$ . No statistically significant difference in MVA was found between the 2D TTE and CMR ( $P = 0.90$ ) [6]. A very strong correlation was found also between both methods in MVA assessment ( $r = 0.971$ ,  $P < 0.0001$ ). The mean difference of MVA between both methods was  $0.018 \text{ cm}^2$ .

Our results are also concordant with another study done by Lanjewar et al on 30 patients with rheumatic mitral stenosis in India. The mean CMR MVA, was  $1.71 \pm 0.44 \text{ cm}^2$  (range: 0.5 - 2.40  $\text{cm}^2$ ); and the mean MVA by 2D TTE,  $1.31 \pm 0.30 \text{ cm}^2$  (range: 0.6 to 2.50  $\text{cm}^2$ ) [10]. The correlation between planimetric MVA by CMR and 2D TTE was very good ( $r = 0.81$ ,  $p < 0.05$ ). The

mean absolute difference between both methods was  $0.12 \pm 0.23 \text{ cm}^2$  ( $p < 0.05$ ), with a slight overestimation (by 7.6%) of the CMR MVA when compared to 2D TTE MVA.

Our study results and the previous studies mentioned before are also in line with a study done by Djavidani, et al; where they compared MVA by CMR, 2D TTE, and invasively by Gorlin-formula at the catheterization laboratory in 22 patients with mitral stenosis in Germany [11]. The correlation between planimetric CMR MVA and 2 D TTE MVA was very good ( $r = 0.81$ ,  $p < 0.0001$ ). The mean absolute difference between CMR MVA and 2 D TTE MVA was  $0.13 \pm 0.24 \text{ cm}^2$  ( $p < 0.05$ ), resulting in a slight overestimation of CMR MVA as compared with 2D TTE MVA by 8.1%.



**Figure 2:** A sixty years old man with symptomatic MS. MV leaflets were severely calcified by TTE parasternal long axis view (a) and parasternal short axis view (b) at mid diastole. (c) CMR can clearly identify the tips and measure MVA, which was  $0.9 \text{ cm}^2$ .

### Limitation of the study

The small number of patients included in this study is considered a limitation, but other studies were done on also a smaller number of patients.

Exclusion of AF in the study patients, however it is more common in MS patients, but the aim of exclusion of AF patients was to avoid averaging of measurements which may lead to bias.

We didn't compare our MVA results with Gorlin Formula (the gold standard method for MVA), because this invasive method is not done routinely for MS patients and restricted only in special circumstances where there is discrepancies between echocardiographic measurements and clinical status. However the Gorlin method is invasive and has several pitfalls and technical limitations [12,13].

We used the conventional 2D planimetry method is the reference method because other Doppler methods including PHT, PISA, and continuity equation have their own limitations [14,15].

### Recommendations

CMR is a non-invasive imaging modality that can be used as an alternative to TTE in cases with poor acoustic windows or in calcific mitral valves.

### Conclusion

Cardiovascular magnetic resonance is a non-invasive imaging  
*World J Clin Med Img, 2022*

modality that can provide a reliable assessment of MVA with comparable results with echocardiography.

### References

1. Woldu, B., & Bloomfield, G. S. (2016). Rheumatic heart disease in the twenty-first century. *Current Cardiology Reports*, 18(10), 1-11.
2. Watkins, D. A., Johnson, C. O., Colquhoun, S. M., Karthikeyan, G., Beaton, A., Bukhman, G., ... & Roth, G. A. (2017). Global, regional, and national burden of rheumatic heart disease, 1990–2015. *New England Journal of Medicine*, 377(8), 713-722.
3. Lilly, L. S. (2012). *Pathophysiology of heart disease: a collaborative project of medical students and faculty*. Lippincott Williams & Wilkins.
4. La Canna, G. (2016). Heart valve disease (mitral valve disease): anatomy and morphology of the mitral valve. *The EACVI Textbook of Echocardiography*, 267.
5. Helvacioğlu, F., Yildirimtürk, O., Duran, C., Yurdakul, S., Tayyareci, Y., Ulusoy, O. L., & Aytekin, S. (2014). The evaluation of mitral valve stenosis: comparison of transthoracic echocardiography and cardiac magnetic resonance. *European Heart Journal—Cardiovascular Imaging*, 15(2), 164-169.
6. Vahanian, A., Baumgartner, H., Bax, J., Butchart, E., Dion, R., Filippatos, G., ... & Tamargo, J. (2007). Guidelines on the management of valvular heart disease: The Task Force on the Management of Valvular Heart Disease of the European Society of Cardiology. *European heart journal*, 28(2), 230-268.



7. Mutnuru, P. C., Singh, S. N., D'Souza, J., & Perubhotla, L. M. (2016). Cardiac MR imaging in the evaluation of rheumatic valvular heart diseases. *Journal of Clinical and Diagnostic Research: JCDR*, 10(3), TC06.
8. Kim, S. S., Ko, S. M., Song, M. G., Chee, H. K., Kim, J. S., Hwang, H. K., & Lee, J. H. (2015). Quantification of stenotic mitral valve area and diagnostic accuracy of mitral stenosis by dual-source computed tomography in patients with atrial fibrillation: comparison with cardiovascular magnetic resonance and transthoracic echocardiography. *The International Journal of Cardiovascular Imaging*, 31(1), 103-114.
9. Charan Lanjewar, B. E., Mishra, N., Jhankariya, B., & Kerkar, P. (2010). Planimetry of mitral valve stenosis in rheumatic heart disease by magnetic resonance imaging. *The Journal of heart valve disease*, 19(3), 357-363.
10. Djavidani, B., Debl, K., Lenhart, M., Seitz, J., Paetzel, C., Schmid, F. X., ... & Luchner, A. (2005). Planimetry of mitral valve stenosis by magnetic resonance imaging. *Journal of the American College of Cardiology*, 45(12), 2048-2053.
11. Hammermeister, K. E., Murray, J. A., & Blackmon, J. R. (1973). Revision of Gorlin constant for calculation of mitral valve area from left heart pressures. *British Heart Journal*, 35(4), 392.
12. Klarich, K. W., Rihal, C. S., & Nishimura, R. A. (1996). Variability between methods of calculating mitral valve area: simultaneous Doppler echocardiographic and cardiac catheterization studies conducted before and after percutaneous mitral valvuloplasty. *Journal of the American Society of Echocardiography*, 9(5), 684-690.
13. Flachskampf, F. A., Weyman, A. E., Gillam, L., Chun-Ming, L., Abascal, V. M., & Thomas, J. D. (1990). Aortic regurgitation shortens Doppler pressure half-time in mitral stenosis: clinical evidence, in vitro simulation and theoretic analysis. *Journal of the American College of Cardiology*, 16(2), 396-404.
14. Fredman, C. S., Pearson, A. C., Labovitz, A. J., & Kern, M. J. (1990). Comparison of hemodynamic pressure half-time method and Gorlin formula with Doppler and echocardiographic determinations of mitral valve area in patients with combined mitral stenosis and regurgitation. *American heart journal*, 119(1), 121-129.

**Copyright:** ©2022 Ahmed Ibrahim Bedier. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.