

Incidence of Trichomonas Tenax in Diseased Mouth

Hala Nadhim Kadhimi¹ and Nadham Kadham Mahdi^{2,*}¹University of Al-Watania, Thi-Qar, Iraq²Department of Microbiology, College of Medicine, University of Basrah, Iraq***Corresponding Author**

Dr. Nadham K Mahdi, Department of Microbiology, College of Medicine, University of Basrah, Central Post Office-42001, P. O. Box 1565, Ashar, Basrah, Iraq.

Submitted: 2023 June 19; **Accepted:** 2023 July 14; **Published:** 2023 July 20**Citation:** Kadhimi, H.N. & Mahdi, N.K. (2023). Incidence of Trichomonas Tenax in Diseased Mouth. *J Oral Dent Health*, 7(3), 198-200.**Abstract****Objective:** Is to report the incidence of *T. tenax* in relation to oral health, age, sex and residency.**Materials and Methods:** Two hundreds patients with diseased mouths were examined by wet mount preparation and Giemsa's stained methods for identification of *Trichomonas tenax*.**Results:** Its frequency was 9% among the examined swabs. There were 50% infection rate for both males as well as females. The highest incidence (11.25%) was found among 21-40 years old while the lowest (6.66%) in 41-60 years old. There were no significant variation in relation to residency. The association between *T. tenax* infection and dental caries was noticed at a rate of 66.7%. In addition, different rates for *T. tenax* were observed among patients with some other oral diseases as bleeding gum, gingivitis and periodonitis.**Conclusion:** This parasite should be considered as a potential etiological agent in diseased mouth, especially in individuals with poor oral hygiene.**Keywords:** Incidence, Mouth diseases, Trichomonas tenax**1. Introduction**

Trichomonas tenax is a non-pathogenic oral protozoan parasite of human being. It has a worldwide distribution and may be found in upto 26% of patients with dental caries or pyorrhea and in upto 11% of those healthy mouths [1]. There were studies that related to its incidence in patients with chronic periodontitis [2-4]. Transmission is through saliva, air droplet spray, and kissing or use of contaminated dishes and drinking water [1].

Several studies have reported its incidence in the world including Iraq (20%), Iran, (20.7%), Turkey (2.17%), Egypt (19%), Malaysia (32%), Nigeria (35%), Italy (40%), Hungary (38.3%) and France (35.5%) [3,5-13].

The aim of the study is to record the incidence of *T. tenax* in relation to oral health, age, sex and residency.

2. Patients and Methods

A sterile swab was rubbed around the surface of teeth for 200 patients with diseases mouth. Samples were collected during 3 months (June-August, 2022). This study was carried out at College of Dentistry, College of Medicine and private clinics,

Basrah, Iraq. Informed consent has been obtained from all involved patients. The work has proved by the Ethical Committee of the College of Medicine, Basrah, Iraq (No. 2022-371). All patients provided informed consent. There were 100 males and 100 females. Their ages ranged between 3 – 60 years. People receiving anti-protozoan drugs were excluded from the study.

The collected samples were examined directly by wet mount preparation and Giemsa's stained methods [14].

3. Results

Trichomonas tenax was found in 18(9.0%) patients with diseased mouths (Table 1). Its frequency (11.25%) was noticed in a substantial level among age group of 21-40 years (Table 1). There were equal incidence rates for both males and females. Furthermore, there was no significant variation as far as residency is concerned.

The association between *T. tenax* infection and dental caries was noticed at a rate of 66.7%. In addition, different rates for *T. tenax* were observed among patients with some other oral diseases as bleeding gum, gingivitis and periodonitis (Table 2).

Variable (years)	No. examined	No. (%) positive
Age:		
3-20	105	8 (7.6)
21-40	80	9 (11.25)
41-60	15	1(6.66)
Sex:		
Male		
Female	100	9 (50)
8 (7.6)	100	9 (50)
Residency:		
Urban	120	11 (9.16)
Rural	80	7 (8.75)
Total	200	18 (9.0)

Table 1: Incidence of *Trichomonas tenax* according to different variables

Oral disorders	<i>Trichomonas tenax</i> , No (%)
Dental caries	12 (66.7)
Bleeding gum	3 (16.7)
Gingivitis	2 (11.1)
Periodonitis	1 (5.5)
Total	18 (100)

Table 2: The association between *Trichomonas tenax* positivity and oral disorders

4. Discussion

The peak of infection (11.25%) was detected in patients of 21-40 years of age. Similar findings with higher incidence rates have been reported elsewhere but dissimilar with other study [5,9,15-17]. The prevalence of parasitic infection among population may be attributed to lack of health and cultural awareness, lack of attention to oral and dental hygiene, lack of adequate hygiene guidance, and the spread of bad habits such as the use of matchstick and other tooth cleaning products that lead to increased oral parasites.

However, the low incidence observed in this study can be due to a better degree of oral hygiene, teeth brushing and proper restorations of decayed teeth and periodontal problems. In addition, the decrease or absence of *T. tenax* in older people may be related to the unfavorable conditions for its existence for instance, in toothless mouths, or due to educational level factor to maintain better oral hygiene and dental care.

Both sexes showed an equal incidence rate for *T. tenax*. This result is in agreement with previous study at the same locality [5]. Other workers have stated a high incidence among males than females while vice versa have reported a higher rates among females rather males [9,15-17]. The explanations for that are all related to the practice degree for oral health including teeth.

Present study hasn't shown significant effect of residence. Similarly, in Iran have not seen significant differences as far as residence is concerned [17]. Teeth brush is an excellent practice to get a healthy clean mouth and teeth to prevent suitable

conditions for the growth and survival of *T.tenax* [9].

5. Conclusion

In conclusion, this parasite should be considered as a potential etiological agent in diseased mouth, especially in individuals with poor oral hygiene.

Author Contributions: N.K. Mahdi, diagnosis of the protozoan parasite and wrighting up. H.N. Kadhim, sampling and diagnosis of the oral diseases.

Declaration of Interests: The authors have no conflicts of interest to declare.

Funding: The authors declare that this work has received no financial support.

References

1. Soulaby, R.J.L. (1969). Helminth, arthropoda and protozoa of domesticated animals. 2nd ad. London Bailliere, Tindall and Caaell, p. 588.
2. Sarowska, J., Wojnicz, D., Kaczkowski, H. & Jankowski, S. (2004). The occurrence of Entamoeba gingivalis and Trichomonas tenax in patients with periodontal disease. Adv Clin Exp Med, 13(2), 291-297.
3. Athari, A., Soghandi, L., Haghighi, A. & Kazemi, B. (2007). Prevalence of Oral trichomoniasis in Patients with Periodontitis and Gingivitis Using PCR and Direct Smear. Iran. J Publ Health, 36(3), 33-37.
4. Bisson, C., Dridi, SM. & Machouart, M. (2019). Assessment

- of the role of *Trichomonas tenax* in the etiopathogenesis of human periodontitis: A systematic review. *PLoS ONE*, 14(12), e0226266.
5. Mahdi, N.K., AL-Saeed, AT. (1993). *Trichomonas tenax* in Basrah Iraq. *J Pakistan Med Assoc*, 43(12), 261-263.
 6. Ibrahim, S. & Abbas, R. (2012). Evaluation of *Entamoeba gingivalis* and *Trichomonas tenax* in patients with periodontitis and gingivitis and its correlation with some risk factors. *J Bagh Coll Dentistry*, 24, 158-162.
 7. Gharavi, M.J., Hekmat, S., Ebrahimi, A. & Jahani, MR. (2006). Buccal cavity protozoa in patients referred to the Faculty of Dentistry in Tehran, Iran. *Iranian J Parasitol*, 1, 43-46.
 8. Abualqomsaan, M., Töz, S.O., Yolasiğmaz, A. & Turgay, N. (2010). The investigation of *Entamoeba gingivalis* and *Trichomonas tenax* in a group of patients with periodontal disease. *Turkiye Parazitoloji Dergisi*, 34, 91-94.
 9. Hamadto, H.H.A., Ibrahim, A.H., El-Hayawan, F., Abdallah, I. & Abd El-Maboud, O.I., et al. (2014). Relation between *Trichomonas tenax* and pulmonary diseases. *The Egyptian Journal of Medical Sciences*, 35(2), 633-652.
 10. Honigherg, B.M. (1971). *Trichomonada* of Importance in Human Medicine. In *Parasitic Protozoa*, Vol II (J.P. Kreier ed.). Academic Press, New York, San Francisco and London.
 11. Onyido, A.E., Amadi, E.S., Olofin, I., Onwumma, A.A. & Okoh, I.C, et al. (2011). Prevalence of *Entamoeba gingivalis* and *Trichomonas tenax* among dental patients attending Federal School of Dental Technology and Therapy clinic, Enugu, Nigeria. *Nature and Science*, 9(9), 59-62.
 12. Matteo, F., Viganò, L. & Casu, C. (2018). *Trichosoma tenax* and *Entamoeba gingivalis*: pathogenic role of protozoic species in chronic periodontal disease development. *J Human Virology and Retrovirology*, 6(1), 81-82.
 13. Feki, A. & Molet, B. (1990). Importance of *Trichomonas tenax* and *Entamoeba gingivalis* protozoa in the human oral cavity. *Rev. Odontostomatol*, 19(1), 37-45.
 14. John, D.T., Petri, W.A., Markell. & Voge. (2006). *Medical Parasitology*. 9th Ed. Philadelphia, Saunders, pp. 421-422.
 15. Bracamonte-Wolf, C., Orrego, PR., Muñoz, C., Herrera, D. & Bravo, J., et al. (2019). Observational cross-sectional study of *Trichomonas tenax* in patients with periodontal disease attending a Chilean University dental clinic. *BMC Oral Health* 19, 207-216.
 16. Mahmmmed, S.A. & Al-Waaly, A.B.M. (2019). Prevalence of *Trichomonas tenax* in Karbala Governorate. 2nd International Science Conference IOP Conf. Series: Journal of Physics: Conf. Series 1294.
 17. Mahmoudvand, H., Sepahvand, A., Niazi, M., Momeninejad, N. & Sepahvand, S.M, et al. (2018). Prevalence and risk factors of oral cavity protozoa (*Entamoeba gingivalis* and *Trichomonas tenax*) among patients with dental cavity caries. *Journal of Research in Medical and Dental Science*, 6(5), 42-46.

Copyright: ©2023 Dr. Nadham K Mahdi, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.