

New Representation of the Euler Constant "e"

Thomas Clapies* 

Independent Researcher, France

*Corresponding Author

Thomas Clapies, Independent Researcher, France.

Submitted: 2026, May 11; Accepted: 2026, Jun 01; Published: 2026, Jun 12

Citation: Clapies, T. (2026). New Representation of the Euler Constant "e". *J Electrical Electron Eng*, 5(3), 01-02.

Abstract

This article reminds us that certain fundamental constants are associated with functions, in particular the base of the exponential function. We use certain remarkable identities associated with the exponential function and Lambert's function W (also called Product log) to obtain a new representation of the Euler constant $e = \exp(1)$ using a function [1,2].

Demonstration

According to Euler's equation

$$e^{i\pi} = -1 = i^2$$

we deduce that

$$\pi = -2i \ln i$$

$$\text{And } {}^iW_0\left(-\frac{\pi}{2}\right) = i\frac{\pi}{2}$$

therefore, relationship (1)

$$i = \frac{-\pi/2}{W_0\left(-\frac{\pi}{2}\right)} = \frac{2}{\pi} W_0\left(-\frac{\pi}{2}\right)$$

with W_0 the Lambert function (the main branch of the Product log). It is also accepted that

$$i^i = e^{-\frac{\pi}{2}}$$

therefore, relation (2) :

$$e = \left(e^{-\frac{\pi}{2}}\right)^{-\frac{2}{\pi}} = i^{-\frac{2i}{\pi}} = i^{-\frac{2i}{2i \ln i}} = \frac{1}{i \ln i}$$

Since pi and e are related to i, we deduce that e is related to pi: (1) combined with (2)

$$e = -\frac{\pi}{2W_0\left(-\frac{\pi}{2}\right)} \frac{1}{\ln\left(-\frac{\pi}{2W_0\left(-\frac{\pi}{2}\right)}\right)}$$

In addition, we can see (graphically, or by technically complex calculation based on the derivation properties of the W and ln functions) that

$$\forall x \in \mathbb{R}, \quad \frac{d}{dx} \left(-\frac{x}{2W_0\left(-\frac{x}{2}\right)} \frac{1}{\ln\left(-\frac{x}{2W_0\left(-\frac{x}{2}\right)}\right)} \right) = 0$$

so

$$\forall x \in \mathbb{R}, f(x) = -\frac{x}{2W_0\left(-\frac{x}{2}\right)} \frac{1}{\ln\left(-\frac{x}{2W_0\left(-\frac{x}{2}\right)}\right)} = e$$

to be compared, for example, with long-established trigonometric functions such asii

$$g(x) = \cos x^2 + \sin x^2 = 1, \forall x \in \mathbb{R}_{hG}$$

$$h(x) = \tan^{-1} \frac{1}{x} + \tan^{-1} x = \frac{\pi}{2}, \forall x \in \mathbb{R}^+$$

References

1. Lambert W function – Wikipedia (special values)
2. Tangent arc - Wikipedia (wikipedia.org)

Copyright: ©2026 Thomas Clapies. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.