

## Palliative Care and Surgery

Sinisa Franjic\*

Independent Researcher

\*Corresponding Author

Siniša Franjić, Independent Researcher

Submitted: 2025, Dec 01; Accepted: 2026, Jan 02; Published: 2026, Jan 16

**Citation:** Franjic Sinisa. (2026). Palliative Care and Surgery. *Int J Cancer Res Ther*, 11(1). 01-05.

### Abstract

Palliative care is a special type of medical care. It is pointed at progressing the quality of life of patients enduring from serious illnesses. It makes a difference them and their families to adapt more effectively with the issues that such diseases bring. The fundamental objective of palliative care is to make the quiet feel way better. In this manner, it anticipates or treats the indications and side impacts of the illness and treatment, but the accentuation is on the passionate, social, commonsense and otherworldly issues that the infection can cause. Palliative surgical strategies in oncological patients are those that dispense with complications that are a result of harmful disease. Their fundamental reason is to progress the quality of life of patients, and in some cases draw out survival. Palliative surgical strategies must be carefully weighed between the conceivable benefits of the anticipated result of the method itself, but moreover the advancement of complications.

**Keywords:** Palliation, Palliative Care, Surgery, Patient, Health

### 1. Introduction

The word “palliative” comes from a Latin word palliat, which means “cloaked”; as the root word suggests, palliative care points to alleviate side effects that cause pain and enduring when source control may not be conceivable [1]. The express “Palliative care” was to begin with named in 1973 by Dr. Balfour Mount—the Canadian specialist frequently credited as the father of palliative care in North America—to be utilized instep of “Hospice” which had a negative intention. Whereas the term is of later development, the concept of palliative care dates back to early medicine when the center, in the nonappearance of compelling corrective approaches, was basically on easing pain and enduring. It is as it were a cutting edge conception that accomplishing a remedy is of sole importance—enabled by uncommon mechanical progresses in therapeutic, surgical, and basic care therapies—as caring for patient’s consolation has been moderately de-emphasized.

#### 1.1. Frailty

Frailty is a geriatric disorder signifying misfortune of physical and cognitive save for which numerous scales and apparatuses have been created to evaluate [2]. Records of side effects, clutters, and physical confinements such as automatic weight misfortune, self-

reported fatigue, action level, hold quality, and strolling speed are utilized to offer assistance decide the level of slightness. Whereas there are different measurements to evaluate slightness, in any case of how it is measured the nearness of preoperative slightness has been connected with expanded length of remain, chance of complications, and postoperative mortality. With the understanding of the imperatives that expanding feebleness has on favorable results, particular care must be made to portray the objectives of care for the quiet through the continuum of their care. Patients’ wants for the sorts of treatments to get may alter as the quiet moves to the diverse stages of their care. It is basic to guarantee the patient’s values and inclinations stay at the center of the decision-making handle. Results that require to be surveyed, and re-assessed as the patient’s condition advances, are long-term side effects, utilitarian status, living area, and certainly probability of survival. It is officeholder on the intense care specialist to adjust the treatment arrange with the patient’s generally health care goals. Undoubtedly the intense care specialist will experience patients with dynamic, hopeless, and terminal disease forms. Palliative care must be recognized as an fundamental component of the armamentarium when managing with patients with surgical crises.

---

Improved information of this key component of care is crucial when caring for elderly patients.

## 1.2. Palliative Care

Palliative care has been characterized in different domains [1]. For occurrence, the World Health Organization characterizes as “the add up to dynamic care of patients whose infection is not responsive to corrective treatment. Control of torment, of other indications, and of mental, social, and otherworldly issues, is fundamental. The objective of palliative care is accomplishment of the best quality of life for patients and their families”. Of note, this definition emphasizes that palliative care is a “total” (i.e. biopsychosocialspiritual) and “active” (i.e. not latently holding up for death) care demonstrate that points to accomplish indication control when source control is not conceivable and care for quiet and family when remedy is not conceivable. Surgical palliative care has been sub-defined as beginning assessment and nearby control of infection, taken after by control of release, hemorrhage, and pain, and finishing with remaking and rehabilitation.

While diverse definitions may give one of a kind sees that encourage in general understanding, uncertain characterizations of palliative care have also conflated the part of surgery and its signs and results in patients, particularly with progressed cancer. Not at all like conventional corrective treatments, surgical palliative treatments cannot be characterized by the type of operation (e.g. the same surgery can be utilized to accomplish a particular reason in diverse patients) or the degree of disease (e.g. net edge examination may be unessential to the objective of the surgery)—the key distinction lies in the aim of care and individualized care. Hence, surgical palliation or palliative surgery is best characterized as an operation or method purposely performed with the essential deliberate of moving forward quality of life or calming side effects in a persistent with serious infection such as progressed threat. And making strides quality of life may cruel tending to the basic etiology such as obstacle, dying, or aperture and indications may incorporate torment, queasiness, heaving, anorexia, or jaundice. Essentially, the viability of palliative surgery is best characterized by the length of patient-acknowledged indication determination, or maybe than the length of by and large survival.

Surgical methods can be seen as either corrective or palliative [3]. Healing strategies are planned for patients with seizures convincingly localized to a particular cortical locale which is secure to expel. The objective in healing surgery in this manner is total resection of the influenced cortex, and methods incorporate front worldly lobectomy, particular amygdalohippocampectomy, neocortical resections, and hemispherectomy. Palliative surgery is utilized in circumstances where a seizure center is either not recognized or cannot be securely evacuated. For example, patients with intrinsic syndromic epilepsies such as Lennox–Gastaut disorder encounter life-threatening generalized seizures for which there is no identifiable center. The objective of palliative surgery hence is lessening in seizure recurrence and seriousness. Common palliative strategies incorporate situation of a vagus nerve stimulator and corpus callosotomy.

## Examples

When cure is no longer an choice, surgery can be viable for relieving indications that create in the progressed stages of cancer [4]. Cases of palliative surgery include:

- Treatment of a fungating breast wound
- Relief of bowel hindrance in ovarian cancer
- Debulking of a tumor to control release, the arrangement of fistulae, or hemorrhage
- Avoidance of hemorrhage when a tumor is squeezing on a imperative blood vessel
- Prophylactic or restorative sticking of metastases in long bones
- Spinal surgery to anticipate or stabilize spinal rope compression
- Debulking of tumor to control penetration of nerves which cause neuropathic pain
- Inclusion of intraepidural or intrathecal catheters for spinal opioids or neuro-lytic pieces for pain in such cancers as pancreas or gallbladder cancer

The point of palliative surgery is to diminish enduring and to minimize the side effects of the infection, so that if the quality of life will not be made strides, or if there is an pointless chance of horribleness or mortality, at that point palliative surgery ought to not be considered. People and their spouse/carers require to be educated of the points of palliative surgery, so that they can be reasonable in their desires of the surgery and make an educated choice. A few people may deny palliative surgery, favoring to spend what time they have cleared out without submitting themselves to a healing center remain or the dangers of surgery, while others may welcome the surgical mediation and see it as a treatment that may draw out their life. Once more, people ought to be upheld in their choice, as the signs and side effects that require palliative surgery are constantly those of dynamic disease, and as such practical desires of surgery ought to be emphasized, and the quiet upheld to get it these. Palliative surgery needs to be considered on an person premise and the choice based on each individual's side effects and their current quality of life against how surgery might move forward that quality.

## 1.3. Triangle

Functional Assessment of Cancer Therapy-G data—an objective measure that scores each patient’s physical, utilitarian, social, and passionate prosperity along with relationship with doctor—showed high scores in the preoperative period and kept up until death [1]. The Palliative Triangle was also assessed as a device for cautious understanding determination. Importantly, 1 or 2 meetings between patient, family, and specialist enduring 60–90 min took put some time recently making a choice for palliative surgery and the utilize of the Palliative Triangle was affirmed with each going to specialist. Generally, more than 90% of patients detailed side effect change or determination, and palliative operations were related with lower 30-day postoperative dreariness (20.1%) and mortality (3.9%) than already distributed comes about, appearing the promising esteem of the Palliative Triangle.

---

The key to the Palliative Triangle is an energetic relationship between the quiet, family, and specialist all through the palliative stage of the treatment—dynamic since the relationship must withstand the lability of patient's state and curveballs and all through since the relationship must be maintained in spite of troubles and detours. In the starting, there needs to be a clear definition of objectives of each patient's treatment, in agreement with the expectation and individualized nature of palliative surgery. At that point, through the elements of the triangle, each patient's values and meaning of life, accessible social and passionate bolster, and current restorative and surgical choices would be shared and weighed. In a perfect world, the reinforced triangle will also offer assistance intervene patient's and/or family's at times improbable desires. Each member of the Triangle is anticipated to fulfill his or her interesting commitment, as all three advance toward the collective picture of trust shaped by the energetic relationship.

#### 1.4. Palliation

Surgical palliation includes different components and bury: surgery with palliative aim and surgery with corrective aim [5]. Surgery with palliative expectation incorporates doing a surgical strategy with an result that trusts to move forward the patients' quality of life, whether that be side effect resolvment or simpler self-management. Surgery with corrective aim is done with the trusts of drawing out the life expectancy and quality of life. Choosing surgery as a treatment choice for palliative care is one that moreover comes with the troublesome assignment of forecast. Whether that be since of the vulnerability or the wide edges for blunder, weighing dangers and benefits is a huge portion when picking a surgical alternative. Whereas each surgery has a level of instability and has the potential for antagonistic results, the choice to seek after a surgical alternative for palliative care comes with a more prominent degree of hazard. Seeking after a surgical treatment course may not appear to be the best choice as it would not fundamentally draw out life span, but it may be viable at moving forward the patient's quality of life, indeed if for a brief time span. Palliative care choice making includes finding a adjust between doctor obligation of nonmaleficence and quiet independence. Nonmaleficence is regularly prioritized as doctors come with the deliberate of attempting to secure or maybe than make more awful if given a choice. This paternalistic trait is extreme to thrust aside indeed when the understanding is prepared to take hazard for indeed the littlest of benefits.

When surveying for a palliative counsel, there are numerous viewpoints of a patient's care that require to be considered. For case, the patient's determination and life hope play an critical part in deciding what treatment alternatives are indeed attainable given their soundness, but the patient's wanted treatment objectives may be fair as, if not more, vital, and may negate the recommended treatment proposals given their disease status. Palliative care may be advertised for patients that have a genuine or incessant sickness, a declining capacity to total exercises of every day life, individual or family trouble, the require for continuous long-term treatment, and numerous others. Calling for a palliative care counsel can cause trouble and uneasiness for the quiet, family, and care group

as it requires an existential discussion approximately objectives of care for the length of life cleared out accessible and in a few cases, whether experiencing a surgery is the best palliative option.

#### 1.5. Diabetes

Managing diabetes mellitus at the end-of-life rotates around the objective of de-prescribing with the aim of causing no hurt [6]. To date, there is small prove comparing diabetes treatment methodologies in patients with both diabetes and progressed infection, and a few unanswered questions stay as how to best approach glucose administration toward the end of life. Patients with steady and dynamic progressed illness with reasonable wholesome admissions (such as metastatic cancers, dementia, cardiomyopathy) may have the guess of a few months to a year, patients with approaching passing with organ or framework failure with anorexia or cachexia (as seen in cases of fulminant liver failure, bone marrow failure) may have the forecast of a few days or weeks, and patients effectively passing on may have the guess of a few hours to days. Diabetes administration ought to be custom-made to the quiet based on the seriousness of their illness state with the objective to de-escalate their diabetes care and unwind their blood glucose targets for palliative side effect control, whereas dodging the chance of DKA (diabetic ketoacidosis) and hyperosmolar hyperglycemic state (HHS). For patients with steady and dynamic progressed illness, diabetes administration may stay the same if the understanding is educating on how to screen and oversee extraordinary glycemc outings of hypoglycemia and hyperglycemia. In cases of renal failure, liver failure, anorexia, and dynamic weight misfortune, verbal diabetes specialists and/or affront may likely require measurements decreases. Patients with type 1 diabetes ought to proceed long-acting affront for as long as conceivable to maintain a strategic distance from DKA or with dosage diminishments to dodge any hypoglycemia. For patients with looming passing with organ or framework failure with anorexia or cachexia, the objective of administration is to dodge hypoglycemia due to expected parchedness, liver, and renal failure. Verbal anti-diabetes operators may require critical dosage diminishments or may require to be halted. Patients with T1DM would require critical diminishments in their long-acting affront dosage with disposal of pre-meal and possibly remedial affront. SMBG (self-monitoring of blood glucose) checks are by and large disposed of in patients with T2DM and utilized as it were where a choice needs to be made for administration in patients with T1DM. As in the past two stages, a agreement on administration is missing for patients effectively biting the dust. Most specialists in this case would basically pull back all verbal hypoglycemics and halt affront in most cases of diabetes mellitus. At this point, care is centered on patient's consolation and preliminary loss counseling for caretakers and patients, where appropriate.

#### 1.6. Patient

The overriding fight in palliative care is the choice between the physician's obligation to maintain a adjust between nonmaleficence and advantage versus regarding quiet independence [5]. Whereas conducting a palliative surgical strategy seem offer assistance make strides the patient's quality of life, it may also demonstrate

---

hindering for the patient's continuous treatment or length of survival. Subsequently, the struggle of deciding whether to seek after nonmaleficence or quiet independence arises. Patient independence and doctor obligation to dodge hurt are two imperative standards that direct choice making. Understanding independence alludes to the right of a persistent to make their possess choices approximately their healthcare, based on their values, convictions, and inclinations. Doctor obligation, on the other hand, alludes to the obligation of a doctor to act in the best intrigued of their patients and give care that meets the most noteworthy measures of restorative hone. Doctor obligations advancing nonmaleficence and advantage have been the center of medication for decades; be that as it may, there has been a move in the final few years from the conventional "paternalistic" practice of medicine to an emphasis on "patient-centered care". This system shifts the center from telling the persistent what to do to encouraging shared choice making with the quiet based on their objectives and values. It implies the doctor is no longer the despot of persistent healthcare but or maybe an master that is included in the discourse and show for exhortation, direction, and most critically, data. The Institute of Medicine identifies this as being "responsive to person understanding inclinations, needs, and values". In palliative care, doctors frequently discover themselves choosing whether to back the patients' inclinations or doing what is considered to be the most elevated standard of restorative hone. One way to discover a adjust between the two is by prioritizing the guideline of educated consent.

Informed assent includes giving patients with all the data they require to make an educated choice approximately their care. This incorporates data almost all the potential dangers, benefits, and the capacity of each treatment alternative that is not skewed by doctor conclusion. The objective with educated assent is that the understanding may at that point make an taught choice with respect to their restorative care that best serves their objectives and values in life, which are profoundly particular to and shift for each person. If the understanding chooses the dangers and results of a surgical intervention are worth the potential benefits, at that point it is their choice to take those dangers. Whereas a specialist may not concur with the treatment choice, if they are willing and able to total the surgery, it is the best treatment course to be taken. By giving patients with data almost their alternatives, suppliers can regard independence whereas moreover satisfying their obligation to act in the patient's best interest.

### 1.7. Emergency

Palliative care begins well some time recently confirmation to intensive care units (ICU), in the sense that it can be characterized as treatment that does not specifically change the illness process(es), driving to the clinical introduction: for illustration, this may be basically giving compelling absense of pain [7]. Multidisciplinary palliative care that is more formal will gotten to be the fundamental need when treatment alternatives are distinguished as pointless. Tolerating that assist clinical care is pointless is exceptionally troublesome for numerous specialists, and frequently for the near family of the quiet as well. Convincing the current elderly cohort

of patients that medication does not give immortality is as a rule direct, but the over- hopeful depiction of recuperation from close death on TV programs is confounding for their relatives and more youthful family individuals. Making sufficient time to teach them into understanding reality may be difficult. On event, this makes it troublesome to move from treatment to palliation. The timing of execution of these choices will depend on the acknowledgment by the understanding and their family that there is no assist advantage to be picked up and that advance frameworks back will as it were drag out the prepare of biting the dust. This may take no time at all or a few days. During the last mentioned prepare, the care group will moreover require passionate bolster to proceed what they may accept is pointless interventional management. The palliative care group, seriously care specialists, and the understanding, with their carers, will investigate alternatives to guarantee that all human rights of the persistent are in put. This will incorporate absense of pain for symptomatic control of pain, religious support, financial and social advice, and every day reviews to monitor progress. Arranging for their inescapable death and the future care for their family are zones that require to be talked through and any back that may offer assistance ought to be identified. One of the most extraordinary and squeezing circumstances for the appraisal of worthlessness is when a cardiac capture happens in the clinic. This habitually includes elderly patients who have critical pre- existing therapeutic issues. The victory of revival driving to release from the clinic is exceptionally low. Around 5% of all patients will survive, but this falls with expanding age. No survivors are likely after the age of 70, in spite of the fact that they may be revived sufficient to be conceded to the ICU. Of those who are conceded, 50% have critical neurological harm, and the others create misery or other push- related disorders. In the confront of these shocking results, numerous elderly patients select to order DNAR (Do Not Attempt to Resuscitate) in the occasion of a cardiac arrest. These wishes may be formally communicated by the quiet or more casually through their relatives. If the ward staff are not mindful of these DNAR records, resuscitation will be attempted.

## 2. Conclusion

The positive affect of science and innovation has empowered the application of advanced strategies, moved forward treatment, amplified life hope and diminished mortality. Be that as it may, the number of individuals enduring from hopeless illnesses is expanding. The increment in dementia, dangerous, unremitting and other infections adversely influences the quality of life. Palliative surgical methods require complex adjustment to the patient's recently created health condition and specific care. Depending on the patient's needs and complaints, strategies are most regularly performed to meet the patient's needs and guarantee their quality of life. Methods must not decline the patient's condition, so the choice of method must be such that it carries as few complications as possible.

## References

1. Shin, J., & Miner, T. J. (2024). Palliative Surgery: When To and When Not To?. *In Recent Strategies in High Risk Surgery* (pp. 673-692). Cham: Springer Nature Switzerland.

- 
2. Britt, L. D., & Martyak, M. (2023). Acute care surgery in the geriatric patient population: general principles. In *Acute Care Surgery in Geriatric Patients* (pp. 1-5). Cham: Springer International Publishing.
  3. Doherty, G. M. (2024). *CURRENT diagnosis and treatment surgery*. McGraw Hill Professional.
  4. Poston, G. J., Beauchamp, R. D., & Ruers, T. J. M. (Eds.). (2007). *Textbook of surgical oncology* (pp. 1-4). Informa Healthcare.
  5. Mistry, K., Stover, A., & Ganai, S. (2024). How Should Surgical Palliative Success Be Defined?. In *Recent Strategies in High Risk Surgery* (pp. 693-701). Cham: Springer Nature Switzerland.
  6. Presswala, L. S., Farooki, A., & Flory, J. (2024). Diabetes Care in the Oncologic Population. In *Diabetes Management in Hospitalized Patients: A Comprehensive Clinical Guide* (pp. 355-370). Cham: Springer International Publishing.
  7. Dodds, C., Kumar, C. M., & Servin, F. (2016). *Anaesthesia for the elderly patient*. Oxford University Press.

*Copyright:* ©2026 Siniša Franjić. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.