

## Prevalence of Vitamin A Deficiency And Associated Factors Among Children Aged 6-59 Months In Dera District, Northwest Ethiopia: A Cross-Sectional Study

Yidsrail Hune<sup>1\*</sup>, Hirut Asaye<sup>2</sup>, Ayenew Negesse<sup>1</sup>, Habtamu Temesgen<sup>1</sup>, Tadesse Yirga<sup>3</sup>, Biachew Asmare Fenta<sup>1</sup>, Amare Bekalu<sup>1</sup> and Habtamu Mekonnen<sup>1</sup>

<sup>1</sup>Department of Human Nutrition and Food Science, College of Health Sciences, Debre Markos University, Debre Markos, Ethiopia

<sup>2</sup>Department of Applied Human Nutrition, Faculty of Chemical and Food Engineering, Bahir Dar University, Bahir Dar, Ethiopia

<sup>3</sup>Department of Pediatrics and Child Health Nursing, College of Health Sciences, Debre Markos University, Debre Markos, Ethiopia

### \*Corresponding author

Yidsrail Hune, Department of Human Nutrition and Food Science, College of Health Sciences, Debre Markos University, Debre Markos, Ethiopia

Submitted: 20 Oct 2021; Accepted: 26 Oct 2021; Published: 30 Oct 2021

**Citation :** Yidsrail Hune, Hirut Asaye, Ayenew Negesse, Habtamu Temesgen, Tadesse Yirga, Biachew Asmare Fenta, Amare Bekalu, Habtamu Mekonnen (2021) Prevalence of Vitamin A Deficiency And Associated Factors Among Children Aged 6-59 Months In Dera District, Northwest Ethiopia: A Cross-Sectional Study. *J Pediatr Neonatal Biol* 6(2): 35-43.

### Abstract

**Background:** Vitamin A is a fat-soluble vitamin. It comes retinol from animal sources or beta-carotene from plant source. Vitamin A contains breast milk after the post-weaning period and their increased nutrient demand of children from 6-59 months, they are considered highly affected segments of the community. Hence, study assesses vitamin A deficiency and associated factors among children aged 6-59 months in Dera district, Northwest Ethiopia.

**Methods:** Cross-sectional study design was conducted via multistage sampling techniques. Data were coded and entered into Epi-data version 3.1 and exported to Statistical Package for the Social Sciences (SPSS) version 20 for analysis. The bivariate and multiple variable logistic regression analyses were fitted.

**Results:** Total response rate 94.7%. Prevalence of vitamin A deficiency among children age from 6-59 months was 7.8% (95% CI: 4.7, 11.3). Anti natal care (ANC) follow-up (AOR = 0.446; 95 % CI; 0.155, 0.980), and birth interval (AOR= 0.392: 95 % CI; 0.107, 0.839), reduces the odds of developing vitamin A deficiency; whereas age group of 36-47 months (AOR= 1.911: 95 % CI; 1.305, 11.969) increases the odds of its deficiency.

**Conclusion:** Age of children, birth interval, ANC follow-up, Post natal care (PNC) follow-up, and place of residence were associated factors for vitamin A deficiency. Therefore more efforts should be encouraged to produce and purchase a variety of foods rich in vitamin A.

**Keywords:** Vitamin A, Dera District, 6-59 Months Old Children.

### Background

Vitamin A Deficiency (VAD) can be defined clinically or subclinically. Xerophthalmia is the clinical spectrum of ocular manifestations of vitamin A deficiency; these range from the milder stages of night blindness and Bitot spots to the potentially blinding stages of corneal xerosis, ulceration and necrosis (keratomalacia). The various stages of xerophthalmia are regarded both as disorders and clinical indicators of vitamin A deficiency. VAD is among the top 10 risk factors contributing to the global burden of disease among

preschool-age children in resource-limited countries [1, 2]. (Globally, around 33.3 % of preschool-age children (One third (190 million) of the world's preschool children) were Vitamin A deficient, and 15.3% of pregnant women in populations at risk of VAD. An estimated 44.4% of preschool children in Africa (56.4 million children) are vitamin A deficient and are supposed to contribute to over 1 million childhood deaths a year [3, 4]. Recent reviews showed that the sub-Saharan Africa region has the highest rate of VAD (48%) prevalence in children <5 years [3].

Vitamin A deficiency alone is responsible for almost 6% of child deaths under the age of 5 years in Africa and 8% in South-East Asia [3]. Vitamin A supplementation in children 6–59 months of age living in developing countries is associated with a reduced risk of all-cause mortality and a reduced incidence of diarrhea [5].

The consequences of VAD are magnified by poverty and the higher prevalence of the infectious disease and it is an underlying cause for nearly one-fourth of global child mortality from measles, diarrhea, and malaria [6, 7]. This mortality risk worsens among children born in sub-Saharan African countries, which face 16.5 and 1.8 times higher probability of dying before the age of 5 years compared to children born in developed regions and Southern Asia respectively [8].

In Ethiopia national prevalence of vitamin A deficiency estimated based on retinol adjusted for inflammation among preschool children was found 13.9%. And the lowest prevalence was observed in Addis Ababa; almost all preschool children in this city administration were not at risk of vitamin A deficiency. Among the region's the prevalence of vitamin A deficiency of preschool children who live in Harari was the highest as compared to other regions at a prevalence of 21.0% and moderate prevalence of vitamin A deficiency was observed among preschool children who live in the Amhara region (10.3%) [9]. Hence based on WHO classification, this prevalence can be categorized as a moderate public health problem in Ethiopia. Likewise, the prevalence of Vitamin A deficiency can be considered as a moderate public health problem in all regions, except Harari and Addis Ababa. Vitamin A deficiency is a severe public health problem among the Harar preschool children at a prevalence of 21%. On the other hand vitamin A deficiency is not a public health problem among the Addis Ababa preschool children [9]. In Ethiopia Mortality in children who are blind from Keratomalacia or who have the corneal disease is reported to be from 50% to 90% and measles mortality associated with VAD is increased up to 50% [10, 11].

The problem of vitamin A deficiency in developing countries is inextricably linked to poverty. Low incomes constrain families' abilities to purchase the diverse foods they need for a nutritious and balanced diet with adequate amounts of vitamin A [12]. Maternal vitamin A deficiency results in low concentrations of vitamin A in breast milk, predisposing infants to deficiency. This is exacerbated by an inadequate dietary intake of vitamin A during the complementary feeding period. Finally, illness can worsen vitamin A status because of a reduced food intake from anorexia, malabsorption, and increased excretion in urine [13].

To redress vitamin A deficiency, megadose vitamin A capsules are administered twice yearly to preschool children in many developing countries, often in conjunction with the Expanded Program of Immunization (EPI). Children can receive micronutrients from foods, fortified food, and direct supplementation. But inadequate intake of this micronutrient and the presence of infections such as measles and diarrheal diseases in children are the cause of Vitamin A deficiency (VAD) [14].

VAD can occur in individuals of any age. However, vitamin A deficiency is a disabling and potentially fatal public health problem

for children under 5 years of age. VAD-related blindness is most prevalent in children under 3 years of age [10]. This period is very sensitive and critical characterized by high requirements for vitamin A to support early rapid growth, the transition from breastfeeding to dependence on other dietary sources of the vitamin, and increased frequency of respiratory and gastrointestinal infections. The increased mortality risk from concurrent infections extends at least to 6 years of age and is associated with both clinical and subclinical VAD [15].

Children have protein-energy malnutrition and anemia affected by VAD, VAD causes night blindness and Bitot's spot during childhood period this indicator of increased morbidity and mortality in preschool children. So children are more vulnerable to illness, both infectious and non-infectious [16].

Seven national regional states of Ethiopia showed that the prevalence rate of Bitot's spots was highest in Amhara (3.2%), followed by Afar (2.1%), Oromiya (1.5%), Addis Ababa (1.4%), Harari (1.2%), and Dire Dawa (1.1%) regions and it was significantly associated with male sex, rural residence and greater age [17]. In Ethiopia and other African countries, poverty, sub-optimal nutrition, unsanitary living conditions, and poor health care access exacerbate the risk of developing multiple micronutrient deficiencies [18]. 26% of the young children consume vitamin A rich foods, while only 4% receive food from at least four food groups [19]. Therefore, the purpose of this study is to provide information regarding the prevalence and factors associated with Vitamin A Deficiency among 6-59 months children.

### **Rationale of the Study**

Ethiopian national nutrition program targeted to improve the nutritional status of children from birth up to ten years. Thus prevention and control of micronutrient deficiencies is one of the initiatives. It has been confirmed that improving the vitamin A status of deficient children significantly reduces the risk of mortality from measles by 50%, from diarrhoea by 40%, and overall mortality by 25-30%. Therefore the elimination of VAD is considered a key element for improving the survival, wellbeing, growth and development of children. Thus, research showing the burden and determinants of VAD are of paramount importance. However, research study based on information regarding night blindness, Bitot's spot and vitamin A intake among pre-school children is limited, particularly to this study area. In general this study will give theoretical as well as practical significance for health care provider, program planner, researchers, policy makers to use as baseline data to conduct large scale on the level and factors to develop the intervention plan that makes preschool children vitamin A deficient.

### **Objectives of the Study**

#### **General objective**

- To assess the prevalence of VAD and associated factors among children aged 6-59 months in Dera wereda, South Gonder, Ethiopia.

#### **Specific objectives**

- To determine the prevalence of VAD among children aged 6-59 months in Dera wereda.
- To identify factors associated with VAD among children aged 6-59 months in Dera wereda.

## Methods

### Study Design and Setting

A community-based cross-sectional study was conducted in Dera Woreda, South Gonder Zone from December 1/2018 to January 1/2019. Dera Woreda has located 600 km from Addis Ababa in the Northwest and 40 km from Bahir Dar in the Northeast. The estimated population of the Woreda in 2017/18 is about 154,886. The estimated number of the under-five year children is 22,567. The total number of estimated children aged 6-59 months were 19,282. Quantitative data were collected in the Community mothers who have a child age 6-59 months.

All participants were allowed to ask questions throughout data collection and could refuse to answer questions or stop the interview at any moment.

### Sample Size Determination and Sampling Procedure

The sample size was determined using a single population proportion formula

$$N = \frac{\left(\frac{z_{\alpha}}{2}\right)^2 p(1-p)}{d^2} [28],$$

where, p represents the proportion of children who took the Vitamin A Deficiency, which was 8.6% taken from the study done in Dembia [20]. To get the optimum sample size, 5% margin of error (d) was considered with 95% confidence interval.

$$N = \frac{(1.96)^2 0.086(1 - 0.086)}{0.05^2} = 120.7 = 121$$

We added 10% for the non-response rate and multiplied by 2 because of the design effect. Then the final sample size was 267.

Multistage sampling technique was implemented. of the 39 administrative Kebeles in Dera Woreda, nine were selected randomly by lottery method. Among those kebele 20-30 % of the population proportional allocation was done to select the desired samples from each selected Kebele. The sampling frame, based on a community-based health information system of the family folder in health posts, was constructed. Lists of all mothers having children age 6-59 months with Community Health Information System (CHIS) number in selected kebeles were used to select the respondents. Finally, we used computer-generated random sampling to select the desired sample from the sampled frame of each selected kebele. other-child pair having a child aged 6 to 59 months who has a mental problem, critically ill were excluded.

### Data Collection Procedure and Measurements

Questionnaires were first prepared in English and translated into Amharic Version, which later back-translated into English. Amharic version questionnaires were used to collect data at the community level. A structured questionnaire was used to collect data on socio-demographic characteristics, maternal and child characteristics, and a Pre-test was done on 5% of the samples, two weeks before the actual data collection at Fegera Woreda. Data on Vitamin A deficiency were collected through face-to-face interviewing and with direct observation of children's eyes of children age 6-59 months.

### Assessment of Dietary Diversity and Quasi-Food Frequency

Food consumption studies can provide important information at an earlier stage before the clinical manifestation has appeared. To determine the food consumption of the children twenty-four-hour recall method was used. One single twenty-hour recall was collected for every participant and dietary diversity was calculated. Determination of dietary diversity score (DDS) was done by asking the mother/caregiver to list all foods consumed by the child in the previous 24 hours preceding the survey. In the case of a mixed dish, mothers were asked to list the ingredients of the food items. Then reported food items were classified into seven food groups like grains, legumes, vitamin A-rich fruits and vegetables, other fruits and vegetables, egg, dairy products (milk, yogurt, and cheese), and meat. Considering four food groups as the minimum acceptable dietary diversity, a child with a dietary diversity score (DDS) of less than four was classified as having poor dietary diversity; otherwise, they were considered to have good dietary diversity [21].

### Assessment of Vitamin A Deficiency

A detailed ophthalmic examination was carried out by clinical optometrists was used to assess the clinical signs of vitamin-A deficiency and by using all data collectors in hand pictures of those VAD, such as Bitot's spot, Conjunctival xerosis, corneal xerosis, corneal ulceration, and corneal scar in order to identify each VAD. However, during data collection time the history of night blindness was confirmed by asking mothers about their children using the local word for night blindness "dafint" or "chicken eye". Information on whether a child faced any difficulty while playing or in identifying objects in dim light, especially at sunset, was gathered. In addition to this Six data collectors (2 clinical optometrists & 4 clinical nurses) and two supervisors (public health officers) were participating in the data collection process. One day of training was provided to data collectors and supervisors about how to collect the data and how to ask the investigators.

The Dependent variable of this study is Vitamin A deficiency and Independent variables are Socio-demographic and economic characteristics, Health utilization, and Child factors.

### Operational Definition

#### Optimal Dietary Diversity

Dietary diversity was defined as optimal if children (aged 6-59 months) received foods from at least four of seven food groups (1) Grains, roots, and tubers, (2) Legumes and nuts, (3) Dairy products, (4) Flesh foods, (5) Eggs, (6) Vitamin-A rich fruits and vegetables, (7) Other fruits and vegetables, within the preceding 24 hours of the interview.

#### Sub-Optimal Dietary Diversity

Suboptimal dietary diversity:-If a child aged 6-59 months receive foods from three or below three food groups from seven food groups [22].

VAD- Vitamin A deficiency can be defined clinically or subclinically in our work to give working definition for VAD if a child has a history of night blindness and during the physical examination if the child has one signs of vitamin A deficiency, such as Bitot's spot, Conjunctival xerosis, corneal xerosis, corneal ulceration, and

corneal scar.

The terms ‘Women and mothers’ are used interchangeably in this study.

### Statistical Analysis

Data were entered into EpiData version 3.1 with double entry for verification. The analysis was performed using SPSS version 20.0. Descriptive data presented by using Frequency and cross-tabulation. Both the bivariate and multivariable logistic regression analyses were used to assess the association between dependent and independent variables. Independent variables that showed  $P < 0.2$  at 95% CI in the bivariate logistic regression analysis were included in the multivariable logistic regression model.  $P < 0.05$ , with 95% CI, was considered to declare the variables significantly associated with the dependent variable.

**Table 1: Sociodemographic and economic characteristics of mothers having a child aged 6-59 months in Dera Woreda, North-west Ethiopia, 2019/20(N=267)**

Variables		Frequency	
		N	%
Maternal age	15-35 year	213	84.2
	36-49 year	40	15.8
Maternal religion	Orthodox	237	97.3
	Muslim	16	6.3
	Others	-	-
Maternal ethnicity	Amhara	253	100
	Others	-	-
Maternal marital status	Single	11	4.3
	Married	215	85
	Divorced	25	9.9
	Windowed	2	0.8
Place of residence	Urban	165	65.2
	Rural	88	34.8
Maternal education level	Unable to read and write	186	73.5
	Read and write only	19	7.5
	Primary school (1-8 grade)	27	10.7
	Secondary school (9-12 grade)	15	5.9
	Higher education	6	2.4
Father’s education	Unable to read and write	132	52.2
	Read and write only	73	28.9
	Primary school (1-8 grade)	23	9.1
	Secondary school(9-12grade)	20	7.9
	Higher education	4	1.6
Maternal Occupation	Housewife	209	82.6
	Government employed	9	3.6
	Merchant	23	9.6
	Farmer	9	3.6

### Results

Of the total 267 sampled mother-children Pairs takers who had children 6-59 months, 253 of them participated in the study with a response rate of 94.7 %.

### Socio-Demographic Characteristics of Mothers

More than half of the participants 165 (65.2%) were living in the urban kebele of Dera Woreda. Of the respondents of the study, 237 (93.7%) orthodox religious followers, and 100% were Amhara in the ethnic group. Almost 105 (41.5%) of the study participants live in a household that has seven and above household members. The majority of 209 (82.6%) of the respondents were housewives; married 215 (85%) and illiterate 186 (73.5%). About half 173 (52.2%) of the fathers were unable to read and write and the majority 211 (83.7%) of them were farmers in their occupation (Table1).

	Student	3	1.2
Fathers' occupation	Farmer	211	83.7
	Government employed	15	6
	Private employed	1	0.4
	Merchant	23	9.1
	Student	1	0.4
Household family size	<=4	66	26.1
	5-6	82	32.4
	>=7	105	41.5
Availability of Media sources* at the households	Yes	62	24.5
Exposure to available media sources	Yes	44	17.4
Available farmland	Yes	211	83.3
Cultivating vegetables	Yes	176	69.5
Mother involved in the decision-making	Yes	110	43.4

\*Includes Radio and Television (TV)

#### Characteristics of the Children Aged 6-59 Months

About more than half 160 (63.2%) of children were male and 68 (26.9%) were found in the age group of 36-47 months. The majority of 189 (74.7%) of them was a second and above child for the

interviewed mothers. About 108 (42.6%) of them had received a growth-monitoring service and 208 (82.2 %) of them received a measles vaccine (Table2).

**Table 2: Selected health-related characteristics of children aged 6-59 months in Dera Woreda, Northwest Ethiopia, 2019/20(N=267)**

Variables		Frequency	
		N	%
Child sex	Male	160	63.2
	Female	93	36.8
Child age in month	6-23	58	22.9
	24-35	60	23.7
	36-47	68	26.9
	48-59	67	26.5
Child live at home	<2	64	25.3
	>=2	189	74.7
Growth monitoring service	Yes	108	42.6
Birth interval	< 2years	24	9.5
	>=2 years	229	90.5
The morbidity for the last two weeks	Yes	15	5.9
Type of morbidity	Diarrhea	24	37.5
	Fever	15	5.9
	Cough	24	37.5
	Others	9	14
BCG Vaccinated	Yes	232	91.6
Pentavalent1 vaccinated	Yes	232	91.6

<b>Pentavalent3 vaccinated</b>	Yes	215	84.9
<b>Measles vaccinated</b>	Yes	208	82
<b>Growth monitoring practice follow up</b>	Yes	108	42.6

BCG: Bacillus Calmette-Guérin, Pentavalent: A vaccine against Diphtheria, Pertussis, Tetanus, Hepatitis B, and Haemophilus influenza type B

### Health Utilization and Dietary Related Characteristics of Mothers


The majority of 143 (56.52%) of mothers had no antenatal care (ANC) visits for the current children. The majority of 225 (88.9%) of children took vitamin A supplementation in the last 6 months. The majority of 225 (88.9%) of children's had dietary diversity scores below and equal to 4 and also 176 (69.5%) three fourth of mothers delivered at the clinic (Table3).

**Table 3: Health service utilization and dietary practice of mothers in Dera-Woreda, Northwest Ethiopia, 2019/20(n=267)**

Variables		Frequency	
		N	%
ANC follow up for the index child	Yes	110	43.8
Place of delivery	Home	70	27.2
	Clinic	176	69.5
	Hospital	5	1.9
Mode of delivery	Spontaneous vaginal delivery	228	90
	Assisted deliveries	25	9.8
PNC follow up for the index child	Yes	108	42.7
Vitamin A supplementation	Yes	225	88.9
Dietary diversity score	<=4	225	88.9
	>4	28	11.1

### Prevalence of Vitamin A Deficiency

The prevalence of Vitamin A deficiency among children age from 6-59 months was 7.8% with 95% CI (4.7, 11.3), (figure 1)



**Figure 1: The prevalence of vitamin A deficiency**

### Factors Associated With Vitamin A Deficiency Among Children Aged 6-59 Months

In the bivariable analyses age of children, Sex, place of residence, ANC visit, PNC follow-up, Cultivate dummy vegetable and the birth interval were significantly associated with Vitamin A deficiency. Those variables were significant in bivariable analysis, including multivariable analysis. In multivariable analysis age of

children, sex of children, birth interval, ANC, and PNC follow-up were significantly associated with Vitamin A deficiency. The birth interval of greater than two years was 60.8% times less likely to suffer from VAD compared to counterparts (AOR=0.392,95% CI:0.107,0.839). On the other hand, age groups 36-47 months were 1.91 times more like to have VAD compared to the age group 24-35 months (AOR 1.911,95%CI: 1.305,11.969). Likewise, moth-

ers who had ANC visits had 55.4 % times less likely to develop VAD compared to mothers who did have ANC visits (AOR=0.446, 95%CI: 0.155,0.980). Similarly, children from mothers who had

PNC visits had 63.3 % times less likely to develop VAD compared to children from mothers who did not attend PNC (AOR=0.337, 95%CI: 0.111, 0.924)(Table 4).

**Table 4: The bivariate and multivariable logistic regression analysis showing factors associated with VAD among children aged 6-59 months in Dera Woreda, Northwest Ethiopia**

Variables		VAD		Crude Odds Ratio (COR)(95% CI)	AOR (95% CI)
		No	Yes		
Sex	Male	143	17	0.280(0.080,0.984) ***	0.296(0.077,1.141)
	Female	90	3	1	1
Age of children in a month	6-23	47	11	0.271(0.081,0.905) ***	0.381(0.099,0.774) *
	24-35	57	3	1.206(0.259,5.623)**	1.206(1.037,6.129)**
	36-47	66	2	2.095(0.371,11.844)	1.911(1.305,11.969)
	48-59	63	4	1	1
Place of residence	Urban	146	19	0.088(0.012,0.671)*	
	Rural	87	1	1	
ANC	Yes	97	13	0.384(0.148,0.998)*	0.446(0.155,0.980)*
	No	136	7	1	1
PNC follow up	Yes	94	14	0.290(0.108,0.781)**	0.337(0.111,0.924)**
	No	139	6	1	1
Cultivate dummy vegetable	Yes	157	19	0.109(0.014,0.827)*	
	No	76	1	1	
Birth interval	Within >2 years	19	5	0.266(0.087,0.813)*	0.392(0.107,0.839)*
	Within 2 years	214	15	1	1

Significant at \**p*-value < 0.05, \*\**p*-value < 0.01, and \*\*\**p*-value < 0.001

## Discussion

This study tried to assess the prevalence of vitamin A deficiency and associated factors among children age 6-59 months. The overall prevalence of VAD was 7.8%. The finding of this study in line with the study done in Demba, 8.6% [20]. But the current study finding was higher than the study done in Mali 5.4% and the reports of WHO cut off point, 1.56 % [23, 24]. The discrepancy might be due to the existence of differences in the socio-economic status of the study area, living condition, lack of adequate dietary diversity (11.1%) and lack of maternal literacy in the study area (18.6%) [25]. Adequate diets and maternal literacy are of greater value for children to resist vitamin A deficiency [25].

In this study, Conjunctival xerosis (1.6%), Bitot's spots (2.4%), Corneal Ketosis (0.4%), corneal ulceration (1.6%), and Corneal scar (1.6%) were the sign of VAD reported.

The prevalence of Bitot's spots was higher than the study done in different parts of India and Mali (2%), but lower than the study done in Amhara, Ethiopia (3.2%) and higher than the study done in different parts of Ethiopia (2.1%, 1.5%, 1.2%, 1.4%, and 1.1%) [17, 23, 26]. This difference might be due to socio-demographic

factors, place of residence, maternal health care, the purchasing power of the community, food price, child health care, low intake of beta carotene-rich and vitamin A containing foods [27].

This study also identified that the age of children, birth interval, antenatal care, and postnatal care were factors significantly associated with vitamin A deficiency.

In this study age of the child was a statistically significant factor for vitamin A deficiency. Those children whose age group 24-35 months was 1.21 times more likely affected by vitamin A deficiency than 48-59 months age in Dera District. This finding in line with a study done in Indian, different parts of Ethiopia, and North West Ethiopia [17, 20, 26]. This occurs might be in Ethiopia most children stop breastfeeding and continue family food after the age of 24 months. This leads to an inadequate intake of vitamin A reach foods from their diet that leads to a deficiency of vitamin A. Additionally, undernutrition is common during the periods of complementary feeding practices in Ethiopia. On the other hand, it might be due to physiological changes to support growth and development during this period needs additional micronutrients.

The presence of antenatal care follow up was significantly associated with vitamin A deficiency in the Dera district, North West Ethiopia. Those mothers who had ANC follow up care during pregnant of the index child were 55% decreases in the presence of vitamin A deficiency in their children. This finding is supported by the study done in North West Ethiopia [20]. This occurs due to during ANC follow-up each pregnant woman is consoled about diversified feeding practices for both them and their children. Therefore pregnant women who had ANC follow up have a better understanding and knowledge about child feeding practices and also gain adequate knowledge about vitamin A rich foods during ANC follow-up periods [28-30].

Additionally, Vitamin A deficiency was associated with the presence of postnatal care during postnatal periods. Mothers who had PNC visits during postnatal periods were 64% times less likely to develop vitamin A deficiency among children age 6-59 months. This result was supported by the study done in Indian and North West Ethiopia [20, 26]. The reason might be due to during the postnatal periods there are counseling sessions about infant and child feeding practice that results in the increased awareness about adequate dietary diversity feeding practices and increase knowledge towards vitamin A rich foods.

The birth interval was also factored in significantly associated with vitamin A deficiency. The children delivered within equal and greater than two years of the birth interval were 60% times less likely to develop vitamin A deficiency than delivered within two years of birth interval. This in line with the study findings done in Bahir Dar [28, 29]. The reason might be due to this interval the children's had freely breastfeeding in addition to family food and a good mother to child interaction.

## Conclusion

The overall prevalence of vitamin A deficiency in this study was lower than the national and regional prevalence. According to this study, vitamin A deficiency is a mild public health problem in the study area. Age of child, birth interval, ANC follow up and PNC follow-up were factors associated with Vitamin A deficiency in Dera District, North West Ethiopia. Therefore, special focus on children ages greater than 24 months. Additionally, maternal health services especially ANC, PNC, and family planning services to decrease the prevalence of vitamin A deficiency in children age 6-59 months in Dera District, North West Ethiopia.

## Abbreviations

ANC Ante-Natal Care, ANRS Amhara National Regional State, CHIS Community Health Information System, DDS Dietary Diversity Score, EDHS Ethiopia Demographic Health Surveys, FAO Food and Agriculture Organizations, MUAC Mid Upper Arm Circumference, PNC Post Natal Care, SPSS Statistical Package for Social Science, VAD Vitamin A Deficiency, VAS Vitamin A Supplementation, WHO World Health Organization.

## Declarations

### Ethics approval and consent to participate

We obtained ethical clearance from the Ethical review committee of the Faculty of chemical and food engineering of Bahir Dar University. Verbal informed consent, which was prepared in written

form and dictated to the respondents during data collection, was obtained from the study participants after explaining the purpose of the study and the benefits. In the case of minors (participants below 16 years old), verbal assent was obtained from their parents or guardians. Participation was voluntarily and the data were kept anonymous. We used verbal consent as the majority of the respondents were unable to read and write. The ethical review committee has approved this consent. Respondents were interviewed voluntarily, anonymously and confidentiality also was assured.

## Acknowledgments

We thank Bahir Dar University Faculty of chemical and food engineering for providing ethical clearance, data collectors, supervisor, and study participants for their willingness to participate in the study.

## References

1. Gebremedhin S, Loha E, Abebe Y, Dese G (2009) Assessment of vitamin A supplementation coverage and its association with childhood illness in Boloso Sore Woreda, Wolaita Zone, SNNP Region, Ethiopia. *Ethiop J Health Dev* 23: 223-228.
2. Arimond M, Ruel MT (2002) Progress in developing an infant and child feeding index an example using the Ethiopia Demographic and Health Survey 2000.
3. WHO (2009) Global Prevalence of Vitamin A Deficiency in Populations at Risk 1995–2005: WHO Global Database on Vitamin A Deficiency. World Health Organization 2009: 1-68
4. Y T Asrat, A M Omwega, J W G Muita (2002) Prevalence of vitamin A deficiency among pre-school and school-aged children in Arssi Zone, Ethiopia. *East Afr Med J* 79: 355-359.
5. Oumer MM, Mengestu Z, Yitayih S, Mequantant M, Molla A (2020) The determinant of National Vitamin A supplementation in Ethiopia: A Multilevel Logistic Regression Analysis.
6. Keith P West Jr (2002) Extent of vitamin A deficiency among preschool children and women of reproductive age. *J Nutr* 132: 2857-2866.
7. Ezzati M, Lopez AD, Rodgers AA, Murray CJ (2004) Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors. World Health Organization 2004: 1-2258.
8. You D, Jin NR, Wardlaw T (2012) Levels & trends in child mortality.
9. EPHI (2016) Ethiopian national micronutrient survey report. 2016: 1-126.
10. Sommer A (1994) Vitamin A deficiency and its consequences: A field guide to their detection and control. World Health Organization 1994: 1-88.
11. G D Hussey, M Klein (1990) A randomized, controlled trial of vitamin A in children with severe measles. *N Engl J Med* 323: 160-164.
12. Haidar J (2011) Common micronutrient deficiencies among food aid beneficiaries: Evidence from refugees in Ethiopia. *Ethiop J Health Dev*. 25: 222-229.
13. Miller M, Jean Humphrey, Johnson E, Marinda E, Brookmeyer R, et al. (2002) Why do children become vitamin A deficient? *J Nutr* 132: 2867-2880.
14. Mofokeng MJ (2013) Nutritional status and dietary intake patterns of children aged 7-13 years in Qwa-Qwa. 2013: 1-199.
15. Beaton GH (1993) Effectiveness of Vitamin A Supplementation



- tion in the Control of Young Child Morbidity and Mortality in Developing Countries – Nutrition policy discussion paper No. 13. 1993: 1-166.
16. Avula L, Madhavan K Nair, Nimmathota Arlappa, Pullakhandam R, Nagalla B, et al. (2011) Prevalence of ocular signs and subclinical vitamin A Deficiency and its determinants among rural pre-school children in India. *Public Health Nutr* 15: 568–577.
  17. Demissie T, Ahmed Ali, Mekonen Y, Haider J, Umata M (2010) Magnitude and distribution of vitamin A deficiency in Ethiopia. *Food Nutr Bull* 31: 234-241.
  18. Grantham McGregor S, Yin Bun Cheung, Santiago Cueto, Glewwe P, Richter L, et al. (2007) Developmental potential in the first 5 years for children in developing Countries. *Lancet* 369: 60-70.
  19. Abrha T, Girma Y, Haile K, Hailu M, Hailemariam M (2016) Prevalence and associated factors of clinical manifestations of vitamin a deficiency among preschool children in asgede-tsimbla rural district, north Ethiopia, a community based cross sectional study. *Arch Public Health* 74: 1-8.
  20. Tariku (2016) Vitamin-A deficiency and its determinants among preschool children. *BMC Res Notes* 323: 3-5.
  21. WHO (2008) Indicators for assessing infant and young child feeding practices. 2008: 1-176.
  22. WHO/UNICEF (2003) Global strategy for infant and young child feeding. 2003: 1-37.
  23. Schemann JF, Banou AA, Guindo A, Joret V, Traore L, et al. (2002) Prevalence of undernutrition and vitamin A deficiency in the Dogon Region, Mali. *J Am Coll Nutr* 21: 381-387.
  24. WHO (2009) Global prevalence of vitamin A deficiency in populations at risk 1995-2005. 2009: 1-68.
  25. Jones G, Steketee RW, Black RE, Bhutta ZA, Morris SS, et al. (2003) How many child deaths can we prevent this year? *Lancet* 362: 65-71.
  26. N Arlappa, A Laxmaiah, N Balakrishna, R. Harikumar, Mallikharjuna Rao Kodavanti, et al (2010) Micronutrient deficiency disorders among the rural children of West Bengal, India. *Ann Hum Biol* 38: 281-289.
  27. BLGK AH (2016) protein energy malnutrition, vitamin A deficiency and night blindness in Bangladeshi children. *J pub Health Nutr* 1: 1-14.
  28. Yalew WA, Abitew DB (2014) Breastfeeding practices of mothers of under five years old children in Bahir Dar City, Ethiopia: a descriptive cross sectional community based survey. *Int J Med Sci Public Health* 3: 532-536.
  29. Haile D, Biadgilign S, Azage M (2015) Differentials in vitamin A supplementation among preschool-aged children in Ethiopia: evidence from the 2011 Ethiopian Demographic and Health Survey. *Public health* 129: 748-754.
  30. Daba G, Beyene F, Fekadu H, Garoma W (2013) Assessment of knowledge of pregnant mothers on maternal nutrition and associated factors in Guto Gida Woreda, East Wollega Zone, Ethiopia. *J Nutr Food Sci* 3: 1-7.

**Copyright:** ©2021 Yidsail Hune, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.