

Soft Tissue Wound Healing by Laser

Mohammad Nazrul Islam*

Head-Biomedical Science and Medical, Biotechnology Department, Shaheed Suhrawardy Medical College and Hospital, Dhaka-1207, Bangladesh

*Corresponding author

Mohammad Nazrul Islam, Head-Biomedical Science and Medical, Biotechnology Department, Shaheed Suhrawardy Medical College and Hospital, Dhaka-1207, Bangladesh, Tel: +880 9130800-19; Ext: 240 (Office), +880 1733381819; E-mail: bdbiomedical@gmail.com

Submitted: 30 Nov 2017; Accepted: 23 Mar 2018; Published: 30 Apr 2018

Abstract

Background: In 1967 a few years after the first working laser was invented, Endre Mester in Semmelweis University Budapest, Hungary wanted to find out if laser might cause cancer. He took some mice, shaved the hair off their backs, divided them into two groups and gave a laser treatment with a low powered ruby laser to one group. They did not get cancer and to his surprise the hair on the treated group grew back more quickly than the untreated group. That was how "laser biostimulation" effects were discovered.

(Effect of laser on hair Growth of mice (in Hungarian). Mester, E. Szende, B. and Tota, J.G. (1967). Kiserl Orvostud 19. 628-631).

Purpose of the work: The effects of pulsed monochromatic light, with fixed pulsations and wavelengths, on the healing of pressure ulcers were evaluated in this prospective, randomized, controlled study.

Method: A placebo-controlled, double-blind study using low energy photon therapy (LLLT) was performed in ten patients with bed sore on the back. Treatment was given three times a week for 10 weeks, using monochromatic (red) optical sources; diode 660nm (GaAl- 660). The patients who were randomized to placebo treatment received sham therapy from an identical-appearing light source from the same delivery system.

Results: Ten patients with bed sore were randomized to receive LLLT or placebo therapy. At the conclusion of the study, the percentage of the initial ulcer area remaining unhealed in the LLLT and placebo groups was 24.4% and 84.7%, respectively ($P = 0.0008$). The decrease in ulcer area (compared to baseline) observed in the LLLT and placebo groups was 193.0 mm² and 14.7 mm², respectively ($P = 0.0002$). One patient dropped out of the study, complaining of lack of treatment efficacy; he was found to be randomized to the placebo group. There were no adverse effects.

Conclusions: In this placebo-controlled, double-blind study LLLT was an effective modality for the treatment of bed sore which were resistant to conventional medical management.

The results are encouraging as pulsed monochromatic light increased healing rate and shortened healing time. This will positively affect the quality of life in elderly patients with pressure ulcers.

Keywords: Bed sore Healing, Soft Tissue Healing, Decubitus Ulcer, Ulcer Healing, Wound Healing, Low Level Laser, Laser Therapy.

Pretreatment photograph of wound showing irregular margins, Necrotic tissue and pus discharge.

Visualization of treatment Progress by LLLT (LED- Ga-Al- As 660 nm) of a patient- (Chronological Picture View of a Laser (LLLT) Treated Patient).

1ST DAY, 24TH OCTOBER/2009

Pretreatment photograph of wound showing irregular margins, Necrotic tissue and pus discharge.



Figure 1: At the end of 1st Week



Figure 2: At the end of 2nd week.



Figure 3a: At the end of 3rd week.



Figure 3b: At the end of 3rd week.



Figure 4: At the end of 4th week.



Figure 5: At the end of 5th week.



Figure 6: At the end of 6th week.



Figure 7: At the end of 7th week.



Figure 8b: At 9th week.



Figure 10a: At 10th week.



Figure 10b: At The End of 10th Week.



Figure 10c: At The End Of 10th Week

References

1. Bipolar Sternomastoid Tenotomy for infantile (Congenital) Torticollis. The Journal of Bangladesh Orthopedic Society 24 (2009).
2. A Case report on treatment of 'Shepherd Crook deformity' in fibrous dysplasia. The Journal of Bangladesh Orthopedic Society 24 (2009).
3. Effects of Low Level Laser (Diode-830nm) Therapy On - Human Bone Regeneration. Journal of Surgical Sciences, Society of Surgeons of Bangladesh (Official Publication) (2011) 15: 34-40.
4. Management of chronic Osteomyelitis of Femur with Interlocking Intramedullary Nail in a Young Man-A Case Report. www.banglajol.info/index.php/JSF. Journal of Science Foundation (2013) 11.

-
5. Low Level Laser Therapy (LLLT) on Soft Tissue Healing: A Case Report with Literature Review. www.banglajol.info/index.php/JSSMC (2013) 5: 117-121.
 6. Open Reduction and Internal Fixation of Displaced Fractures of Lateral Malleolus by tension Wiring and Plating in Bi-malleolar Fractures in Adults. Journal of Shaheed Suhrawardy Medical College (2013) 5: 81-83.
 7. Low Level Laser Therapy (LLLT) On Soft Tissue Healing. ISSN 1728-2152 (2014) 18. Journal of Surgical Sciences (2014) 18: 38-42 @ 2012 Society of Surgeons of Bangladesh.
 8. Effects of Low Level Laser (Diode-830nm) Therapy on - Human Bone Regeneration. Journal of Biotechnology and Biomaterials. (2016) 6 (10th Asia-pacific Conferences Issue-2016, Thailand, Bangkok).

Copyright: ©2018 Mohammad Nazrul Islam. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.