

Species can base their impression of themselves on the number of sensory neurons they have and their IUCN level

Aaron Goetz

University of Nevada Reno: Reno, NV, US

***Corresponding author:**

Aaron Goetz, Species can base their impression of themselves on the number of sensory neurons they have and their IUCN.

Submitted: 18 Oct 2022; **Accepted:** 04 Nov 2022; **Published:** 23 Nov 2022**Citation:** Goetz, A. (2022). Species can base their impression of themselves on the number of sensory neurons they have and their IUCN level. *J Math Techniques Comput Math*, 1(2), 151-153.**Abstract**

Humans have the most sensory neurons of any terrestrial species, 18.83B, with more than twice as many as the runner up western gorilla. There are three aquatic species with more than humans and seven more with more than the western gorilla. The killer whale with more than twice as many as humans, long-finned and short-finned pilot whale with a little less than twice humans. With that many sensory neurons it would be assumed that those would be the species with the most emotional disturbance from anthropogenic influence. There is no data on killer whale endangerment, and both long and short finned pilot whales are LC, ranking them 23rd and 24th most disturbed species. It is important for us to consider what the species means to them in evaluating what species we should care about, it would be beneficial to base that on what species care about themselves the most.

Introduction

Humans have the most sensory neurons of any terrestrial species, 18.83B, with more than twice as many as the runner up western gorilla. There are three aquatic species with more than humans and seven more with more than the western gorilla. The killer whale with more than twice as many as humans, long-finned and short-finned pilot whale with a little less than twice humans. With that many sensory neurons it would be assumed that those would be the species with the most emotional disturbance from anthropogenic influence. There is no data on killer whale endangerment, and both long and short finned pilot whales are LC, ranking them 23rd and 24th most disturbed species.

Methods

Species sensory associated neuron and total neuron quantities from wiki were columnized on an excel. There are 146 species that

have their sensory neuron quantities documented on wiki and 110 of those have their total quantities on wiki. A percentage of sensory neurons was then calculated for each of the 110 species that have both. The 146 species were checked on the IUCN for their level. 26 species are without IUCN document so 120 species have both sensory neuron quantities and an IUCN level. For an IUCN level .01 was used for LC, .25 for NT, .5 for VU, .75 for EN and .99 for CR. That percentage multiplied by the quantity of sensory neurons totals a level of IUCN sense for each of the 120 species. Each of the 120 species levels of IUCN sensory use was percentagized comparatively to the species with the most combined sensory neurons and IUCN level. A weighted level of IUCN sense was also calculated for the 110 species that have both sensory and total neuron quantities, only 92 of those with an IUCN level, multiplying the IUCN level and the percentage of sensory neurons to total.

These variables were then put into R as a MLR using $y_i = \beta_0 + \beta_1 + \beta_2 \dots \beta_i + \epsilon_i$, $\epsilon_i \sim \text{Normal}(0, \sigma^2)$ to determine the role of each of the columns as variables with the response being the total level of IUCN sense. $y = \text{total.level.of.iucn.sense}$

$\text{sens} = \text{sensory.associated.neurons}$

tot= total.neurons

perc= percentage.sens

iucn= iucn.level

name= name

weightediucn= weighted.level.of.iucn.sense

AIC scores were compared for $y_1 \sim \text{sens} + \text{perc} + \text{iucn}$, $y_2 \sim \text{perc} + \text{iucn} + \text{weightediucn}$, $y_3 \sim 1$ (null), $y_4 \sim \text{iucn} + \text{sens} + \text{weightediucn}$, $y_5 \sim \text{weightediucn} + \text{tot} + \text{iucn}$, and $y_6 \sim \text{sens} + \text{iucn}$.

Results

The three species with more sensory neurons than humans has not had their total neurons quantified, so humans have the most total neurons of all neural measured species with, 86B total, while the western gorilla has 33.4B and the orangutan 32.6B. Five of the 146 species that have sensory neuron totals are CR with the IUCN

an eight are EN. Of the 14 species with the most disturbed ranked (having the most neurons combined with IUCN level), 5 are VU, 6 EN and 3 CR. The species with the most disturbed rank is an EN, the blue whale. After the ninth ranked species the percent of IUCN sense goes less than half of the top ranked species.

Table 1: Top 28 most disturbed species IUCN sense = sensory associated neurons x IUCN level

1	name	sensory associated neurons	total neurons	percentage sens	iucn level	weighted level of iucn sens	total level of iucn sense	percent of sense
2	Blue whale	15,000,000,000			0.75		11250000000	1
3	Western gorilla	9,100,000,000	33400000000	0.27245509	0.99	0.269730539	9009000000	0.8008
4	Orangutan	8,300,000,000	32600000000	0.254601227	0.99	0.252055215	8217000000	0.7304
5	Fin whale	15,000,000,000			0.5		7500000000	0.666666667
6	Chimpanzee	7,400,000,000	28000000000	0.264285714	0.75	0.198214286	5550000000	0.493333333
7	African elephant	5,600,000,000			0.99		5544000000	0.4928
8	Pygmy chimpanzee or bonobo	7,250,000,000			0.75		5437500000	0.483333333
9	Asian elephant	6,775,000,000	2.57E+11	0.026361868	0.75	0.019771401	5081250000	0.451666667
10	Walrus	3,929,000,000			0.5		1964500000	0.174622222
11	Guenon	2,500,000,000			0.75		1875000000	0.166666667
12	Mandrill	3,102,000,000			0.5		1551000000	0.137866667
13	Hyacinth macaw	2,944,000,000			0.5		1472000000	0.130844444
14	Pigtail Macaque	2,531,000,000			0.5		1265500000	0.112488889
15	Kea	1,281,000,000	2149000000	0.596091205	0.75	0.447068404	960750000	0.0854
16	Horse	1,200,000,000			0.75		900000000	0.08
17	Giraffe	1,731,000,000	10750000000	0.161023256	0.5	0.080511628	865500000	0.076933333
18	Tufted capuchin	1,140,000,000	3691000000	0.308859388	0.75	0.231644541	855000000	0.076
19	Bonnet macaque	1,660,000,000	3780000000	0.439153439	0.5	0.21957672	830000000	0.073777778
20	Grey parrot	850,000,000	1566000000	0.542784163	0.75	0.407088123	637500000	0.056666667
21	Raccoon	453,000,000	2148000000	0.210893855	0.99	0.208784916	448470000	0.039864
22	Crab-eating macaque	800,960,000	3440000000	0.232837209	0.5	0.116418605	400480000	0.035598222
23	Long-finned pilot whale	37,200,000,000			0.01		372000000	0.033066667
24	Short-finned pilot whale	35,000,000,000			0.01		350000000	0.031111111
25	Tarsius	310,000,000			0.99		306900000	0.02728
26	Lion	545,240,000	4667000000	0.116828798	0.5	0.058414399	272620000	0.024232889
27	Risso's dolphin	18,750,000,000			0.01		187500000	0.016666667
28	Goeldi's marmoset	357,130,000	636,000,000	0.561525157	0.5	0.280762579	178565000	0.015872444

The results from the AIC scores y1-y6 have y4 with the best covariates, with y1 only 45.11795 away (the others are more than 5,000 away). This shows that combining the iucn and the sens variables are most effective in the 3 variable equations.

Table 2: $\epsilon_4 \sim \text{Normal}(0, \sigma^2)$

```

Residual standard error: 8.334e-07 on 8 degrees of freedom
(55 observations deleted due to missingness)
Multiple R-squared: 1, Adjusted R-squared: 1
F-statistic: 3.453e+30 on 82 and 8 DF, p-value: < 2.2e-16

```

lm(formula = y ~ iucn + sens + weightediucn)

```

Residuals:
    Min       1Q   Median       3Q      Max
-1.574e-06  0.000e+00  0.000e+00  0.000e+00  1.574e-06

```

```

Coefficients:
(Intercept)  1.118e+07  1.664e-06  6.719e+12
iucn         1.700e+07  7.943e-05  2.140e+11

```

Discussion

It is important for us to consider what the species means to them in evaluating what species we should care about, it would be beneficial to base that on what species care about themselves the most.

Copyright: ©2022 Aaron Goetz. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.