

Survey of Animal Fauna in Gohayna District at Sohag Governorate, Egypt

Abd El-Aleem S. S. Desoky, Mohamed M. Abd-Allah

Plant Protection Department, Faculty of Agriculture, Sohag University, Egypt.

*Corresponding author

Abd El-Aleem S. S. Desoky, Plant Protection Department, Faculty of Agriculture, Sohag University, Egypt.

Submitted: 30 Aug 2021; Accepted: 13 Sep 2021; Published: 25 Sep 2021

Citation: Abd El-Aleem S. S. Desoky, Mohamed M. Abd-Allah. (2021). Survey of Animal Fauna in Gohayna District at Sohag Governorate, Egypt. *J Vet Heal Sci*, 2(1), 12-14.

Abstract

The present work was aimed to identify of animal fauna in Gohayna District, Sohag Governorate, Egypt during 2020/2021season. The results were revealed the presence of four species of rats included the white bellied rat, *Rattus frugivorus* the dominant species from, *Rattus alexandrines*, the Nile grass rat, *Arvicanthis niloticus*. *Rattus norvegicus* and *Gerbillus* sp. Also, found that *Mustela* sp., and long-eared hedgehog, *Hemiechinus auritus*. These results to be used in the development of a future plan in effective strategy for implementation of animal pest's management programs in cultivated in Sohag Governorate.

Keywords: *Arvicanthis Niloticus*, *Rattus R. Alexandrines*, *Rattus R. Frugivorus*, *Rattus Norvegicus* *Mustela* sp., *Hemiechinus Auritus*

Introduction

Rodentia is one of the most important mammalian orders which has a great number of rodent species with their effect on the environment. Directly, through their destructive feeding habits and indirectly by a stable food items for many predators in the food chains. In Egypt changes in the agro-ecosystem, during the last 40 years, have had a great effect on the distribution and abundance of field rodent population (El-Sherbiny, 1987). Rodents are implicated in many types of damage, including crop and tree damage, structural property and cable damage, disease transmission [1].

Rodents are known for their high reproductive potential; however, there is much variability among species as to the age at first reproduction, size of litters, and the number of litters per year. Under favorable conditions, populations of some species such as the microtines (e.g., voles) can irrupt, going from less than 100 per ha to several thousand per ha in the period of a few months [2]. There are many interesting dynamics to various rodent populations that should be understood to better facilitate their management and to reduce damage. The population goes through an annual cycle that may include high and low densities, active and inactive periods, reproductive and non-reproductive periods, and dispersal periods. To avoid inclement periods, some species exhibit winter dormancy (hibernation), and some species have summer dormancy (estivation) during hot, dry periods. Some species exhibit multi-year cycles; for example, the microtines often reach population peaks (irruptions)

every 3-5 years [3, 4].

All rodents require food, shelter, and water. The shelter provides protection from predators, inclement weather, and a favorable place to bear and rear their young. Although rodents require water, those water requirements vary greatly by species. Because rodent food and cover (i.e., vegetation) can be influenced by human activities, there has been considerable development of strategies to reduce populations and damage by manipulating vegetation. We will discuss some of these habitat management approaches, but caution that many of them have not been thoroughly investigated or tested on a large scale [5]. The present work was aimed to identify of animal fauna in Gohayna District, Sohag Governorate, in Upper Egypt, to be used in the development of a future plan in effective strategy for implementation of animal pest's management programs in newly reclaimed land in Egypt.

Materials and Methods

Location of the study site:

- The study sites in Gohayna district northward away from Sohag Governorate, about 50 kilometers.

Animal pest's species were collected from the above-mentioned sites by applying the common wire traps. Each trap was baited by bread and distributed twice every 15 days at 6 pm Next morning at 7 am during study period, traps were checked and animals were

identified and recorded for data processing. The captured of animal pests were classified and recorded.

Results and Discussion

Data in Table 1 show the species composition of rodents trapped from two different areas in animal fauna in Gohayna district, Sohag Governorate, Egypt. Species recorded were the white bellied rat, *Rattus frugivorus*, the grey bellied rat, *Rattus alexandrinus* and the Nile grass rat, *Arvicanthis niloticus*. Norway rat, *Rattus norvegicus* and *Gerbillus* sp. Also, found that *Mustela* sp., and long-eared hedgehog, *Hemiechinus auritus*.

In the Houses region found that the presence of four species of rats included the white bellied rat, *R. r. frugivorus* the dominant species from the *Rattus alexandrinus*, Nile grass rat, *A. niloticus*, *Rattus r. frugivorus* and *Mustela* sp., at the study area. *Rattus r. frugivorus* the dominant species, this may be due to the presence of attributed to the availability of food and shelter as well as prefers trees for nesting in houses. Also, this may be due to the inter-specific competition between this species and other species.

On the other hand, in the fields crops the Nile grass rat, *A. niloticus* the dominant species from *R. r. frugivorus*, *Rattus r. alexandrinus*, and its recorded long-eared hedgehog, *Hemiechinus auritus* However, *R. norvegicus* was not captured by the traps

The results similar with Ali (1985) recorded six species of rats and mice in Sohag Governorate [6]. The species density percentages were arranged quantitatively in the following descending order *R. norvegicus* (35.17%), *A. niloticus* (19.86%), *R. r. frugivorus* (19.39%), *R. r. alexandrinus* (13.88%), *M. musculus*, (11.00%), *Acomys cahirinus* (0.72%).

Abdel-Gawad (1987) found those, three rodent species *A. niloticus* (Desm.), *R. r. frugivorus* and *Gerbillus* sp. Desm., in Wady El-Assiuty area, Assiut Governorate [7]. He noticed that, *A. niloticus* (Desm.) preferred areas planted with the field crops and some parts under cultivation beside old cultivated land while *R. r. frugivorus* was found in citrus orchards and around fanners buildings, whereas *Gerbillus* sp. Desm., lived in desert and semi-desert parts where wild plants and weeds were grown.

Desoky et al., (2014) finding agrees with the results show in the experimental station of the Faculty of Agriculture [8], El-Kawther city, Sohag University, found that the presence of three species of rats included the Lesser garbia, *Gerbillus* sp. was recorded (1.08%) from newly reclaimed area; the Nile grass rat, *A. niloticus* (4.44%). This may be attributed to the availability of food in neighbored field crops and vegetables plantations also, the white bellied rat, *R. r. frugivorus* the dominant specie (94.27 %.) This may be due to several factors e.g., intra-specific competition, fecundity increasing and in habitat the ecosystems in which poultry buildings established in the faculty farm the presence of palm trees in the preparation of farm animal production, or poultry farm nearby, this provides shelter and increase in feed stores.

The differences in species composition of rodents depending on locality, neighboring, habitat type, inter specific compotation and preferred food [8]. Identification of rodent species in the study

area can be used in the development of a future plan in effective strategy for implementation of rodent management programs in newly reclaimed land in Egypt [9-11].

Table 1: List of animal fauna collected in Gohayna district, Sohag Governorate in Upper Egypt during 2020/2021 season.

Animal fauna	Study area		Common name
	Houses	Fields	
<i>Rattus frugivorus</i>	+++	++	white bellied rat, date palm rat
<i>Rattus alexandrinus</i>	++	+	the grey bellied rat
<i>Arvicanthis niloticus</i>	+	+++	Field rat, grass rat, Nile rat, Nile grass rat
<i>Rattus norvegicus</i>	+	+	street rat, sewer rat, Norway rat, brown Norway rat, Norwegian rat.
<i>Gerbillus</i> sp	-	+	Lesser garbia
<i>Mustela</i> sp	+	-	weasel
<i>Hemiechinus auritus</i>	-	+	long-eared hedgehog

+++ = High population
 ++ = Moderately population
 + = Slightly population
 - = Absent

References

1. Witmer, G. W., Campbell III, E. W., & Boyd, F. (1998, January). Rat management for endangered species protection in the US Virgin Islands. In Proceedings of the Eighteenth Vertebrate Pest Conference (1998) (p. 22).
2. O'brien, J. M. (1994). Voles. Pages B177–B182 in S. Hygnstrom, R. Timm, and G. Larson, editors. Prevention and control of wildlife damage. Cooperative Extension Service, University of Nebraska, Lincoln, NE, USA.
3. Edge, W. D., Wolff, J. O., & Carey, R. L. (1995). Density-dependent responses of gray-tailed voles to mowing. The Journal of wildlife management, 245-251.
4. Wolff, J. O., Schaubert, E. M., & Edge, W. D. (1997). Effects of Habitat Loss and Fragmentation on the Behavior and Demography of Gray-Tailed Voles: Efectos de la Pérdida y Fragmentación de Hábitat Sobre el Comportamiento y la Demografía de Ratones de Campo de Cola Gris. Conservation Biology, 11(4), 945-956.
5. Barras, S. C., & Seamans, T. W. (2002). Habitat management approaches for reducing wildlife use of airfields. In Proceedings of the Vertebrate Pest Conference (Vol. 20, No. 20).
6. Ali, M. K. (1985). Studies on rodents and their ectoparasites in Sohag Governorate (Doctoral dissertation, M. Sc. Thesis Fac. Agric., Assiut Univ).
7. Abdel-Gawad, K. H., & Maher Ali, A. (1987). Seasonal

-
- distribution of rodent species and their associated ectoparasites in the new cultivated land. *Egyptian Journal of Wildlife and Natural Resources*, 9, 1-11.
8. Desoky, A., Baghdadi, S., & Ahmed, H. (2014). POPULATION DENSITY AND SEASONAL DISTRIBUTION OF RODENT SPECIES AT SHEEP FARMING IN EL-KAWTHER REGION, SOHAG GOVERNORATE, EGYPT. *Journal of Plant Protection and Pathology*, 5(10), 903-907.
 9. Desoky, A. S. S., Ali, A. M., Abd El-Gawad, K. H., & Abdel-Galil, F. A. (2006). Management strategies for rodents within different ecosystems (Doctoral dissertation, Assiut University).
 10. El-Sherbiny, A. H. (1987). Cyclic fluctuation in rodent population: Review of current researches. *Egypt wild and not resources*, 9, 17.
 11. Abdel-Gawad, K. H. (2010). Rodent species composition in the present compared with past, the 5 Scientific Conferences for Agric. Assiut Univ. Oct, 16-17.

Copyright: ©2021 Abd El-Aleem S. S. Desoky, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.