

## The Biophysical Modeling of the Nephrogenesis in the Perinatal Periods

Janos Vincze, Gabriella Vincze-Tiszay

Health Human International Environment Foundation,  
Budapest, Hungary

### Corresponding author

Janos Vincze, Gabriella Vincze-Tiszay, Health Human International Environment Foundation, Budapest, Hungary. Email: [ndp@t-online.hu](mailto:ndp@t-online.hu)

Submitted: 15 July 2020; Accepted: 20 July 2020; Published: 29 July 2020

### Abstract

Our experiments analyzed the microscopic picture of kidneys of patients who died during gestational age or infancy. We wanted to investigate nephrogenesis in its ontogenetic development. We only conducted the analysis until the age of three, because the one million nephron number characteristic of the kidney in adulthood is already developed by the age of three. This perinatal time interval was divided into five periods. Intrauterine are the early fetal (3–5 months) and late fetal periods, neonates (0–1 months), infants (1–12 months) and young children (1–3 years). The specific case numbers are not taken into account, only general remarks are communicated. We found that the structural and functional unit of the kidney is the nephron, which undergoes three initial stages: pro-, meso- and metanephros. There are two periods of accumulation: late fetal and infant age. After birth, the number of incomplete renal corpuscles is negligible. Nephrogenesis occurs from the outside to the inside. By the age of three, the volume of the cortex doubles that of the marrow. In further periods of ontogenesis, the nephron count remains unchanged, only the volume of the nephrons increases. When modelling kidney function, we found that the body is “resting” some of the nephrons, because it is during this time that the cell walls resulting from cavitation are regenerated.

**Key words:** Nephrogenesis, Biophysical Modelling, Perinatal Periods, Ontogenetic Development

### Introduction

It is an old finding that the more an organ works, the higher its oxygen consumption. The kidneys, which weigh less than two percent of the body (about 150 grams), require one tenth of the body's total oxygen consumption. To meet this need, 800 liters of blood flow daily through the two kidneys. The kidney excretes excess substances through urine: decomposition products of substances foreign to the metabolism and organism (medicines, nutritional supplements). The kidneys also play an important regulatory role [1] (fluid and water balance, pH, blood pressure, hematopoiesis [2], bone metabolism). The kidney regulates certain ions of the body ( $\text{Na}^+$ , K and chlorides) and its total water content, and thus regulates the osmotic pressure [3] of the extracellular space. In addition to regulating ionic balance, the kidney also plays an important role in the excretion of metabolites resulting from metabolism. The end product of proteolysis is urea (residual nitrogen), while the end product of nucleic acid decomposition is uric acid. Creatinine is the metabolite of muscle metabolism. The kidneys also play an important role in the excretion of many drugs and other foreign substances and in the elimination of red blood cell metabolites (bilirubin) in liver disease. In this article we do not deal with several types renal failure [4, 5], including kidney cancer.

### The prenatal periods

We distinguish two main stages of the ontogenesis of the individual: pre- and postnatal development; while within these we know several major periods. Initial signs of the final form of the kidneys

are already evident on the 8<sup>th</sup> week of embryonic development, i.e. they are characteristic of the early fetal period (3–5 intrauterine month), whereas the final development of the kidneys (the number of nephrons characteristic of the kidney is constant) is completed approx. by age 3. The kidneys of dead fetuses and young children were used to create sections and examined for microscopic images. Based on these fact, the following 5 major developmental periods were studied to monitor the microscopic structure of the kidneys: early fetal (3–5 embryonic months), late fetal (6–9 embryonic months) periods, and neonates (0 to 30 days after birth), infancy (1–12 months) and infancy (years 1–3).

During ontogeny, three renal organs develop relatively sequentially in time and space: pro-, meso- and metanephros. At week 3, the most cranial renal organ develops in the embryo from somite 8–10 on week 3, and it persists for 10 to 14 days. Before the involution of pronephros, mesonephros, which develops from the epithelium of the somatic meso-nephrene 11–30, appears on week 4. [6] The involution of the mesonephros begins in month 2, progresses in the cranio-caudal direction, and ends in month 4. Metanephros is the renal organ developing in the most caudal way, appearing on week 7–8 on the area of somite 31–32.

The morphological and functional unit of metanephros is the nephron that consists of:

- 1.) Malpighian corpuscle
- 2.) Primary (proximal) convoluted tubules

- 3.) The ascending and descending limb of loop of Henle
- 4.) Secondary (distal) convoluted tubules
- 5.) intercalary tubule

At the beginning of the early fetal period there is no clear separation between the cortex and the marrow, and the structure of the lobes in the cortex is incomplete. In the third month, in the cortex - which is approx. one fifth of the marrow - only the outer zone is observed, the inner zone appears only in the fourth month. The outer zone of the cortex contains a large number of small renal corpuscles, the inner zone is characterized by a small number of large glomeruli, while the middle zone is characterized by transient sized renal corpuscles.

In the outer zone, the differentiation of the metanephrogenic tissue to the tubules and vesicles formed by these cells is observed, i.e. this is the nephrogenic zone. The tubules show intrusions, which are considered to be the initiators of the Malpighian corpuscles. These are called incomplete renal corpuscles. This incomplete form dominates the outer zone of the cortex. These renal corpuscles are grouped multifocally. Completely formed renal corpuscles are found in both the outer and inner zones, but their number is much higher in the inner zone than in the outer layer. During this period, metanephros globally contains more incomplete than complete renal corpuscles.

The cortex and the marrow is structured for the late fetal period, the linear ratio of the two is 1:4. In the cortex, the three zones can be clearly demarcated. The incomplete renal corpuscles are multifocally located in the outer zone. Transient forms between incomplete and complete renal corpuscles are also observed, in which the Bowman's capsule completely surrounds the glomerulus but the vascular pole has not yet formed. The middle and inner zones are clearly dominated by complete renal corpuscle, in which the capillary loops are fully developed, and their lumen displays red blood cells. The proximal and distal convoluted tubules found near renal corpuscles can be distinguished by staining procedures. One of the most characteristic features of the late fetal period is the significant increase in the number of nephrons, so this stage of the ontogeny of methanephros can be called the first accumulation period [7].

### The postnatal periods

Birth is the only compatible big leap in an individual's life. In the neonatal period, the ontogenesis of metanephros represents a tremendous change in quality, since during the intrauterine development, besides the kidney, the placenta also played a role, whereas after birth, the kidney alone fulfills this function. In the functional realization of the homeostasis of the internal environment, the hydro electrolytic balance is substantially modified, taking into account that the newborn must adapt to the gaseous medium rather than the liquid medium. During this period, the water balance is extremely unstable, as the kidney has not reached full maturity, neither structurally nor functionally, and its antidiuretic hormone response is also poor. The regulatory mechanisms that regulate fluid balance are extremely fragile, and diseases that disrupt fluid balance are more common. It follows from this that the newborn has to adapt to the changed conditions extremely quickly after birth, and that is why functional changes are far more significant than anatomical changes.

The maturation process of the nephrons accelerates over time, which explains why the proportion of incomplete renal corpus-

cles is significantly reduced compared to the previous period. This morphological structuring of the kidney can be understood as a response to altered functional requirements. During this period, the linear ratio of the cortex to the marrow is 1:3, and the lobes and lobuli are clearly structured. The diameter of the lumen of the canals increases from the outside to the inside of the marrow, whereas in earlier periods it was characterized by an approximately uniform diameter. One of the most important features of this period is that this is the last stage in which incomplete renal corpuscles are still present. [8] Infant age is characterized primarily by morphological changes, including easily distinguishable cortical zones, well-structured lobes and lobuli, and the exclusive presence of complete renal corpuscles.

Despite the fact that only complete renal corpuscles exist, kidney development has not yet been completed. The reasons for the apparent absence of incomplete renal corpuscles are: the relative long duration of infancy (330 days) compared to the maturation time of the nephrons (approximately 1-2 days); the large number of pre-existing renal corpuscles compared to the developing ones (approx. 1,000: 1). So, out of 500,000 to 1,000,000 renal corpuscles, only 1 to 10 are in an incomplete state at a given moment. [9] This period is characterized by a significant increase in the number of nephrons per unit area, so this period can be called the second accumulation phase. This accumulation is of such extent that it represents a significant increase compared to the former, but the kidney mass triples only by the age 3 compared to the newborn kidney, while the body weight triples by the age of 1. During the infant period, the period is primarily characterized by dimensional growth of the nephrons, while numerical growth is secondary. At this age, microscopic images of the kidney show a striking resemblance to those of the adult kidney. The cortex and marrow and their elements are well structured and clearly distinguishable. The ratio of cortex to marrow is 1:2, which is characteristic of adulthood as well.

Infant age is characterized primarily by morphological changes, including easily distinguishable cortical zones, well-structured lobes and lobuli, and the exclusive presence of complete renal corpuscles. Table I.

**Table I. The characteristics data of kidney**

Periods	Kidney-volum (cm3)	Ratio of cortex/marrow	Nefron number
Neonatal	6,5	1,37/1	300.000
Infancy	21,0	1,37/1	650.000
Little child	55,0	2/1	1000.000

In this period, the total number of nephrons in the kidneys reaches one million, and this number does not change in adulthood, but after the age of 60, the total number of nephrons decreases by approx. 1% per year. [10] Nephrons are formed in a multifocal, asynchronous, spatially and temporally unequal manner. Asynchronism of nephrogenesis is indicated by two accumulation periods of nephron growth. Multifocal character and asynchronism, as well as sequential location, indicate the morphological heterogeneity of the kidney, resulting in functional heterogeneity.

Human kidney development in perinatal periods is influenced by three important parameters: periods of ontogenesis, gender and zone-based localization. [11] The ontogenetic development of the

---

kidney, in the light of the nephron number, in the perinatal periods is strongly influenced by age (ontogenetic periods), moderately influenced by the distribution based on zones, while gender has a less significant effect on nephrogenesis.

### References

1. Rose B D (1989) Clinical Physiology of Acid-Base and Electrolyte Disorders. McGraw Hill Book Co., N. Y, 1989.
2. Kashtan C E, Segal Y (2011) Genetic disorders of glomerular basement membranes. Nephron Clin. Pract. 118: c9.
3. Thompson C J, Baylis P H (1987) Thirst in diabetes insipidus: clinical relevance of quantitative assessment. Q. J. Med. 65: 853-862.
4. D'Agati V D, Kaskel F J, Falk R J (2011) Focal segmental glomerulosclerosis. N. Eng. J. Med 365: 2398.
5. Kambham N (2012) Postinfectious glomerulonephritis. Adv. Anat. Pathol 19: 338.
6. Vincze J (2018) Medical Biophysics. NDP P, Budapest.
7. Vincze J (2015) The Biophysics is a Borderland Science. Second Ed. NDP P, Budapest.
8. Vincze J (2007) Interdisciplinarity, NDP P, Budapest.
9. Vincze J (2020) Biophysics of the Human Apparatus, NDP P, Budapest.
10. Vincze J (2006) Biophysics of the structure and of the function in the human kidneys onto-genesis. NDP P, Budapest.
11. Vincze J (2017) Biophysics of the Excretory Apparatus. Second Ed, NDP P, Budapest.

**Copyright:** ©2020 Janos Vincze. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.