

The Mathematics of Extratropical Cyclones Originating in the South of the South American Continent

Ricardo Gobato^{1*}, Abhijit Mitra² and Sana Ahmed³

¹Green Land Landscaping and Gardening, Seedling Growth Laboratory, Brazil

²Department of Marine Science, University of Calcutta, 35 B. C Road, Kolkata, India

³Department of Oceanography, Techno India University, West Bengal, EM 4 Salt Lake, Sector V, India

*Corresponding Author

Ricardo Gobato, Green Land Landscaping and Gardening, Seedling Growth Laboratory, Brazil.

Submitted: 2024, Mar 20 Accepted: 2024, Apr 08 Published: 2024, Apr 22

Citation: Gobato, R., Mitra, A., Ahmed, S. (2024). The Mathematics of Extratropical Cyclones Originating in the South of the South American Continent. *Space Sci J*, 1(1), 01-08.

Abstract

Extratropical cyclones are common in the South Atlantic. They generally arise with the passage of cold fronts to the south of the South American continent, crossing the south of Chile and Argentina, in the regions of Puerto Natales, Punta Arenas (Chile), Rio Galegos, Rio Grande, El Calafate and Ushuaia (Argentina). The extratropical cyclone analyzed presented at its peak the very characteristic shape of a Chinese dragon. Many others were also analyzed. They present a mathematical form of a double Cotes Spiral curve. Here called the Dragonhead cyclone (DC), indicate a structural similarity with spiral galaxies, especially NGC 5247, in the constellation Virgo. With an area of influence and a size of around 3,247 thousand km² at its peak, it moved quickly in a WNW direction, with an average speed of 76 km/h, with winds of 84 km/h at 100 km from the nucleus, as it passed north of the Falkland Islands (Malvinas Islands). In the data collected (Gobato et al., 2018-2023), and analyzed from the Dragonhead cyclone, it is clear that all extratropical cyclones that appear south of the South American continent, below 40° latitude, have the shape of a spiral curve, like the spiral galaxy. Most of these are in the form of a double Cote's spiral curve.

Keywords: Cyclones, South American, Spiral Galaxy, NGC 5247, Falkland Islands, Dragonhead

1. Introduction

1.1 Cyclone

A cyclone is a large air mass that rotates around a strong center of low atmospheric pressure, counterclockwise in the Northern Hemisphere and clockwise in the Southern Hemisphere as viewed from above (opposite to an anticyclone). A subtropical cyclone is a weather system that has some characteristics of a tropical cyclone and some characteristics of an extratropical cyclone [1-4,27,29]. They can form between the equator and the 50th parallel. These storms usually have a radius of maximum winds that is larger than what is observed in purely tropical systems, and their maximum sustained winds have not been observed to exceed about 32 m/s (64 knots). Subtropical cyclones sometimes become true tropical cyclones, and likewise, tropical cyclones occasionally become subtropical storms [1,5-9,26,27]. Subtropical cyclones in the Atlantic basin are classified by their maximum sustained surface winds: Subtropical depressions have surface winds less than 18 m/s (35 knots), while subtropical storms have surface winds greater than or equal to 18 m/s [9-29,39].

Tropical cyclones are compact, circular storms, generally some 320 km (200 miles) in diameter, whose winds swirl around a

central region of low atmospheric pressure. The winds are driven by this low-pressure core and by the rotation of the Earth, which deflects the path of the wind through a phenomenon known as the Coriolis force. As a result, tropical cyclones rotate in a counterclockwise (or cyclonic) direction in the Northern Hemisphere and in a clockwise (or anticyclonic) direction in the Southern Hemisphere.

1.2 NGC 5247 Spiral Galaxy

Spiral galaxies form a class of galaxy originally described by Edwin Hubble in his 1936 work *The Realm of the Nebulae* and, as such, form part of the Hubble sequence [40-43]. Most spiral galaxies consist of a flat, rotating disk containing stars, gas and dust, and a central concentration of stars known as the bulge. These are often surrounded by a much fainter halo of stars, many of which reside in globular clusters [54].

Spiral galaxies are named by their spiral structures that extend from the center into the galactic disc. The spiral arms are sites of ongoing star formation and are brighter than the surrounding disc because of the young, hot OB stars that inhabit them [42,43,47,54].



Figure 1: HAWK-I image of NGC 5247 Galaxy
Source: [55]

The Figure (1) shows NGC 5247, a grand design barred spiral galaxy, located 60–70 million light-years away. The galaxy lies face-on towards Earth, thus providing an excellent view of its pinwheel structure and multiple arms. It is in the zodiacal constellation of Virgo (the Maiden) [55].

The image was made in infrared light with the HAWK-I camera on ESO's Very Large Telescope at Paranal Observatory in Chile. HAWK-I is one of the most powerful infrared imagers in the world, and this is one of the sharpest and most detailed pictures of this galaxy ever taken from Earth. The filters used were Y (shown here in blue), J (in light blue), H (in green), and K (in red). The field of view of the image is about 6.4 arc minutes [55].

NGC 5247 is a face-on unbarred spiral galaxy located some

60 million light years away in the constellation Virgo. It is a member of the Virgo II Groups, a series of galaxies and galaxy clusters strung out from the southern edge of the Virgo Super cluster. This grand design spiral galaxy displays no indications of distortion caused by interaction with other galaxies. It has two spiral arms that bifurcate after wrapping halfway around the nucleus. The disk is estimated to be 4.9 ± 2.0 kly (1.5 ± 0.6 kpc) in thickness and it is inclined by roughly 28° to the line of sight [56].

1.3 Cote's Spiral

In physics and in the mathematics of plane curves, a Cote's spiral (also written Cotes' spiral and Cotes spiral) is one of a family of spirals classified by Roger Cotes [29-36].




Figure 2: Shapes of curves called Cote's spiral gives the solution to the central orbit problem
Source: [58-61].

2. Development

After an analysis of the different types of spirals, it was concluded that the form that came closest the spiral, Figure (1, 2, 4-7), is a double Cotes's Spiral, for the case $\mu < h^2$, Figure (2) [29-36].

It was determined the mathematical equation of the shape of the

$$\ddot{r} = -\mu[r] - 3\dot{\theta}^2 r \quad (1)$$

where μ is a positive constant. There are three solution regimes,

$$r = A \sec(k\theta + \epsilon) \quad (2)$$

where $k^2 = 1 - \frac{\mu}{h^2}$, when $\mu < h^2$,

where A and ϵ are constants, and h is the specific angular momentum [26,37,38].

Analyzing the shape of the Spiral shape called "Cotes's Spiral." for $\mu < h^2$ [24, 25], it appears that adding two constants to Equation (2) makes the necessary adjustments for the Isobaric ones. In the case of DC, the spiral that gives the solution to a radial force law is given by Equation (1).

An adjustment in Equation (2) is necessary to obtain the graph of Figure (7). Then, adding the constants

$B \neq 0$ and C where for $\mu < h^2$.

$$r = B A \sec(k\theta + \epsilon) + C \quad (3)$$




Figure 3: The Figure (3) shows isobaric and wind lines at 500 m height, in the south of the American continent
Source: [57, Adapted Authors]



Figure 4: Image of the Dragonhead cyclone (DC), at 8 am (UTC), February 7, 2024
Source: [57, Adapted Authors]



Figure 5: Image of the DC, at 12 am (UTC), February 7, 2024
Source: [57, Adapted Authors]




Figure 6: Image of the DC, at 6 pm (UTC), February 7, 2024
Source: [57, Adapted Authors]




Figure 7: Graphic plot of the dimensions and shape of the Dragonhead cyclone, as at 18 am (UTC), February 7, 2024
Source: [Authors]

3. Analysis and Results

With winds coming from the Pacific Ocean, the cold front crosses southern Chile and Argentina. Forming from the region of Puerto Natales, Punta Arenas (Chile), El Calafate, Rio Galegos, Rio Grande and Ushuaia (Argentina), as shown in Figure 1, 4h UTC on February 7, 2024. The cold coming from the Pacific collides with a mass of hot air coming from the north of the province of Santa Cruz and central-south of Chubut.

February 7, 2024	Coordinates	Pressure (hPa)
08:00 am (UTC)	52°46'08"S 67°49'48"W	978
06:00 pm (UTC)	51°33'22"S 65°14'24"W	970

Table 1: Dragonhead cyclone coordinates/Pressure

Source: [57,Authors]

It had a pressure of 978 hPa, February 7, 2024 at 08:00 am (UTC), 100 km north of the nucleus Figure 6, with winds of 76 km/h, with a nucleus at 970 hPa. Taking off in the WNW direction.

It appears that the spiral shape of the Dragonhead is very similar to the shape of the spiral galaxy NGC 5247, presenting the same shape.

The analogous shape of Dragonhead cyclone and the NGC 5247 Galaxy, studied here is clear. These present a double Cote's spiral. Studied by Lindblad, but with the Cote's spiral form, (Gobato et al. 2022) [8,9,11].

The Figure (2) show shapes of curves called Cote's spiral gives the solution to the central orbit problem.

The Table (1) shows the coordinates of Dragonhead cyclone, February 07, 2024, am/pm, a central vortex pressure of 951 hPa, with an approximate dimension of 10 km.

The Figure (3) shows isobaric and wind lines at 500 m height, in the south of the American continent.

The pressure at 500m height in the cities of El Calafate, Puerto Natales, Ushuaia, Rio Grande, Rio Galegos, are 989, 990, 984, 982, 979 and 979 hPa, respectively. In the Falkland Islands (Malvinas Islands) it is between 984 and 990 hPa.

The Figures (4-6) show the trajectory of the dragonhead cyclone between 08:00 am (UTC) and 06:00 pm, February 7, 2024.

The Figure (7) it represents the dimensions of the points collected from Figure (6), in km, using the Isobaric [30,31] found in Figures (3-6).

Other spiral galaxies, vortex storms, have also been analyzed [48-54].

4. Conclusions

Extratropical cyclones are common in the South Atlantic. They generally arise with the passage of cold fronts to the south of

On February 7, 2024 at 18:00 UTC, the cyclone reached an area of influence of 3,247 thousand km². It had a head/neck of elliptical dimensions measuring 639 km by 552 km, whose core was located at 45°36'S, 21°31'W, north of the Falkland Islands. With winds of 84 km/h, it had already traveled 640 km in 10 hours, with an average of 64 km/h.

the South American continent, crossing the south of Chile and Argentina, in the regions of Puerto Natales, Punta Arenas (Chile), Rio Galegos, Rio Grande, El Calafate, and Ushuaia (Argentina). The extratropical cyclone analyzed, presented at its peak the very characteristic shape of a Chinese dragon. Mathematical analyzes of the shape of a double Cotes's Spiral. Here called the Dragonhead cyclone, indicate structural similarity with spiral galaxies, especially NGC 5247, in the constellation Virgo. With an area of influence and a size of around 3,247 thousand km² at its peak, it moved quickly in a WNW direction, with an average speed of 76 km/h, with winds of 84 km/h at 100 km from the nucleus, as it passed north of the Falkland Islands (Malvinas Islands).

In the data collected and analyzed from the Dragonhead cyclone, it is clear that all extratropical cyclones that appear south of the South American continent, below 40° latitude, have the shape of a spiral curve, like the spiral galaxy. Most of these are in the form of a double Cote's spiral curve [6-21].

References

- Gobato, R., & Mitra, A. Double spiral galaxies and the extratropical cyclone in South Georgia and the South Sandwich Islands.
- Elsner, J. B. (2020). Continued increases in the intensity of strong tropical cyclones. *Bulletin of the American Meteorological Society*, 101(8), E1301-E1303.
- Landsea, C. (2009). Subject:(A6) what is a sub-tropical cyclone?. *Atlantic Oceanographic and Meteorological Laboratory*.
- Gobato, R., & Mitra, A. (2021). Tornado in Guatambu, Santa Catarina, Southern Brazil, Late Winter 2021 (Case Study).
- Edwards, R. (2006). The online tornado FAQ. *Storm Prediction Center. National Oceanic and Atmospheric Administration*.
- Gobato, R., Mitra, A., & Valverde, L. (2022). Tornadoes analysis Concordia, Santa Catarina, Southern Brazil, 2022 season. *Aeronautics and Aerospace Open Access Journal*, 6, 184-188.
- Gobato, R., Mitra, A., & Mullick, P. (2023). Extratropical cyclone in the South Georgia and south sandwich islands

- and double spiral galaxies. *Phys Astron Int J*, 7(3), 157-161.
8. Gobato, R., Mitra, A., Heidari, A., & Risso Gobato, M. R. (2022). Spiral galaxies and powerful extratropical cyclone in the Falklands Islands. *Physics & Astronomy International Journal*, 6(2), 48-51.
 9. Gobato, R., Mitra, A., Heidari, A., & Risso Gobato, M. R. (2022). Spiral galaxies and powerful extratropical cyclone in the Falklands Islands. *Physics & Astronomy International Journal*, 6(2), 48-51.
 10. Gobato, R., Mitra, A., Heidari, A., & Gobato, M. R. R. (2022). Extratropical cyclone in the falklands islands and the spiral galaxies. *Sumerianz Journal of Scientific Research*, 5, 32-42.
 11. Gobato, R., Mitra, A., & Mullick, P. (2023). Double Spiral Cote's Galaxies NGC 1566 and the Southern Pinwheel, and Cyclone in the South Sandwich Islands. *American Journal of Astronomy and Astrophysics*, 10(1), 1-8.
 12. Gobato, R., Mitra, A., & Heidari, A. (2022). Vortex storms in the west of santa catarina. *Biomedicine and Chemical Sciences*, 1(2), 41-46.
 13. Gobato, R., Heidari, A., & Mitra, A. (2020). Mathematics of the Extra-tropical Cyclone Vortex in the Southern Atlantic Ocean. *J Climatol Weather Forecast*, 9, 268.
 14. Bluestein, H. B. (2013). Severe convective storms and tornadoes. *Springer*, 10, 978-3.
 15. Gobato, R., Gobato, M. R. R., & Heidari, A. (2018). Evidence of tornadoes reaching the countries of rio branco do ivai and rosario de ivai, southern brazil on june 6, 2017. *Climatol Weather Forecasting*, 6(4).
 16. Gobato, R., Gobato, M. R. R., & Heidari, A. (2018). Evidence of tornadoes reaching the countries of rio branco do ivai and rosario de ivai, southern brazil on june 6, 2017. *Climatol Weather Forecasting*, 6(4).
 17. Gobato, R., Gobato, M. R. R., & Heidari, A. (2019). Storm Vortex in the Center of Paran State on June 6, 2017: A Case Study. *Sumerianz Journal of Scientific Research*, 2(2), 24-31.
 18. Gobato, R., Heidari, A., Mitra, A., & Gobato, M. R. R. (2020). Vortex cotes's spiral in an extratropical cyclone in the southern coast of brazil. *Archives in Biomedical Engineering and Biotechnology-ABEB*, 4(5), 1-4.
 19. Gobato, R., Heidari, A., Mitra, A., & Gobato, M. R. R. (2020). Vortex cotes's spiral in an extratropical cyclone in the southern coast of brazil. *Archives in Biomedical Engineering and Biotechnology-ABEB*, 4(5), 1-4.
 20. Heidari, A., Gobato, R., Mitra, A., & Gobato, M. R. R. (2020). Cotes's Spiral Vortex in Extratropical Cyclone Bomb South Atlantic Oceans. *Aswan University Journal of Environmental Studies*, 1(2), 147-156.
 21. Gobato, R., Gobato, A., & Fedrigo, D. F. G. (2016). Study of tornadoes that have reached the state of Parana. *Parana J. Sci. Educ*, 2(1), 1-27.
 22. Gobato, R., Mitra, A., Heidari, A., & Gobato, M. R. R. Spiral Galaxies and Extratropical Cyclone.
 23. Casey, J. (1893). *A treatise on the analytical geometry of the point, line, circle, and conic sections: containing an account of its most recent extensions, with numerous examples.* Hodges, Figgis, & Company Limited.
 24. Sharipov, R. (2011). Course of analytical geometry. *arXiv preprint arXiv:1111.6521*.
 25. De Leon, M., & Rodrigues, P. R. (2011). *Methods of differential geometry in analytical mechanics.* Elsevier.
 26. Vasquez, T. (2002). *Weather forecasting handbook.* Garland, TX, USA: Weather graphics technologies.
 27. Bosart, L., & Bluestein, H. (Eds.). (2013). *Synoptic-dynamic meteorology and weather analysis and forecasting: a tribute to Fred Sanders* (Vol. 33, No. 55). Springer Science & Business Media.
 28. Gobato, R., & Heidari, A. (2020). Cyclone bomb hits southern brazil in 2020. *Journal of Atmospheric Science Research*, 3(3), 8-12.
 29. Rafferty, J. P. (Ed.). (2010). *Storms, violent winds, and earth's atmosphere.* Britannica Educational Publishing.
 30. Krasny, R. (1986). A study of singularity formation in a vortex sheet by the point-vortex approximation. *Journal of Fluid Mechanics*, 167, 65-93.
 31. Saffman, P. G. (1995). *Vortex dynamics.* Cambridge university press.
 32. Sokolovskiy, M. A., & Verron, J. (2000). Four-vortex motion in the two layer approximation: Integrable case. *Regular and Chaotic dynamics*, 5(4), 413-436.
 33. Heidari, A., Gobato, R., Mitra, A., & Gobato, M. R. R. (2020). Cotes's Spiral Vortex in Extratropical Cyclone Bomb South Atlantic Oceans. *Aswan University Journal of Environmental Studies*, 1(2), 147-156.
 34. George, J. J. (2014). *Weather forecasting for aeronautics.* Academic press.
 35. Gobato, R., & Mitra, A. (2023). Cote's Spiral in Neptune Great Dark Spot (GDS).
 36. Anderson Jr, J. D. (1991). *Fundamentals of Aerodynamics,* McGraw-Hill. New York.
 37. Gobato, R., Mitra, A., & Mullick, P. (2023). Double Spiral Cote's Galaxies NGC 1566 and the Southern Pinwheel, and Cyclone in the South Sandwich Islands. *American Journal of Astronomy and Astrophysics*, 10(1), 1-8.
 38. Whittaker, E. T. (1964). *A treatise on the analytical dynamics of particles and rigid bodies.* CUP Archive.
 39. Heidari, A., Gobato, R., Mitra, A., & Gobato, M. R. R. (2020). Cotes's Spiral Vortex in Extratropical Cyclone Bomb South Atlantic Oceans. *Aswan University Journal of Environmental Studies*, 1(2), 147-156.
 40. Fischer, R. (1993). *Fibonacci applications and strategies for traders* (Vol. 4). John Wiley & Sons.
 41. Toomre, A. (1977). Theories of spiral structure. In: *Annual review of astronomy and astrophysics. Volume 15.*(A78-16576 04-90) Palo Alto, Calif., Annual Reviews, Inc., 1977, p. 437-478. *NSF-supported research.*, 15, 437-478.
 42. Gobato, R., & Mitra, A. (2023). Cote's Spiral in Neptune Great Dark Spot (GDS).
 43. Nezlin, M. V., & Snezhkin, E. N. (2013). *Rossby vortices, spiral structures, solitons: astrophysics and plasma physics in shallow water experiments.* Springer Science & Business Media.
 44. Gobato, R., Mitra, A., & Mullick, P. (2023). Double Spiral Cote's Galaxies NGC 1566 and the Southern Pinwheel, and Cyclone in the South Sandwich Islands. *American Journal*

-
- of Astronomy and Astrophysics*, 10(1), 1-8.
45. Gobato, R., Mitra, A., & Mullick, P. (2023). Double Spiral Cote's Galaxies NGC 1566 and the Southern Pinwheel, and Cyclone in the South Sandwich Islands. *American Journal of Astronomy and Astrophysics*, 10(1), 1-8.
46. Gobato, R., Mitra, A., & Mullick, P. (2023). Extratropical cyclone in the South Georgia and south sandwich islands and double spiral galaxies. *Phys Astron Int J*, 7(3), 157-161.
47. Lindblad, B. (1964). On the circulation theory of spiral structure. *Astrophysica Norvegica*, v. 9, p. 103, 9, 103.
48. Gobato, R., & Heidari, A. (2020). Vortex hits southern brazil in 2020. *J Cur Tre Phy Res App*, 1(2), 109-112.
49. Mendonça Junior, J. A. M. D. (2023). *Galáxias starburst como fontes de raios cósmicos ultra-energéticos: contribuição para o espectro de energia e composição* (Master's thesis, Universidade Tecnológica Federal do Paraná).
50. Gobato, R., Mitra, A., & Mullick, P. (2023). Double Spiral Cote's Galaxies NGC 1566 and the Southern Pinwheel, and Cyclone in the South Sandwich Islands. *American Journal of Astronomy and Astrophysics*, 10(1), 1-8.
51. Gobato, R., & Mitra, A. Double spiral galaxies and the extratropical cyclone in South Georgia and the South Sandwich Islands.
52. Serebriakova, N., Tkachenko, A., Gebruers, S., Bowman, D. M., Van Reeth, T., Mahy, L., ... & Aerts, C. (2023). The ESO UVES/FEROS Large Programs of TESS OB pulsators-I. Global stellar parameters from high-resolution spectroscopy. *Astronomy & Astrophysics*, 676, A85.
53. Islands, F. (2010). South Georgia and the South Sandwich Islands. *Turks and Caicos Islands*.
54. Gobato, R., & Mitra, A. Double spiral galaxies and the extratropical cyclone in South Georgia and the South Sandwich Islands.
55. Hisaindee, M. N., Mudhawa, A. I., & Suresh, E. S. M. (2022, October). Vulnerability Mapping of Residential Buildings with Respect to Cyclonic Wind in Mauritius Using Remote Sensing and QGIS. In *Proceedings of the 2nd International Symposium on Disaster Resilience and Sustainable Development: Volume 1-Multi-hazard Vulnerability, Climate Change and Resilience Building* (pp. 15-27). Singapore: Springer Nature Singapore.
56. Gobato, R., Heidari, A., Mitra, A., & Gobato, M. R. R. (2020). Vortex cotes's spiral in an extratropical cyclone in the southern coast of brazil. *Archives in Biomedical Engineering and Biotechnology-ABEB*, 4(5), 1-4.
57. Whittaker, E. T. (1964). *A treatise on the analytical dynamics of particles and rigid bodies*. CUP Archive.
58. Fischer, R. (1993). *Fibonacci applications and strategies for traders* (Vol. 4). John Wiley & Sons.
59. Gowing, R. (2002). *Roger Cotes-natural philosopher* (Vol. 50). Cambridge University Press.
60. De Barros, J. A., De Mendonça, J. P. R. F., & Pinto-Neto, N. (2007). Realism in energy transition processes: an example from Bohmian quantum mechanics. *Synthese*, 154, 349-370.

Copyright: ©2024 Ricardo Gobato, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.