

The Role of Type IV Secretion System in Brucella Virulence: A Review

Bayeta Senbeta Wakjira

National Animal Health Diagnostic and Investigation Center,
Sebeta, Ethiopia

*Corresponding author

Bayeta Senbeta Wakjira, National Animal Health Diagnostic and Investigation Center, Sebeta, Ethiopia.

Submitted: 02 Oct 2021; Accepted: 20 Oct 2021; Published: 21 Dec 2021

Citation: Bayeta Senbeta Wakjira. (2021). The Role of Type IV Secretion System in Brucella Virulence: A Review. *J Vet Heal Sci*, 2(1), 30-35.

Abstract

Brucellosis is a bacterial disease of domestic and wild animals caused by the genus *Brucella* which has great public health importance globally. In general, *Brucella* species do not express toxins or virulence factors that cause direct damage to the host. Instead, this pathogen's strategy is to persist long enough in the infected host until transmission can occur, which in the natural hosts is usually through abortion, sexual contact, or shedding of bacteria in milk. One important Brucellavirulence factor for intracellular survival and persistence in the host is the type IV secretion system (T4SS). The type IV secretion system (T4SS) is one of numerous secretion systems used by microbes to transfer macromolecules through the cell membrane, such as proteins and DNA. It's the most versatile secretion system, transporting monomeric proteins, multi-subunit protein poisons, and nucleoprotein complexes, and it's found in Gram-positive and Gram-negative bacteria, as well as some Archaea. The type IV secretion system is a major *Brucella* virulence factor for intracellular survival and host persistence. This review will go through the present state of knowledge on the *Brucella* type IV secretion system, including its architecture and regulation, as well as the newly discovered effector substrates that this system delivers into host cells.

Keywords: Type IV Secretion System, Virulence Factor, Brucella

Introduction

Brucellosis is an economically important disease in production animals worldwide [1] and is one of the most common zoonotic infections. *Brucella* species have evolved to avoid the host's immune system and infection is usually characterized by long-term persistence of the bacteria. One important *Brucella* virulence factor for intracellular survival and persistence in the host is the type IV secretion system.

Type IV secretion systems (TFSS) are a newly identified family of multiprotein complexes that secrete macromolecules [2]. T4SS is encoded by the virB operon in *Brucella*, which is made up of 12 genes (virB1–12) on chromosome II. The promoter upstream of virB1 controls transcription of the vir B operon [3,4]. The vir B operon was discovered in *B. suis* for the first time. Following that, it was discovered to be substantially conserved in all *Brucella* spp. for which genomic sequences were available, indicating that this operon may play a significant role [3]. For a long time, it was assumed that *Brucella* spp. did not have the virulence factors found in other bacteria [5]. However, in recent years, various virulence factors that are essential for infection, including lipopolysaccharide

[6], β -cyclic glucan [7], BvrS/BvrR [8] and outer membrane proteins (Omps) [9] have been identified.

In *Brucella*, VirB T4SS is another major virulence factor that mediates intracellular survival and controls the host immunological response to infection. As a result, the role of the type IV secretion system in *Brucella* virulence factor will be discussed in depth in this review, including current knowledge of architecture and regulation, as well as the newly found effector substrates that this system delivers into host cells.

Regulation of the Brucella T4SS

When *Brucella* enters the host cell, it is exposed to novel environmental circumstances that encourage the production of VirB proteins. T4SS is only required by *Brucella* during specific and relatively short periods of intracellular infection, hence its expression must be tightly regulated. This period begins several hours after the bacterium is taken into a host cell and ends after phagolysosome destruction has been avoided and a replication ready vacuole in the Endoplasmic reticulum has been created. After uptake by host cells, acidification of the *Brucella*-containing

vacuole induces T4SS expression [10].

This occurs after the phagosome transiently fuses with early and late endosomes and lysosomes [11]. Nutrient deprivation could possibly be a signal for virB gene expression. After switching the bacteria from a rich medium to a minimal media at a low pH, virB gene expression can be strongly induced in culture [12]. Around 5 hours after infection of host cells, virB gene expression reaches its peak [13].

Architecture of the Brucella T4SS

The Brucella T4SS is a complex of 12 proteins that assembles in the envelope of the bacterium. Once assembled, this mechanism can transport effector proteins, from Brucella into infected host cells. The virB operon, which includes genes virB1 through virB12 and is regulated by a major promoter upstream of virB1, encodes the Brucella T4SS (Figure 1). The majority of VirB proteins are similar to T4SS components seen in other bacteria, such as Agrobacterium tumefaciens and Bordetella pertussis [3]. Brucella strains lacking a functional T4SS are highly attenuated in macrophages and mice and in the natural host, the goat [3,4,14–16].

Insertions in virB genes or genes affecting virB expression have repeatedly been discovered in transposon mutagenesis screens aimed to uncover Brucella virulence factors [17-21]. This highlights the significance of the T4SS in Brucella pathogenicity.

However, not all VirB proteins are required for the T4SS to function properly. By eliminating each virB gene in *B. abortus* one at a time, it was discovered that virB1, virB7, and virB12 are not required for *B. abortus* persistence in mice [14,16,22]. In Brucella or other bacteria that possess a T4SS, VirB1, VirB7, and VirB12 are not known to be part of the T4SS core translocation apparatus or the pilus (Figure 1). As a result, these proteins may assist or increase T4SS function but are not required. VirB1, for instance, is a lytic transglycosylase that destroys peptidoglycan to make room for the T4SS to form Höppner [23]. However this function could be redundant as the Brucella genome encodes other similar enzymes [14].

The core structure of a T4SS was shown to be composed of the proteins VirB8, VirB9 and VirB10 [24,25]. This appears to be also the case in Brucella as these three proteins interact with each other [26]. Other proteins, such as VirB6 and VirB7, may aid in the development of the core complex by stabilizing it [27,28]. T4SS function is also dependent on the ATPases VirB4 and VirB11, which provide energy for T4SS assembly and transport of effector proteins [29,30]. The major component VirB2 and the minor component VirB5 make up the pilus [31] and assembly of the pilus is mediated by VirB3 [32]. The pilus is thought to attach to the host cell surface and to create a pore, through which effectors are translocated [33].




Figure 1: Schematic representation of the virB operon and the T4SS it encodes [34]

In vitro in macrophages or in vivo in mice, the genes listed in blue are definitely essential for Brucella virulence. In mice, the genes highlighted in orange were determined to be non-essential for virulence. Direct regulators of the virB operon are also displayed,

along with their (approximate) binding areas. Grey indicates putative promoters and intergenic areas containing putative promoters. Substrate proteins are shown in green and green arrows show putative translocation pathways. OM, outer membrane, IM,

inner membrane.

Function of Brucella T4SS during infection

All T4SS vir B genes are preserved in all Brucella species sequenced demonstrating that this system is vital for Brucella. The significance of the T4SS for Brucella virulence has been demonstrated experimentally, largely in vitro in macrophages and in vivo in a mouse infection model [3,4,15,18,19]. Brucella wild-type bacteria can persist and reproduce in both models, however T4SS mutants are slowly eliminated. In vivo, *B. melitensis* and *B. abortus* wild-type bacteria induce innate immune responses in mice at later stages of infection, whereas T4SS mutants do not [35]. This verified prior findings on Brucella stealthy character during infection of their hosts, but it also indicated that later during infection, the T4SS serves as a signal that is recognized by the innate immune system, either directly or indirectly. (for review see [36]) Brucella may actively translocate a molecule into host cells via the T4SS that activates the innate immune system with the purpose of polarizing the immune response to Th1 by increasing interferon gamma production or causing granulomas to develop [36,37].

It's also plausible that a variation in intracellular trafficking and growth causes the difference in immune activation between Brucella wild-type and T4SS-deficient strains. The Brucella T4SS is essential for the maturation of the Brucella phagosome into an ER-derived compartment, according to experiments utilizing cultivated cells [11,38,39]. Despite the fact that Brucella phagosomes transiently fuse with early and late endosomes and lysosomes, a subset of intracellular Brucella with a T4SS is able to exclude endosomal and lysosomal markers from their phagosomes and avoid breakdown in phagolysosomes [11]. Brucella phagosomes, on the other hand, acquire ER markers such as calreticulin. This process of excluding endosomal and lysosomal markers and acquiring ER markers is completed approximately 12 to 24 hours after Brucella infection of a host cell and requires the T4SS and presumably its translocated effectors. Brucella then starts to multiply to high numbers inside host cells, while Brucella virB mutants never reach the ER derived vacuole and are killed in phagolysosomes [11,38,39].

Effectors

The genes encoding Brucella effectors identified to date are scattered across the two Brucella chromosomes (Figure 2). In Bartonella species, the virB genes are located together with the genes encoding the effector substrates [40]. Since a similar situation does

not exist in Brucella, identification of Brucella effectors has proven to be a challenging task.

The first substrates of the Brucella T4SS identified were VceA and VceC [41]. These effectors were found in a screen for Brucella promoters that were activated by VjbR in the heterologous host *E. coli*. Although screening for effector genes among virB-co-regulated genes is a great method of narrowing down potential effector candidates, many candidates could also have been missed. For example regulation of effectors may be under the control of a different regulator (such as directly by BvrR) or regulators downstream of VjbR. Also it could be hypothesized that activation of some effectors is not connected to activation of the virB genes. Brucella may already contain a ready for translocation pool of effectors before entry into host cells, as has been shown for *Legionella pneumophila* [42]. These effectors could be required early during infection of the host cell and expressed constitutively. Since nutrients in the early Brucella phagosome are limited, having a ready pool of effectors would save resources for Brucella for other functions.

Recently, a different strategy to identify Brucella effectors was utilized by screening all proteins of unknown function for eukaryotic-like domains or domains known to be involved in protein-protein interactions [43]. This strategy has proven to be successful in identifying T4SS effectors of other intracellular pathogens, such as *L. pneumophila* and *Coxiella burnetii* [44,45]. Using this strategy, 6 proteins were identified that were translocated into mouse macrophages by *B. abortus*. Translocation into cells of 4 of these Brucella putative effector proteins (BPE123, BPE005, BPE275 and BPE043) was dependent on the VirB T4SS (table 1, where the table).

Recently another protein was found that was translocated in a T4SS-dependent manner into macrophages during infection with *B. abortus*. This protein, named RicA, was found in a screen for Brucella proteins interacting with human proteins predicted to be associated with phagosomes. RicA was demonstrated to interact with Rab2, a GTPase involved in trafficking [46]. Rab2 has been shown before to localize to the Brucella phagosome. Furthermore, it was determined that Rab2 is important for intracellular replication of *B. abortus* [47]. In line with this, RicA, which preferentially binds to GDP bound Rab2, is involved in recruiting this GTPase to the Brucella phagosome [46].




Figure 2: The two chromosomes of *Brucella* species showing the positions of effector genes, virB genes and vjbR in the chromosomes. Effectors, shown in black and blue, have been shown to be translocated by the type IV secretion system into host cells. Those in purple have not been shown to be translocated to date. Genes shown in blue have been shown to be regulated by vjbR [34].

Although T4SS was identified in *Brucella* spp. a proximately 15 years ago, the effectors of T4SS remained uncharacterized until recently. The first two effectors, VirB-co-regulated effector (Vec) A and VecC, were identified while screening for genes whose transcription was co-regulated by the virB operon regulator, VjbR [41]. Screening of the interactions between human proteins and predicted *Brucella* proteins using a yeast two hybrid (Y2H) system led to the identification of the Rab2 interacting conserved protein A (RicA)-Rab2 interaction pair. Subsequently, RicA was identified as a T4SS-dependent effector [46].

Genome wide bioinformatics screening for putative T4SS effector proteins using distinct filtering criteria led to the identification of nine T4SS-dependent effector proteins [43,48].

Bacterial toll-Interleukin receptor (TIR) domain-containing proteins are thought to be involved in the virulence of *Brucella*; screening for TIR-containing proteins in *Brucella* led to the identification of two proteins, *Brucella* TIR protein (Btp) A and BtpB, as effectors that are translocated by T4SS into host cells [49]. Recently, a secretory protein, secreted effector protein A (SepA), which is encoded within a horizontally transmitted region, was confirmed as a novel T4SS substrate [50]. Together, these 15 effectors constitute the repertoire of T4SS substrates in *Brucella* spp. identified to date.

A candidate protein may be identified as a T4SS effector protein if it fulfills two criteria: the protein must be secreted into host cells and the secretion must be through the T4SS machinery. The former may be validated by TEM-1 lactamase or calmodulin-dependent adenylate cyclase (CyaA) assays and the latter, by constructing a T4SS-deficient mutant. All strategies used to identify effectors in *Brucella* are confined by the limited number of potential proteins; similar problems were encountered during the identification of effectors in other bacteria as well. By extending the screening of potential targets to the whole genome, the Dot/Icm T4SS machinery

of *Legionella pneumophila* and *Coxiella burnetii* were identified to secrete approximately 300 and 100 proteins, respectively [45]. Similarly, extending the screening method described above to the whole *Brucella* genome may lead to the identification of many more effectors.

Conclusions

Brucella relies on multiple virulence factors, including the VirB T4SS, to induce chronic illness brucellosis in people and animals. More than 10 years ago, the T4SS was discovered as a significant *Brucella* virulence component. Other significant human bacterial pathogens, such as *Helicobacter pylori* and *Legionella pneumophila*, have been identified to use a similar T4SS to translocate effector proteins into host cells, thus it was assumed that *Brucella* would do the same. Although a direct role for the T4SS in intracellular phagosome trafficking cannot be ruled out, the *Brucella* T4SS has been implicated in the translocation of a variety of potential effectors in infected host cells. Now that several substrates of the *Brucella* T4SS have been identified, the next step will be to determine the role of these putative effectors in intracellular survival of *Brucella*. Effector function may theoretically be divided into many categories depending on the host cell pathways that *Brucella* is known to disrupt, such as intracellular trafficking of the *Brucella* phagosome, host immune response manipulation, and apoptosis inhibition.

Some *Brucella* effectors may have a single function in the host cell, whereas others may have many functions. RicA probably interferes with *Brucella* phagosome trafficking by recruiting the trafficking GTPase Rab2 to *Brucella* phagosomes, which is one of the probable effectors reported thus far [46]. Although Btp1/TopB has not yet been proven to be a T4SS substrate, the putative effector's known role is inflammation inhibition. Future study is likely to uncover new, and potentially surprising, host cell pathways that are manipulated by *Brucella* effectors, as well as explicate their role in *Brucella* infection.

References

1. Corbel, M. J. (1997). Brucellosis: an overview. *Emerging infectious diseases*, 3(2), 213.
2. Baron, C., O'Callaghan, D., & Lanka, E. (2002). Bacterial secrets of secretion: EuroConference on the biology of type IV secretion processes. *Molecular microbiology*, 43(5), 1359-1365.
3. O'Callaghan, D., Cazevieille, C., Allardet-Servent, A., Boschioli, M. L., Bourg, G., Foulongne, V., ... & Ramuz, M. (1999). A homologue of the *Agrobacterium tumefaciens* VirB and *Bordetella pertussis* Ptl type IV secretion systems is essential for intracellular survival of *Brucella suis*. *Molecular microbiology*, 33(6), 1210-1220.
4. Sieira, R., Comerci, D. J., Sánchez, D. O., & Ugalde, R. A. (2000). A homologue of an operon required for DNA transfer in *Agrobacterium* is required in *Brucella abortus* for virulence and intracellular multiplication. *Journal of bacteriology*, 182(17), 4849-4855.
5. Fugier, E., Pappas, G., & Gorvel, J. P. (2007). Virulence factors in brucellosis: implications for aetiopathogenesis and treatment. *Expert Reviews in Molecular Medicine*, 9(35), 1-10.
6. Lapaque, N., Moriyon, I., Moreno, E., & Gorvel, J. P. (2005). *Brucella* lipopolysaccharide acts as a virulence factor. *Current opinion in microbiology*, 8(1), 60-66.
7. Reth, M. (2002). Hydrogen peroxide as second messenger in lymphocyte activation. *Nature immunology*, 3(12), 1129-1134.
8. López-Goñi, I., Guzmán-Verri, C., Manterola, L., Sola-Landa, A., Moriyón, I., & Moreno, E. (2002). Regulation of *Brucella* virulence by the two-component system BvrR/BvrS. *Veterinary microbiology*, 90(1-4), 329-339.
9. Lim, J. J., Kim, D. H., Lee, J. J., Kim, D. G., Min, W., Lee, H. J., ... & Kim, S. (2012). Protective effects of recombinant *Brucella abortus* Omp28 against infection with a virulent strain of *Brucella abortus* 544 in mice. *Journal of veterinary science*, 13(3), 287-292.
10. Lim, J. J., Kim, D. H., Lee, J. J., Kim, D. G., Min, W., Lee, H. J., ... & Kim, S. (2012). Protective effects of recombinant *Brucella abortus* Omp28 against infection with a virulent strain of *Brucella abortus* 544 in mice. *Journal of veterinary science*, 13(3), 287-292.
11. Starr, T., Ng, T. W., Wehrly, T. D., Knodler, L. A., & Celli, J. (2008). *Brucella* intracellular replication requires trafficking through the late endosomal/lysosomal compartment. *Traffic*, 9(5), 678-694.
12. Rouot, B., Alvarez-Martinez, M. T., Marius, C., Menanteau, P., Guilloteau, L., Boigegrain, R. A., ... & Baron, C. (2003). Production of the type IV secretion system differs among *Brucella* species as revealed with VirB5- and VirB8-specific antisera. *Infection and immunity*, 71(3), 1075-1082.
13. Sieira, R., Comerci, D. J., Pietrasanta, L. I., & Ugalde, R. A. (2004). Integration host factor is involved in transcriptional regulation of the *Brucella abortus* virB operon. *Molecular microbiology*, 54(3), 808-822.
14. Den Hartigh, A. B., Sun, Y. H., Sondervan, D., Heuvelmans, N., Reinders, M. O., Ficht, T. A., & Tsolis, R. M. (2004). Differential requirements for VirB1 and VirB2 during *Brucella abortus* infection. *Infection and immunity*, 72(9), 5143-5149.
15. Zygmunt, M. S., Hagijs, S. D., Walker, J. V., & Elzer, P. H. (2006). Identification of *Brucella melitensis* 16M genes required for bacterial survival in the caprine host. *Microbes and infection*, 8(14-15), 2849-2854.
16. Foulongne, V., Bourg, G., Cazevieille, C., Michaux-Charachon, S., & O'Callaghan, D. (2000). Identification of *Brucella suis* genes affecting intracellular survival in an in vitro human macrophage infection model by signature-tagged transposon mutagenesis. *Infection and immunity*, 68(3), 1297-1303.
17. Hong, P. C., Tsolis, R. M., & Ficht, T. A. (2000). Identification of genes required for chronic persistence of *Brucella abortus* in mice. *Infection and immunity*, 68(7), 4102-4107.
18. Delrue, R. M., Martinez-Lorenzo, M., Lestrade, P., Danese, I., Bielarz, V., Mertens, P., ... & Letesson, J. J. (2001). Identification of *Brucella* spp. genes involved in intracellular survival. *Infection and immunity*, 69(12), 3707-3717.
19. Watarai, M., Kim, S., Erdenebaatar, J., Makino, S. I., Horiuchi, M., Shirahata, T., ... & Katamine, S. (2003). Cellular prion protein promotes *Brucella* infection into macrophages. *The journal of experimental medicine*, 198(1), 5-17.
20. Lestrade, P., Dricot, A., Delrue, R. M., Lambert, C., Martinelli, V., De Bolle, X., ... & Tibor, A. (2003). Attenuated Signature-Tagged Mutagenesis Mutants of *Brucella melitensis* Identified during the Acute Phase of Infection in Mice. *Infection and immunity*, 71(12), 7053-7060.
21. Sun, Y. H., Rolán, H. G., Den Hartigh, A. B., Sondervan, D., & Tsolis, R. M. (2005). *Brucella abortus* virB12 is expressed during infection but is not an essential component of the type IV secretion system. *Infection and immunity*, 73(9), 6048-6054.
22. Höppner, C., Liu, Z., Domke, N., Binns, A. N., & Baron, C. (2004). VirB1 orthologs from *Brucella suis* and pKM101 complement defects of the lytic transglycosylase required for efficient type IV secretion from *Agrobacterium tumefaciens*. *Journal of bacteriology*, 186(5), 1415-1422.
23. Chandran, V., Fronzes, R., Duquerroy, S., Cronin, N., Navaza, J., & Waksman, G. (2009). Structure of the outer membrane complex of a type IV secretion system. *Nature*, 462(7276), 1011-1015.
24. Fronzes, R., Schäfer, E., Wang, L., Saibil, H. R., Orlova, E. V., & Waksman, G. (2009). Structure of a type IV secretion system core complex. *Science*, 323(5911), 266-268.
25. Sivanesan, D., Hancock, M. A., Villamil Giraldo, A. M., & Baron, C. (2010). Quantitative analysis of VirB8–VirB9–VirB10 interactions provides a dynamic model of type IV secretion system core complex assembly. *Biochemistry*, 49(21), 4483-4493.
26. Baron, C., Thorstenson, Y. R., & Zambryski, P. C. (1997). The lipoprotein VirB7 interacts with VirB9 in the membranes of *Agrobacterium tumefaciens*. *Journal of Bacteriology*, 179(4), 1211-1218.
27. Jakubowski, S. J., Krishnamoorthy, V., & Christie, P. J. (2003). *Agrobacterium tumefaciens* VirB6 protein participates in formation of VirB7 and VirB9 complexes required for type IV secretion. *Journal of bacteriology*, 185(9), 2867-2878.
28. Watarai, M., Makino, S. I., Fujii, Y., Okamoto, K., & Shirahata, T. (2002). Modulation of *Brucella*-induced macropinocytosis by lipid rafts mediates intracellular replication. *Cellular microbiology*, 4(6), 341-355.

29. Alvarez-Martinez, C. E., & Christie, P. J. (2009). Biological diversity of prokaryotic type IV secretion systems. *Microbiology and Molecular Biology Reviews*, 73(4), 775-808.
30. Schmidt-Eisenlohr, H., Domke, N., Angerer, C., Wanner, G., Zambryski, P. C., & Baron, C. (1999). Vir proteins stabilize VirB5 and mediate its association with the T pilus of *Agrobacterium tumefaciens*. *Journal of bacteriology*, 181(24), 7485-7492.
31. Shirasu, K., & Kado, C. I. (1993). Membrane location of the Ti plasmid VirB proteins involved in the biosynthesis of a pilin-like conjugative structure on *Agrobacterium tumefaciens*. *FEMS microbiology letters*, 111(2-3), 287-293.
32. Hwang, H. H., & Gelvin, S. B. (2004). Plant proteins that interact with VirB2, the *Agrobacterium tumefaciens* pilin protein, mediate plant transformation. *The Plant Cell*, 16(11), 3148-3167.
33. De Jong, M. F., & Tsolis, R. M. (2012). Brucellosis and type IV secretion. *Future microbiology*, 7(1), 47-58.
34. Roux, C. M., Rolán, H. G., Santos, R. L., Beremand, P. D., Thomas, T. L., Adams, L. G., & Tsolis, R. M. (2007). *Brucella* requires a functional Type IV secretion system to elicit innate immune responses in mice. *Cellular microbiology*, 9(7), 1851-1869.
35. De Jong, M. F., Rolán, H. G., & Tsolis, R. M. (2010). Microreview: innate immune encounters of the (Type) 4th kind: *Brucella*. *Cellular microbiology*, 12(9), 1195-1202.
36. Rolán, H. G., Xavier, M. N., Santos, R. L., & Tsolis, R. M. (2009). Natural antibody contributes to host defense against an attenuated *Brucella abortus* virB mutant. *Infection and immunity*, 77(7), 3004-3013.
37. Celli, J., de Chastellier, C., Franchini, D. M., Pizarro-Cerda, J., Moreno, E., & Gorvel, J. P. (2003). *Brucella* evades macrophage killing via VirB-dependent sustained interactions with the endoplasmic reticulum. *The Journal of experimental medicine*, 198(4), 545-556.
38. Celli, J., Salcedo, S. P., & Gorvel, J. P. (2005). *Brucella* coopts the small GTPase Sar1 for intracellular replication. *Proceedings of the National Academy of Sciences*, 102(5), 1673-1678.
39. Schulein, R., Guye, P., Rhomberg, T. A., Schmid, M. C., Schröder, G., Vergunst, A. C., ... & Dehio, C. (2005). A bipartite signal mediates the transfer of type IV secretion substrates of *Bartonella henselae* into human cells. *Proceedings of the National Academy of Sciences*, 102(3), 856-861.
40. De Jong, M. F., Sun, Y. H., Den Hartigh, A. B., Van Dijk, J. M., & Tsolis, R. M. (2008). Identification of VceA and VceC, two members of the VjbR regulon that are translocated into macrophages by the *Brucella* type IV secretion system. *Molecular microbiology*, 70(6), 1378-1396.
41. Kubori, T., Shinzawa, N., Kanuka, H., & Nagai, H. (2010). *Legionella* metaeffector exploits host proteasome to temporally regulate cognate effector. *PLoS pathogens*, 6(12), e1001216.
42. Marchesini, M. I., Herrmann, C. K., Salcedo, S. P., Gorvel, J. P., & Comerci, D. J. (2011). In search of *Brucella abortus* type IV secretion substrates: screening and identification of four proteins translocated into host cells through VirB system. *Cellular microbiology*, 13(8), 1261-1274.
43. Pan, X., Lührmann, A., Satoh, A., Laskowski-Arce, M. A., & Roy, C. R. (2008). Ankyrin repeat proteins comprise a diverse family of bacterial type IV effectors. *Science*, 320(5883), 1651-1654.
44. Chen, C., Banga, S., Mertens, K., Weber, M. M., Gorbaslieva, I., Tan, Y., ... & Samuel, J. E. (2010). Large-scale identification and translocation of type IV secretion substrates by *Coxiella burnetii*. *Proceedings of the National Academy of Sciences*, 107(50), 21755-21760.
45. de Barsey, M., Jamet, A., Filopon, D., Nicolas, C., Laloux, G., Rual, J. F., ... & De Bolle, X. (2011). Identification of a *Brucella* spp. secreted effector specifically interacting with human small GTPase Rab2. *Cellular microbiology*, 13(7), 1044-1058.
46. Fugier, E., Salcedo, S. P., De Chastellier, C., Pophillat, M., Muller, A., Arce-Gorvel, V., ... & Gorvel, J. P. (2009). The glyceraldehyde-3-phosphate dehydrogenase and the small GTPase Rab 2 are crucial for *Brucella* replication. *PLoS pathogens*, 5(6), e1000487.
47. Myeni, S., Child, R., Ng, T. W., Kupko III, J. J., Wehrly, T. D., Porcella, S. F., ... & Celli, J. (2013). *Brucella* modulates secretory trafficking via multiple type IV secretion effector proteins. *PLoS pathogens*, 9(8), e1003556.
48. Salcedo, S., Marchesini, M. I., Degos, C., Terwagne, M., Von Bargen, K., Lepidi, H., ... & Gorvel, J. P. (2013). BtpB, a novel *Brucella* TIR-containing effector protein with immune modulatory functions. *Frontiers in cellular and infection microbiology*, 3, 28.
49. Döhmer, P. H., Valguarnera, E., Czibener, C., & Ugalde, J. E. (2014). Identification of a type IV secretion substrate of *Brucella abortus* that participates in the early stages of intracellular survival. *Cellular microbiology*, 16(3), 396-410.

Copyright: ©2021 Bayeta Senbeta Wakjira. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.