

Viscoelastic or Viscoplastic Glucose Theory (VGT #151): Health and Longevity Originate from Body Weight Control: Two Case Studies of HbA1C Value via Body Weight, Fasting Plasma Glucose Through Body Weight, Postprandial Plasma Glucose Addition Through Diet, and Postprandial Plasma Glucose Reduction Through Post-Meal Exercise using 6 Different Energy Analysis Tools of 1 from Time Domain, 1 from Space Domain, and 4 from Frequency Domain Based on the GH-Method: Math-Physical Medicine (No. 744)

Gerald C Hsu

EclaireMD Foundation, USA

*Corresponding author

Gerald C. Hsu, EclaireMD Foundation, USA

Submitted: 26 Sep 2022; Accepted: 01 Oct 2022; Published: 10 Oct 2022

Citation: Gerald C Hsu (2022). *Viscoelastic or Viscoplastic Glucose Theory (VGT #151): Health and Longevity Originate from Body Weight Control: Two Case Studies of HbA1C Value via Body Weight, Fasting Plasma Glucose Through Body Weight, Postprandial Plasma Glucose Addition Through Diet, and Postprandial Plasma Glucose Reduction Through Post-Meal Exercise using 6 Different Energy Analysis Tools of 1 from Time Domain, 1 from Space Domain, and 4 from Frequency Domain Based on the GH-Method: Math-Physical Medicine (No. 744)*. *J App Mat Sci & Engg Res*, 6(3), 1-8.

Introduction

When using Google search, a reader can easily locate and read the following important general information regarding health:

“Studies have shown that becoming overweight or obese is a major risk factor in developing type 2 diabetes. Today, roughly 30 percent of overweight people have diabetes, and around 85 percent of diabetics patients are overweight.

Having diabetes means you are more likely to develop heart disease, such as cardiovascular diseases (CVDs) or strokes. People with diabetes are also more likely to have certain other risk factors, such as high blood pressure or high cholesterol, that increase their chances of having a heart attack or a stroke.

Heart disease is common in people with diabetes. Data from the National Heart Association from 2012 shows 65% of people with diabetes will die from some sort of heart disease or stroke.

During Y2020, the total amount of US deaths was 2,506,540 (100%) which includes a new category of 350,831 (14%) from COVID. The subtotal death cases that directly or indirectly resulted from a variety of metabolic disorders are 1,748,553 (70%) This subtotal category includes heart, stroke, kidney, diabetes, cancer and Alzheimer’s.

A note from WHO: An estimated 17.9 million people died from CVDs in 2019, representing 32% of all global deaths. Of these deaths, 85% were due to heart attack and stroke.”

The main path of many diseases and medical conditions is due to “poor lifestyle leading to body weight problems such as being overweight and having obesity, then developing diabetes, hypertension, and hyperlipidemia. This further lead to a variety of metabolic disorder-induced complications, such as heart issues, stroke, kidney problems, even cancers, and Alzheimer’s; therefore, finally leading to death or shortening a person’s lifespan/longevity. *That is why the author has decided to use the phrase, “health and longevity originate from body weight control” as the title. This article particularly focuses on one of the main paths bridges of linking body weight and glucoses.*

Our body weight is easily measured by standing on a weight scale. Body weight (BW) in the early morning is highly correlated with a diabetes patient’s fasting plasma glucose (FPG) value measured first thing in the morning. In the author’s case, the correlation between his BW and FPG over the past 7+ years is higher than 78% (Figure 1). He has also used his BW as the main influential variable for predicting his FPG value in the early morning (there are other 4 secondary influential factors of FPG, i.e. sleep quality, stress situation, weather/room temperature, and sickness) which has achieved an extremely high prediction accuracy of 99.94% with another high correlation of 99.8% (Figure 1). The statistical algorithm for the predicted FPG value is explained in detail in the section of Methods.

Furthermore, FPG also serves as the baseline of postprandial plasma glucose (PPG). Although PPG has about 19 known contribution factors, ***FPG alone contributes 90-95% of the measured PPG value if glucoses are under good control, and***

still contributes 70-75% of the measured PPG value if glucoses are not under good control. In addition, FPG contributes around 25-29% and PPG contributes about 71-75% of the daily estimated average glucose (eAG) via either a finger-pierced test strip measurement or a continuous glucose monitoring (CGM) sensor device. There is definitely a quasi-linear and highly correlated relationship existing between the eAG values and the quarterly laboratory-measured HbA1C values (see Figure 1). The HbA1C values indicate the average eAG level over the past 90-115 days. Clinical doctors usually utilize the HbA1C value to determine the severity of diabetes disease in their patients. **For the author's case, he has used a linear conversion factor of 17.2 to calculate his HbA1C from eAG. Using this linear conversion factor, he has reached a prediction accuracy of 97.6% (lab-tested HbA1C of 6.56 versus predicted HbA1C of 6.70 during 7+ years from 6/1/2015 to 9/30/2022).**

Finally, diabetes is one of the major influential chronic diseases which would develop into many different deadly complications, including heart diseases, such as CVD, stroke, kidney disease, and even cancers.

There are two case studies in this research article regarding the output symptom of PPG versus three input causes of FPG, added PPG from the diet (food), and reduced PPG from post-meal walking exercise (steps):

(1) Real Case: He has measured 8 PPG annual data as the output strain, and corresponding FPG, food, and steps as three input causes (viscosities) from 6/1/2015 to 9/30/2022. Since this 1 symptom of PPG and 3 causes of FPG, food, and steps have the same unit of mg/dL; therefore the normalization factors are 1.0. In this Real Case, due to his stringent lifestyle management efforts (not through medications), his average values are BW = 171 lbs. or BMI = 24.9 (100%), PPG = 115 mg/dL (100%), FPG = 109 mg/dL (100%), food addition = 11 mg/dL (100%), steps reduction = -12 mg/dL (100%), and HbA1C = 6.6%. In other words, he has been quite "healthy" from the viewpoint of diabetic control.

(2) Hypothetical Case: He has assumed to use 110% of BW, 300% of carbs/sugar grams each meal, and 50% of post-meal walking steps from the Real Case during the same period from 6/1/2015 to 9/30/2022 for this hypothetical case. Since the 1 symptom of PPG and 3 causes of FPG, food, and steps have the same unit of mg/dL; therefore, the normalization factors are 1.0 as well. In this Hypothetical Case, his average values are BW = 189 lbs. (110%) or BMI = 27.5, PPG = 157 mg/dL (136%), FPG = 120 mg/dL (110%), food addition = 33 mg/dL (300%), steps reduction = -6 mg/dL (50%), and HbA1C = 8.6%. In other words, he has been quite "unhealthy" from viewpoint of diabetic control.

(3) Note: The differences between the Real Case and the Hypothetical Case originated from the BW difference of 17 lbs. (BMI difference of 2.6). This 10% weight gain along with differences in both food carbs/sugar addition and post-meal walking steps reduction would finally result in a 30% gain in his HbA1C (from 6.6% to 8.6%). This PPG gain from carbs/sugar intake grams (food) and the PPG reduction from post-meal walking exercise (steps) have been calculated using the

reciprocal of Young's modulus in linear elasticity theory of the discipline of engineering material.

This article's data preparation and data processing work is 100% dependent on the author's recently developed and enhanced VGT software tool which has reduced his data work time from 5-6 hours to less than 1 minute. Therefore, he can spend this saved time on results investigation and biophysical phenomenon interpretation.

The specific program enhancements for this particular study have included **the following 4 newly defined variables for fast Fourier transformed (FFT) frequency-domain (FD) analysis results:**

- (1) strain * stress (stress = strain rate * normalized viscosity)**
- (2) strain * normalized viscosity change rate**
- (3) strain * squared normalized viscosity change rate**
- (4) normalized viscosity change rate**

In summary, these 2-case studies have clearly demonstrated the energy-shift pattern among FPG, Food, and Steps. Using the average values of 6 energy percentages, the most prominent finding of the **Real case** is that **FPG has made the most contribution or influence on PPG, 90%. Food contributes or influences 5% and Steps contributes or influences the remaining 5%.**

However, using the average values of 6 energy percentages, the most noticeable finding from the Hypothetical case is that, **FPG has still made the most contribution or highest influence on PPG, 73%. But, Food contributes or influences 23%, and Steps only contributes or influences the remaining 1%.** In other words, the over-consumption of carbs/sugar is going to take away the energy contribution % from both FPG and Steps. FPG's contribution has been reduced by 17% and Steps contribution has been reduced by 4%, while Food contribution has gained 20%. **This clear picture of energy shifting illustrates the importance of the roles of food and steps on HbA1C, regardless of FPG being also raised by 10% from the BW gain of 10%.**

In addition, this article has also demonstrated the high prediction accuracy of FPG using weight as the primary input. This is an important first step toward overall health and longevity concerns.

Methods

The Author's Case of Diabetes and Complications

The author has been a severe T2D patient since 1996. He weighed 220 lb. (100 kg, BMI 32.5) at that time with a one-time glucose reading of 380 mg/dL. By 2010, he still weighed 198 lb. (BMI 29.2) with average daily glucose of 250 mg/dL (HbA1C of 10%). During that year, his triglycerides reached 1161b (hyperlipidemia) and albumin-creatinine ratio (ACR) at 116 (kidney issues). He also suffered from five cardiac episodes within a decade from 1993 through 2003 caused by work stress and diabetes. In 2010, three independent physicians warned him about his urgent need for kidney dialysis treatment and the risk of his life-threatening health situation such as dying from his severe diabetic complications. Other than the cerebrovascular

disease (stroke), he has suffered most of the known diabetic complications, including both macro-vascular & micro-vascular complications, nerve damage as in retinopathy and foot ulcer, as well as a hormonal disturbance, e.g. hypothyroidism.

In 2010, he decided to launch his self-study on endocrinology, diabetes, and food nutrition to save his own life. After developing the metabolism model in 2024, during 2015 and 2016, he developed four prediction models related to diabetes conditions: weight, PPG, fasting plasma glucose (FPG), and A1C. As a result, from using his developed mathematical metabolism index (MI) model in 2014 and those 4 prediction tools, by end of 2016, his weight was reduced from 220 lbs. (100 kg, BMI 32.5) to 176 lbs. (89 kg, BMI 26.0), waistline from 44 inches (112 cm) to 33 inches (84 cm), average finger glucose reading from 250 mg/dL to 120 mg/dL, and lab-tested A1C from 10% to ~6.5%. ***One of his major accomplishments is that he no longer takes any diabetes medications as of 12/8/2015.***

Around that time (2014–2017), he started to focus on preventive medicine instead of blindly trusting and depending on medication treatments only. He also gambled on his belief that most human organs have strong inherent abilities to self-repair themselves through lifestyle improvements by taking good care of them - even though it can only accomplish a certain degree of repairing or healing dependent on certain organ cells and their status of damage, such as pancreatic beta cells.

In 2017, he has achieved excellent results on all fronts, especially glucose control. However, during the pre-COVID period of 2018 and 2019, he traveled to approximately 50+ international cities to attend 65+ medical conferences and made ~120 oral presentations. This hectic schedule inflicted damage to his diabetes control, through dining out frequently, post-meal exercise disruption, jet lag, and along with the overall metabolic impact due to his irregular life patterns through a busy travel schedule; therefore, his glucose control and overall metabolism state were somewhat affected during this two-year heavy traveling period.

Since 1/19/2020, living in a COVID-19 quarantined lifestyle, not only has he written and published ~500 medical papers in 100+ journals, but he has also reached his best health conditions in the past 26 years. By the beginning of 2022, his weight was further reduced to ***168 lbs. (BMI 24.8) along with a 5.8% A1C value (beginning level of pre-diabetes)***, without having any medication interventions or insulin injections. During the period from 1/1/2022 to 8/20/2022, his average FPG is 93 mg/dL, PPG is 113 mg/dL, and daily glucose is 106 mg/dL. These good results are due to his non-traveling, low-stress, and regular daily life routines. Of course, the accumulated knowledge of chronic diseases, various complications, practical lifestyle management experiences, and development of many high-tech tools along with his medical research academic findings have contributed to his excellent health status since 1/19/2020, the beginning date of his self-quarantined life.

On 5/5/2018, he applied a continuous glucose monitoring (CGM) sensor device on his upper arm and checks his glucose

measurements every 5 minutes for a total of ~288 times each day. He has maintained the same measurement pattern to the present day. In his research work, he uses his CGM sensor glucose at a time interval of 15 minutes (96 data per day). Incidentally, ***the average sensor glucoses between 5-minute intervals and 15-minute intervals has only a 0.6% difference (average glucose of 111.86 mg/dL for 5 minutes and average glucose of 111.18 mg/dL for 15 minutes with a correlation of 94% between these two sensor glucose curves) during the period from 2/19/20 to 7/22/22.***

Therefore, over the past 13 years, he could study and analyze his collected 3+ million data regarding his health status, medical conditions, and lifestyle details. He applies his knowledge, models, and tools from mathematics, physics, engineering, and computer science to conduct his medical research work. *His research work has a goal of achieving both “high precision” and “quantitative proof” in the medical findings for the ultimate objectives of “preventive medicine”.*

The following timetable provides a rough sketch of the emphasis in his medical research during each stage:

- 2000-2013: Self-study diabetes and food nutrition, developing a data collection and analysis software.
- 2014: Develop a mathematical model of metabolism, using engineering modeling and advanced mathematics.
- 2015: Weight & FPG prediction models, using neuroscience.
- 2016: PPG & HbA1C prediction models, using optical physics, artificial intelligence (AI), and neuroscience.
- 2017: Complications due to macro-vascular research, such as Cardiovascular disease (CVD), coronary heart diseases (CHD), and stroke, using pattern analysis and segmentation analysis.
- 2018: Complications due to micro-vascular research such as kidney (CKD), bladder, foot, and eye issues (DR).
- 2019: CGM big data analysis, using wave theory, energy theory, frequency domain analysis, quantum mechanics, and AI.
- 2020: Cancer, dementia, longevity, geriatrics, DR, hypothyroidism, diabetic foot, diabetic fungal infection, and linkage between metabolism and immunity, learning about certain infectious diseases, such as COVID-19.
- 2021: Applications of linear elastic glucose theory (LEGT) and perturbation theory from quantum mechanics on medical research subjects, such as chronic diseases and their complications, cancer, and dementia.
- 2022: Applications of viscoelastic/viscoplastic glucose theory (LEGT) on 142 biomedical research cases and 5 economics research cases.

Again, to date, he has spent ~40,000 hours self-studying and researching medicine and he has read 4,000+ published medical papers online. He has collected and calculated more than 3+ million pieces of data regarding his own medical conditions and lifestyle details. In addition, he has written and published 700+ medical research papers in 100+ various medicine, physics, mathematics, and engineering journals. Moreover, he has also given 120+ presentations at 70+ international medical conferences. He has continuously dedicated his time (11-12 hours per day and work each day of a year, without rest during

the past 13 years) and efforts to his medical research work and shared his findings and learnings with other patients worldwide. In addition, he has also spent the past 12 years developing and maintaining a medicine and health software APP on his iPhone which functions as his private numerical laboratory to process the various experimental datasets of his medical conditions and lifestyle details.

Brief Introduction of Math-Physical Medicine (MPM) Research

The author has collected 3+ million data regarding his health condition and lifestyle details over the past 13 years. He spent the entire year of 2014 developing a metabolism index (MI) model using a topology concept, nonlinear algebra, algebraic geometry, and finite element method. This MI model contains various measured biomarkers and recorded lifestyle details along with their induced new biomedical variables for an additional ~1.5 million data. Detailed data of his body weight, glucose, blood pressure, heart rate, blood lipids, body temperature, and blood oxygen level, along with important lifestyle details, including diet, exercise, sleep, stress, water intake, and daily life routines are included in the MI database. In addition, these lifestyle details also include some lifetime bad habits and certain environmental exposures. Fortunately, the author has none of these lifetime bad habits and an extremely low degree of exposure to environmental factors. The developed MI model has a total of 10 categories covering approximately 500 detailed elements that constitute his defined “metabolism index model” which are the building blocks or root causes for diabetes and other chronic disease-induced complications, including but not limited to cardiovascular disease (CVD), chronic heart disease (CHD), stroke, chronic kidney disease (CKD), diabetic retinopathy (DR), neuropathy, foot ulcer, hypothyroidism, dementia, and various cancers. The end result of the MI development work is a combined MI value within any selected period with 73.5% as its dividing line between a healthy and unhealthy state. The MI serves as the foundation for many of his follow-up medical research work.

During the period from 2015 to 2017, he focused his research on type 2 diabetes (T2D), especially glucose, including fasting plasma glucose (FPG), PPG, estimated average glucose (eAG), and hemoglobin A1C (HbA1C). During the following period from 2018 to 2022, he concentrated on researching medical complications resulting from diabetes, chronic diseases, and metabolic disorders which include heart problems, stroke, kidney problems, retinopathy, neuropathy, foot ulcer, diabetic skin fungal infection, hypothyroidism, diabetic constipation, dementia, and various cancers. He also developed a few mathematical risk models to calculate the probability percentages of developing various diabetic complications based on this MI model. From his previous medical research work with 700+ published papers, he has identified and learned that ***the associated energy of hyperglycemic conditions is the primary source of causing many diabetic complications which lead to death.*** Therefore, a thorough knowledge of these energies is important for achieving a better understanding of the dangerous complications.

TD, SD, and FD Analysis Tools

This section has brief descriptions of TD correlation analysis with other observational results, SD VGT analysis with hysteresis loop area's energy results, and FD analysis with frequency curve area's energy results.

First of all, by using a TD analysis tool, we can examine the curves' moving trend and pattern visually along with their correlation numerically. We can also study the extremely high or low data values in the dataset. The visual observation or calculation-derived interpretations are a part of statistical analysis results which can indeed provide some useful hints or even derive some accurate conclusions. However, we must be aware of the limitations of the selected data size and time window and also be cautious of the appropriate statistics tool we choose.

Regarding the TD energy, we can apply the rudimentary definition of physics that ***“the wave carried energy is directly proportional to the square of wave's amplitude”***. However, the ***data quantity %*** of each wave category should be considered and included to obtain a more accurate TD energy value.

The author would like to describe the essence of his developed “hybrid model” that combines both the SD viscoelastic/plastic VGT analysis method and FD FFT analysis method with a comparison against the traditional TD statistical correlation analysis.

It is described in 10 steps in the English language instead of using mathematical equations to explain it. In this article, he has applied both the SD-VGT operations (steps 1-7) and the FD-FFT operations (steps 8-10). As a result, it is aimed at readers who do not have an extensive background in the academic subjects of engineering, physics & mathematics.

The first step is to collect the output data or symptom (strain or ϵ) on a time scale. The second step is to calculate ***the output change rate with time ($d\epsilon/dt$)***, i.e. the change rate of strain or symptom over each period. The third step is to gather the input data or cause (***viscosity or η***) on a time scale. The fourth step is to calculate the time-dependent input or cause (***time-dependent stress or σ***) by multiplying $d\epsilon/dt$ and η together. The “time-dependent input or cause equation” of ***“stress $\sigma = \text{strain change rate of } d\epsilon/dt * \text{viscosity } \eta$ ”*** is the essential part of this “time dependency”. The fifth step is to plot the input-output (i.e. stress-strain or cause-symptom) curve in a two-dimensional space-domain or SD (x-axis versus y-axis) with strain (output or symptom) on the x-axis and stresses (time-dependent inputs, causes, or stresses) on the y-axis.

The sixth step is to calculate the total enclosed area within these stress-strain curves or input-output curves (i.e. the hysteresis loops), which is also an indicator of associated energies (either created energy or dissipated energy) of this input and output dataset. These energy values can also be considered as the degrees of influence on output by inputs. The seventh step

is the assembly of the area values of the selected periods to compare the “historical progression and contribution of medical condition” over certain periods.

For the frequency domain, the eighth step is to define a “hybrid input variable” by using “*strain*stress*” which yields another accurate estimation of energy ratio similar to the SD-VGT energy ratio associated with the hysteresis loop. The ninth step is to present these hybrid models’ results of (*strain*stress*) in TD and then perform the FFT operation to convert them into FD. The enclosed area of the frequency curve (where the x-axis is the frequency and the y-axis is the amplitude of energy) can be used to estimate the total FD-FFT energy. The tenth step is to compare these FD energy results against the SD-VGT energy results, or even TD energy results.

After providing the above 10-step description, the author would still like to use the following set of VGT stress-strain mathematical equations in a two-dimensional SD to address the selected medical variables:

Strain

= ε (*time-dependency characteristics of individual strain value at the present time duration*)

Stress

= σ (*based on the change rate of strain multiplying with a chosen viscosity factor η*)

= $\eta * (d\varepsilon/dt)$

= $\eta * (d\text{-strain}/d\text{-time})$

= (*viscosity factor η using individual viscosity factor at present time duration*) * (*strain at present quarter - strain at previous time duration*)

Some of these inputs (causes or viscosity factors) are further normalized by dividing them or being divided by a normalization factor using certain established health standards or medical pieces of knowledge. Some examples of normalization factors are 6.0 for HbA1C, 120 mg/dL for glucose, 25 for body mass index (BMI), 4,000 steps after each meal, 10,000 or 12,000 steps for daily walking exercise depending on time-period selection, 13 grams to 20 grams of carbs/sugar intake amount per meal depends on time-period selection. If using the originally collected data, i.e. the non-normalized data, it would distort the numerical comparison of the hysteresis loop areas. Using this “normalization process”, we can remove the dependency of the individual unit or certain unique characteristics associated

with each viscosity factor. This process allows us to convert the originally collected variables into a set of “*dimensionless variables*” for easier numerical comparison and result interpretation.

Algorithm of Predicted FPG

Using the 90-day data of the primary influential factor, both BW in the morning (X), and FPG in the early morning (Y).

Applying the “linear interpretation method” to fill the data gaps to ensure the comparable data numbers of both X and Y.

Applying the “least mean square formula” to calculate the value of r. And then, calculate avgX, avgY, sumSX (the summation of squared x values), and sumSY (the summation of squared y values).

Calculate the following:

$sdX = \text{sqrt}(\text{sumSX} / \text{number of } x \text{ values})$

$sdY = \text{sqrt}(\text{sumSY} / \text{number of } y \text{ values})$

$b = r * sdY / sdX$

$a = \text{avgY} - b * \text{avgX}$

After finishing the above procedures, then take today’s morning weight value (if not available, then use the average morning weight over the past 90 days) to calculate:

Predicted FPG = $a + b * \text{weight}$

Regarding the other 4 remaining secondary influential factors of the morning FPG:

Modification 1:

using *sleep scores* and the normalization factor in the Personal Profile setting to adjust.

Modification 2:

Using *weather or room temperature* and the normalization factor in the Personal Profile setting to adjust.

Modification 3:

Using *stress scores* and the normalization factor in the Personal Profile setting to adjust.

Modification 4:

Using *sickness scores* and the normalization factor in the Personal Profile setting to adjust.

Figure 3 reflects 2 SD-VGT analysis results for both Real Case and Hypothetical Case.




Figure 3: 2 SD-VGT analysis results for both Real Case and Hypothetical Case

Figure 4 reveals 6 contribution percentages of 6 different energies (including 1 TD, 1 SD, & 4 FD), and % of energy changes between the Real Case and Hypothetical Case.




Figure 4: 6 contribution % of 6 different energies (1 TD, 1 SD, 4 FD), and % of energy changes between Real Case and Hypothetical Case

Conclusions

In summary, these 2-case studies have clearly demonstrated the energy-shift pattern among FPG, Food, and Steps. Using the average values of 6 energy percentages, the most prominent finding of the Real case is that FPG has made the most contribution or influence on PPG, 90%. Food contributes or influences 5% and Steps contributes or influences the remaining 5%.

However, using the average values of 6 energy percentages, the most noticeable finding from the ***Hypothetical case is that, FPG has still made the most contribution or highest influence on PPG, 73%. But, Food contributes or influences 23%, and Steps only contributes or influences the remaining 1%.*** In other words, the over-consumption of carbs/sugar is going to take away the energy contribution % from both FPG and Steps. FPG's contribution has been reduced by 17% and Steps contribution has been reduced by 4%, while Food contribution has gained 20%. ***This clear picture of energy shifting illustrates the importance of the roles of food and steps on HbA1C, regardless of FPG being also raised by 10% from the BW gain of 10%.***

In addition, this article has also demonstrated the high prediction accuracy of FPG using weight as the primary input. This is an important first step toward overall health and longevity concerns.

The author has applied a math-physical medicine approach that uses mathematics (not statistics), physics (a branch of applied math), and engineering (a branch of applied physics that includes both biology and chemistry). The TD energy has used a rudimentary definition of the wave energy from basic physics, SD viscoelasticity or viscoplastic stress-strain hysteresis loop energy has used more sophisticated time-dependent characters of engineering and material science, while the FD energy has used 4 different newly-defined variables by applying wave theory of physics and complex variable's FFT operation of mathematics.

The following paragraph only reflects the author's personal opinion:

The author has a personal belief that everything and every object

or matter under the sun must follow the basic law of Physics (i.e. principles and theorem); otherwise, they will not be able to exist in this world. The biomedical phenomena are not exceptional cases. Using statistics tools to investigate biomedical phenomena is necessary but not sufficient because statistics tools only scratch the surface. Incidentally, statistics application has many shortcomings. For example, by selecting a different dataset, a distinct time window, or missing some important data which carries some important biophysical characteristics, a different or even a wrong conclusion could be drawn. That is why some researchers even tweak the statistical data to create the results they would like to see. Using the academically sound math-physical methodology, the researcher's responsibility is mainly to interpret or explain the research findings and/or the observed biophysical phenomena, but they cannot change those finding results and phenomena. The author does not believe his viewpoint is biased since, after all, science is science, the only difference is the researcher's depth and width of academic knowledge, plus tools availability and suitability for their specific research subject.

References

For editing purposes, the majority of the references in this paper, which are self-references, have been removed. Only references from other authors' published sources remain. The bibliography of the author's original self-references can be viewed at www.eclairmd.com.

Readers may use this article as long as the work is properly cited, their use is educational and not for profit, and the author's original work is not altered.

For reading more of the author's published VGT or FD analysis results on medical applications, please locate them through three published special editions from the following three specific journals:

- (1) Series of Endocrinology, Diabetes and Metabolism (contact: Patrick Robinson).
- (2) Journal of Applied Material Science & Engineering Research (contact: Catherine).
- (3) Advances in Bioengineering and Biomedical Science Research (contact: Sony Hazi).

Copyright: ©2022 Gerald C. Hsu. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.